arXiv:0707.0889v4 [math.QA] 11 Jul 2008

DEFORMATION THEORY OF REPRESENTATIONS OF PROP(ERAD)S

SERGEI MERKULOV, BRUNO VALLETTE

Abstract. We study the deformation theory of morphisms of properads and props thereby extending Quillen's deformation theory for commutative rings to a non-linear framework. The associated chain complex is endowed with an L_∞-algebra structure. Its Maurer-Cartan elements correspond to deformed structures, which allows us to give a geometric interpretation of these results.

To do so, we endow the category of prop(erad)s with a model category structure. We provide a complete study of models for prop(erad)s. A new effective method to make minimal models explicit, that extends the Koszul duality theory, is introduced and the associated notion is called *homotopy Koszul*.

As a corollary, we obtain the (co)homology theories of (al)gebras over a prop(erad) and of homotopy (al)gebras as well. Their underlying chain complex is endowed an L_∞-algebra structure in general and a Lie algebra structure only in the Koszul case. In particular, we make the deformation complex of morphisms from the properad of associative bialgebras explicit . For any minimal model of this properad, the boundary map of this chain complex is shown to be the one defined by Gerstenhaber and Schack. As a corollary, this paper provides a complete proof of the existence of an L_∞-algebra structure on the Gerstenhaber-Schack bicomplex associated to the deformations of associative bialgebras.

Introduction

The theory of props and properads, which generalizes the theory of operads, provides us with a universal language to describe many algebraic, topological and differential geometric structures. Our main purpose in this paper is to establish a new and surprisingly strong link between the theory of prop(erad)s and the theory of L_∞-algebras.

We introduce several families of L_∞-algebras canonically associated with prop(erad)s, moreover, we develop new methods which explicitly compute the associated L_∞-brackets in terms of prop(erad)ic compositions and differentials. Many important dg Lie algebras in algebra and geometry (such as Hochschild, Poisson, Schouten, Frölicher-Nijenhuis and many others) are proven to be of this prop(erad)ic origin.

The L_∞-algebras we construct in our paper out of dg prop(erad)s encode many important properties of the latter. The most important one controls the deformation theory of morphisms of prop(erad)s and, in particular, the deformation theory of (al)gebras over prop(erad)s. Applications of our results to the deformation theory of many algebraic and geometric structures becomes therefore another major theme of our paper.

On the technical side, the story develops (roughly speaking) as follows: first we associate canonically to a pair, $(\mathcal{F}(V), \partial)$ and $(\mathcal{Q}, d)$, consisting of a differential graded (dg, for short) quasi-free prop(erad) $\mathcal{F}(V)$ on a $\mathbb{S}$-bimodule V and an arbitrary dg prop(erad) $\mathcal{Q}$, a structure of L_∞-algebra on the (shifted) graded vector space, $s^{-1}\mathrm{Hom}^{\mathbb{S}}_\bullet(V, \mathcal{Q})$, of morphisms of $\mathbb{S}$-bimodules; then we prove the Maurer-Cartan elements of this L_∞-algebra are in *one-to-one* correspondence with the set of all dg morphisms,

$$\{(\mathcal{F}(V), \partial) \longrightarrow (\mathcal{Q}, d)\},$$

of dg prop(erad)s. This canonical L_∞-algebra is used then to define, for any particular morphism $\gamma : (\mathcal{F}(V), \partial) \to (\mathcal{Q}, d)$, another twisted L_∞-algebra which controls deformation theory of the morphism γ. In the special case when $(\mathcal{Q}, d)$ is the endomorphism prop(erad), (End_X, d_X), of some dg vector space X, our theory gives L_∞-algebras which control deformation theory of many classical algebraic and geometric structures on X, for example, associative algebra structure, Lie algebra structure, commutative algebra structure, Lie bialgebra structure, associative bialgebra structure, formal Poisson structure, Nijenhuis structure etc. As the case of associative bialgebras has never been rigorously treated in the literature before, we discuss this example in full details; we prove, in particular, that the first term of the canonical L_∞-structure controlling deformation theory of bialgebras is precisely the Gerstenhaber-Schack differential.

We derive and study the deformation complex and its L_∞-structure from several different perspectives. One of them can be viewed as a nontrivial generalization to the case of prop(erad)s of Van der Laan's approach [VdL02] to the deformation theory of algebras over operads, while others are completely new and provide us with, perhaps, a conceptual explanation of the observed (long ago) phenomenon that deformation theories are controlled by dg Lie and, more generally, L_∞ structures.

First, we define the deformation complex of a morphism of prop(erad)s $\mathcal{P} \to \mathcal{Q}$ generalizing Quillen's definition of the deformation complex of a morphism of commutative rings. When $(\mathcal{F}(\mathcal{C}), \partial)$ is a quasi-free resolution of $\mathcal{P}$, we prove that this chain complex is isomorphic, up to a shift of degree, to the space of morphisms of $\mathbb{S}$-bimodule $\mathrm{Hom}^{\mathbb{S}}_{\bullet}(\mathcal{C}, \mathcal{Q})$, where $\mathcal{C}(\simeq s^{-1}V)$ is a homotopy coprop(erad), that is the dual notion of prop(erad) with relations up to homotopy. Since $\mathcal{Q}$ is a (strict) prop(erad), we prove that the space $\mathrm{Hom}^{\mathbb{S}}_{\bullet}(\mathcal{C}, \mathcal{Q})$ has a rich algebraic structure, namely it is a homotopy non-symmetric prop(erad), that is a prop(erad) without the action of the symmetric groups and with relations up to homotopy. From this structure, we extract a canonical L_∞-structure on $\mathrm{Hom}^{\mathbb{S}}_{\bullet}(\mathcal{C}, \mathcal{Q}) \simeq s^{-1}\mathrm{Hom}^{\mathbb{S}}_{\bullet}(V, \mathcal{Q})$. We also obtain higher operations with $m+n$ inputs acting on $\mathrm{Hom}^{\mathbb{S}}_{\bullet}(\mathcal{C}, \mathcal{Q})$ which are important in applications. In the case of, for example, the non-symmetric operad, $\mathcal{A}ss$, of associative algebras the deformation complex is the Hochschild cochain complex of an associative algebra, and the higher homotopy operations are shown to be non-symmetric braces operations which play a fundamental role in the proof of Deligne's conjecture (see [Tam98, Vor00, KS00, MS02, BF04, Kau07]).

Recall that M. Markl proved in [Mar04b] a first interesting partial result, that is for a given minimal model $(\mathcal{F}(\mathcal{C}), \partial)$ a prop(erad) $\mathcal{P}$ concentrated in degree 0, there exists a L_∞-structure on the space of derivations from $\mathcal{F}(\mathcal{C})$ to End_X, where X is a $\mathcal{P}$-(al)gebra. By using a different conceptual method based on the notions of homotopy (co)prop(erad)s and convolution prop(erad)s, we prove here that for any representation $\mathcal{Q}$ of any prop(erad) $\mathcal{P}$, there exists a homotopy prop(erad) structure on the space of derivations from any quasi-free resolution of $\mathcal{P}$ to $\mathcal{Q}$. Moreover this construction is shown to be functorial, that is does not depend on the model chosen. From this we derive functorially the general L_∞-structure.

Another approach of deriving the deformation complex and its L_∞-structure is based on a canonical enlargement of the category of dg prop(erad)s via an extension of the notion of *morphism*; the set of morphisms, $\mathrm{Mor}_{\mathbb{Z}}(\mathcal{P}_1, \mathcal{P}_2)$, in this enlarged category is identified with a certain *dg affine scheme* naturally associated with both $\mathcal{P}_1$ and $\mathcal{P}_2$; moreover, when the dg prop(erad) $\mathcal{P}_1$ is quasi-free, the dg affine scheme $\mathrm{Mor}(\mathcal{P}_1, \mathcal{P}_2)$ is proven to be a *smooth* dg manifold for any $\mathcal{P}_2$ and hence gives canonically rise to a L_∞-structure.

The third major theme of our work is the theory of models and minimal models. To make explicit the deformation complex, we need models, that is quasi-free resolutions of prop(erad)s. We extend the bar and cobar construction to prop(erad)s and show that the bar-cobar construction provides a canonical cofibrant resolution of a prop(erad). Since this construction is not very convenient to work with because it is too big, we would rather use minimal models. We give a complete account to the theory of minimal models for prop(erad)s. We prove that minimal models for prop(erad)s are not in general determined by resolutions of their genus 0 parts, namely dioperads, giving thereby a negative answer to a question raised by M. Markl and A.A. Voronov [MV03], that is we prove that the free functor from dioperads to prop(erad)s is not exact. We provide an explicit example of a Koszul dioperad which does not induce the prop(erad)ic resolution of the associated prop(erad).

A properad is Koszul if and only if it admits a quadratic model. In this case, Koszul duality theory of properad [Val07b] provides an effective method to compute this special minimal model. Unfortunately, not all properads are Koszul. For instance, the properad coding associative bialgebras is not. We include this example in a new notion, called *homotopy Koszul*. A homotopy Koszul properad is shown to have a minimal model that can be explicitly computed. Its space of generators is equal to the Koszul dual of a quadratic properad associated to it. And the differential is made explicit by use of the (dual) formulae of J. Granåker [Gra07] of transfer of homotopy coproperad structure, that is by perturbing the differential. We apply this notion to show that morphisms of homotopy $\mathcal{P}$-algebras are in one-to-one correspondence with Maurer-Cartan elements of a convolution L_∞-algebra.

In the appendix, we endow the category of dg prop(erad)s with a model category structure which is used throughout the text.

The paper is organized as follows. In §1 we remind key facts about properads and props and we define the notion of *non-symmetric prop(erad)*. In §2 we introduce and study the convolution prop(erad) canonically associated with a pair, $(\mathcal{C}, \mathcal{P})$, consisting of an arbitrary coprop(erad) $\mathcal{C}$ and an arbitrary prop(erad) $\mathcal{P}$; our main results is the construction of a Lie algebra structure on this convolution properad, as well as higher operations. In §3 we discuss bar and cobar constructions for (co)prop(erad)s. We introduce the notion of *twisting morphism (cochain)* for prop(erads) and prove Theorem 19 on bar-cobar resolutions extending thereby earlier results of [Val07a] from weight-graded dg properads to arbitrary dg properads. In §4 we recall to the notion and properties of homotopy properads which were first introduced in [Gra07] and we define the notions of *homotopy (co)prop(erad)*. We apply these notions to convolution prop(erad)s. In §5, we give a complete study of minimal models for properads. In §6 we remind geometric interpretation of L_∞-algebras, and then use this geometric language to prove Theorem 62 which associates to pair, $(\mathcal{F}(V), \partial)$ and $(\mathcal{P}, d)$, consisting of quasi-free prop(erad) $(\mathcal{F}(V), \partial)$ and an arbitrary dg prop(erad) $(\mathcal{P}, d)$, a structure of L_∞-algebra on the (shifted) graded vector space, $s^{-1}\mathrm{Hom}_\bullet^{\mathbb{S}}(V, \mathcal{P})$; we also show in §6 full details behind the construction of the above mentioned enlargement of the category of dg prop(erad)s. In §7, we define the deformation complex following Quillen's methods and identify it with $s^{-1}\mathrm{Hom}_\bullet^{\mathbb{S}}(V, \mathcal{P})$ in Theorem 69. We show next how this canonical L_∞-algebra gives rise to twisted L_∞-algebras which control deformation theories of particular morphisms $\gamma : (\mathcal{F}(V), \partial) \to (\mathcal{Q}, d)$. In §8, we illustrate this general construction with several examples from algebra and geometry. We make explicit the deformation complex of representation of the properad of associative bialgebras and prove that it corresponds with the one defined by Gerstenhaber-Schack. In the Appendix, we show that the category of dg prop(erad)s is a cofibrantly generated model category.

Contents

Introduction 1
1. (Co)Properads, (Co)Props and their non-symmetric versions 5
1.1. $\mathbb{S}$-bimodules, graphs, composition products 5
1.2. Properad 6
1.3. Connected coproperad 7
1.4. Free properad and cofree connected coproperad 7
1.5. Props 7
1.6. (Co)triple interpretation 8
1.7. Non-symmetric prop(erad) 8
1.8. Representations of prop(erad)s, gebras 9
2. Convolution prop(erad) 9

2.1. Convolution prop(erad) 9
2.2. Lie-admissible products and Lie brackets associated to a properad 10
2.3. LR-algebra associated to a properad 12
2.4. Lie-admissible bracket and LR-algebra of a convolution properad 13
3. Bar and cobar construction 14
3.1. Infinitesimal bimodule over a prop(erad) 14
3.2. (Co)Derivations 16
3.3. (De)Suspension 17
3.4. Twisting morphism 18
3.5. Bar construction 18
3.6. Cobar construction 19
3.7. Bar-Cobar Adjunction 19
3.8. Props vs properads 20
3.9. Bar-cobar resolution 20
4. Homotopy (Co)prop(erad)s 21
4.1. Definitions 21
4.2. Admissible subgraph 22
4.3. Interpretation in terms of graphs 23
4.4. Homotopy non-symmetric (co)prop(erad) 25
4.5. Homotopy properads and associated homotopy Lie algebras 25
4.6. Homotopy convolution prop(erad) 26
4.7. Morphisms of homotopy (co)prop(erad)s 27
5. Models 30
5.1. Minimal models 30
5.2. Form of minimal models 30
5.3. Quadratic models and Koszul duality theory 31
5.4. Homotopy Koszul properads 32
5.5. Models for associative algebras, non-symmetric operads, operads, properads, props 35
5.6. Models generated by genus 0 differentials 36
6. Homotopy $\mathcal{P}$-gebra 41
6.1. $\mathcal{P}$-gebra, $\mathcal{P}_{(n)}$-gebra and homotopy $\mathcal{P}$-gebra 41
6.2. Morphisms of homotopy $\mathcal{P}$-algebras as Maurer-Cartan elements 41
7. L_∞-algebras, dg manifolds, dg affine schemes and morphisms of prop(erad)s 42
7.1. L_∞-algebras, dg manifolds and dg affine schemes. 42
7.2. Another geometric model for a L_∞-structure 44
7.3. Maurer-Cartan elements in a filtered L_∞-algebra 44
7.4. Extended morphisms of dg props as a dg affine scheme. 45
7.5. Proofs via local coordinate computations 49
7.6. Enlarged category of dg prop(erad)s 51
7.7. Families of natural L_∞-structures on $\oplus\mathcal{P}$ 52
8. Deformation theory of morphisms of prop(erad)s 54
8.1. Basic Definition 54
8.2. Deformation theory of representations of prop(erad)s. 55
8.3. Koszul case and cohomology operations 57
8.4. Definition à la Quillen 58
9. Examples of deformation theories. 61
9.1. Associative algebras. 62
9.2. Poisson structures. 63
9.3. Associative bialgebras 64
9.4. Twisted L_∞-algebras and dg prop(erad)s 70
Appendix A. Model category structure for prop(erad)s 73
A.1. Model category structure on $\mathbb{S}$-bimodules 73
A.2. Transfer Theorem 74
A.3. Limits and Colimits of prop(erad)s 75

A.4. Model category structure 76
A.5. Cofibrations and Cofibrant objects 77
References 80

In this paper, we will always work over a field $\mathbb{K}$ of characteristic 0.

1. (Co)Properads, (Co)Props and their non-symmetric versions

In this section, we recall briefly the definitions of (co)properad and (co)prop. For the reader who does not know what a properad or what a prop is, we strongly advise to read the first sections of [Val07a] before reading the current article since we will make use of the notions everywhere in the sequel. Generalizing the notion of non-symmetric operads to prop(erad), we introduce the notions of *non-symmetric properad* and *non-symmetric prop.*

1.1. **$\mathbb{S}$-bimodules, graphs, composition products.** A *(dg) $\mathbb{S}$-bimodule* is a collection $\mathcal{P} = \{\mathcal{P}(m,n)\}_{m,\,n\in\mathbb{N}}$ of dg modules over the symmetric groups $\mathbb{S}_n$ on the right and $\mathbb{S}_m$ on the left. These two actions are supposed to commute. In the sequel, we will mainly consider *reduced* $\mathbb{S}$-bimodules, that is $\mathbb{S}$-bimodules $\mathcal{P}$ such that $\mathcal{P}(m,n) = 0$ when $n = 0$ or $m = 0$. We use the homological convention, that is the degree of differentials is -1. An $\mathbb{S}$-bimodule $\mathcal{P}$ is *augmented* when it naturally splits as $\mathcal{P} = \overline{\mathcal{P}} \oplus I$ where $I = \{I(m,n\}$ is an $\mathbb{S}$-bimodule with all components $I(m,n)$ vanishing except for $I(1,1)$ which equals $\mathbb{K}$. We denote the module of morphisms of $\mathbb{S}$-bimodules by $\mathrm{Hom}(\mathcal{P},\mathcal{Q})$ and the module of equivariant morphisms, with respect to the action of the symmetric groups, by $\mathrm{Hom}^{\mathbb{S}}(\mathcal{P},\mathcal{Q})$.

A graph is given by two sets, the set V of vertices and the set E of edges, and relations among which say when an edge is attached to one or two vertices (see [GK98] (2.5)). The egdes of the graph considered in the sequel will always be directed by a global flow (*directed graphs*). The edges can be divided into two parts: the ones with one vertex at each end, called *internal edges*, and the ones with just one vertex on one end, called *external edges*. The *genus* of a graph is the first Betti number of the underlying topological space of a graph. A *2-levelled directed graphs* is a directed graph such that the vertices are divided into two parts, the ones belonging to a bottom level and the ones belonging to a top level. In the category of $\mathbb{S}$-bimodule, we define two *composition* products, $\boxtimes$ based on the composition of operations indexing the vertices of a 2-levelled directed graphs, and $\boxtimes_c$ based on the composition of operations indexing the vertices of a 2-levelled directed connected graph (see Figure 1 for an example). Let $\mathcal{G}$ be such a graph with N internal edges between vertices of the two levels. This set of edges between vertices of the first level and vertices of the second level induces a permutation of $\mathbb{S}_N$.



Figure 1. Example of a 2-level graph.

Let $\mathcal{P}$ and $\mathcal{Q}$ be two $\mathbb{S}$-bimodules, their composition product is given by the explicit formula

$$\mathcal{P} \boxtimes \mathcal{Q}(m,\, n) := \bigoplus_{N \in \mathbb{N}^*} \left(\bigoplus_{\bar{l},\, \bar{k},\, \bar{\jmath},\, \bar{\imath}} \mathbb{K}[\mathbb{S}_m] \otimes_{\mathbb{S}_{\bar{l}}} \mathcal{P}(\bar{l},\, \bar{k}) \otimes_{\mathbb{S}_{\bar{k}}} \mathbb{K}[\mathbb{S}_N] \otimes_{\mathbb{S}_{\bar{\jmath}}} \mathcal{Q}(\bar{\jmath},\, \bar{\imath}) \otimes_{\mathbb{S}_{\bar{\imath}}} \mathbb{K}[\mathbb{S}_n] \right)_{\mathbb{S}_b^{\mathrm{op}} \times \mathbb{S}_a},$$

where the second direct sum runs over the b-tuples $\bar{l}$, $\bar{k}$ and the a-tuples $\bar{\jmath}$, $\bar{\imath}$ such that $|\bar{l}| = m$, $|\bar{k}| = |\bar{\jmath}| = N$, $|\bar{\imath}| = n$ and where the coinvariants correspond to the following action of $\mathbb{S}_b^{\mathrm{op}} \times \mathbb{S}_a$:

$$\begin{aligned} &\theta \otimes p_1 \otimes \cdots \otimes p_b \otimes \sigma \otimes q_1 \otimes \cdots \otimes q_a \otimes \omega \sim \\ &\theta\, \tau_{\bar{l}}^{-1} \otimes p_{\tau^{-1}(1)} \otimes \cdots \otimes p_{\tau^{-1}(b)} \otimes \tau_{\bar{k}}\, \sigma\, \nu_{\bar{\jmath}} \otimes q_{\nu(1)} \otimes \cdots \otimes q_{\nu(a)} \otimes \nu_{\bar{\imath}}^{-1}\, \omega, \end{aligned}$$

for $\theta \in \mathbb{S}_m$, $\omega \in \mathbb{S}_n$, $\sigma \in \mathbb{S}_N$ and for $\tau \in \mathbb{S}_b$ with $\tau_{\bar{k}}$ the corresponding block permutation, $\nu \in \mathbb{S}_a$ and $\nu_{\bar{\jmath}}$ the corresponding block permutation. This product is associative but has no unit. To fix this issue, we restrict to connected graphs.
The permutations of $\mathbb{S}_N$ associated to connected graphs are called *connected* (for more details see Section 1.3 of [Val07a]). We denote the set of connected permutations by $\mathbb{S}^c$. We define the *connected composition product* by the following formula

$$\mathcal{P} \boxtimes_c \mathcal{Q}(m,\, n) := \bigoplus_{N \in \mathbb{N}^*} \left(\bigoplus_{\bar{l},\, \bar{k},\, \bar{\jmath},\, \bar{\imath}} \mathbb{K}[\mathbb{S}_m] \otimes_{\mathbb{S}_{\bar{l}}} \mathcal{P}(\bar{l},\, \bar{k}) \otimes_{\mathbb{S}_{\bar{k}}} \mathbb{K}[\mathbb{S}^c_{\bar{k},\, \bar{\jmath}}] \otimes_{\mathbb{S}_{\bar{\jmath}}} \mathcal{Q}(\bar{\jmath},\, \bar{\imath}) \otimes_{\mathbb{S}_{\bar{\imath}}} \mathbb{K}[\mathbb{S}_n] \right)_{\mathbb{S}_b^{\mathrm{op}} \times \mathbb{S}_a}.$$

The unit I for this monoidal product is given by

$$\begin{cases} I(1,\, 1) := \mathbb{K}, & \text{and} \\ I(m,\, n) := 0 & \text{otherwise.} \end{cases}$$

We denote by $(\mathbb{S}\text{-biMod}, \boxtimes_c,\, I)$ this monoidal category.

We define the *concatenation product* of two bimodules $\mathcal{P}$ and $\mathcal{Q}$ by

$$\mathcal{P} \otimes \mathcal{Q}(m,\, n) := \bigoplus_{\substack{m'+m''=m \\ n'+n''=n}} \mathbb{K}[\mathbb{S}_{m'+m''}] \otimes_{\mathbb{S}_{m'} \times \mathbb{S}_{m''}} \mathcal{P}(m',\, n') \otimes_{\mathbb{K}} \mathcal{Q}(m'',\, n'') \otimes_{\mathbb{S}_{n'} \times \mathbb{S}_{n''}} \mathbb{K}[\mathbb{S}_{n'+n''}].$$

This product corresponds to taking the (horizontal) tensor product of the elements of $\mathcal{P}$ and $\mathcal{Q}$ (see Figure 3 of [Val07a] for an example). It is symmetric, associative and unital. On the contrary to the two previous products, it is linear on the left and on the right. We denote by $\mathcal{S}_\otimes(\mathcal{P})$ the free symmetric monoid generated by an $\mathbb{S}$-bimodule $\mathcal{P}$ for the concatenation product (and $\bar{\mathcal{S}}_\otimes(\mathcal{P})$ its augmentation ideal). There is a natural embedding $\mathcal{P} \boxtimes_c \mathcal{Q} \rightarrowtail \mathcal{P} \boxtimes \mathcal{Q}$. And we obtain the composition product from the connected composition product by concatenation, that is $\bar{\mathcal{S}}_\otimes(\mathcal{P} \boxtimes_c \mathcal{Q}) \cong \mathcal{P} \boxtimes \mathcal{Q}$. (From this relation, we can see that $I \boxtimes \mathcal{P} = \bar{\mathcal{S}}_\otimes(\mathcal{P})$ and not $\mathcal{P}$.)

1.2. **Properad.** A *properad* is a monoid in the monoidal category $(\mathbb{S}\text{-biMod}, \boxtimes_c,\, I)$. We denote the set of morphisms of properads by $\mathrm{Mor}(\mathcal{P}, \mathcal{Q})$. A properad $\mathcal{P}$ is *augmented* if there exists a morphism of properads $\varepsilon : \mathcal{P} \to I$. We denote by $\bar{\mathcal{P}}$ the kernel of the augmentation ε and call it the *augmentation ideal*. When $(\mathcal{P}, \mu, \eta,\, \varepsilon)$ is an augmented properad, $\mathcal{P}$ is canonically isomorphic to $I \oplus \bar{\mathcal{P}}$. We denote by $\big(I \oplus \underbrace{\bar{\mathcal{P}}}_{r}\big) \boxtimes_c \big(I \oplus \underbrace{\bar{\mathcal{P}}}_{s}\big)$ the sub-$\mathbb{S}$-bimodule of $\mathcal{P} \boxtimes_c \mathcal{P}$ generated by compositions of s non-trivial elements of $\mathcal{P}$ on the first level with r non-trivial elements of $\mathcal{P}$ on the second level. The corresponding restriction of the composition product μ on this sub-$\mathbb{S}$-bimodule is denoted $\mu_{(r,\, s)}$. The bilinear part of $\mathcal{P} \boxtimes_c \mathcal{P}$ is the $\mathbb{S}$-bimodule $\big(I \oplus \underbrace{\bar{\mathcal{P}}}_{1}\big) \boxtimes_c \big(I \oplus \underbrace{\bar{\mathcal{P}}}_{1}\big)$. It corresponds to the compositions of only 2 non-trivial operations of $\mathcal{P}$. We denote it by $\mathcal{P} \boxtimes_{(1,1)} \mathcal{P}$. The composition of two elements p_1 and p_2 of $\bar{\mathcal{P}}$ is written $p_1 \boxtimes_{(1,1)} p_2$ to lighten the notations. The restriction $\mu_{(1,\,1)}$ of the composition product μ of a properad $\mathcal{P}$ on $\mathcal{P} \boxtimes_{(1,1)} \mathcal{P}$ is called the *partial product.*
A properad is called *reduced* when the underlying $\mathbb{S}$-bimodule is reduced, that is when $\mathcal{P}(m, n) = 0$ for $n = 0$ or $m = 0$.

1.3. **Connected coproperad.** Dually, we defined the notion of *coproperad*, which is a comonoid in ($\mathbb{S}$-biMod, $\boxtimes_c$). Recall that the partial coproduct $\Delta_{(1,\,1)}$ of a coproperad $\mathcal{C}$ is the projection of the coproduct Δ on $\mathcal{C} \boxtimes_{(1,1)} \mathcal{C} := \big(I \oplus \underbrace{\mathcal{C}}_{1}\big) \boxtimes_c \big(I \oplus \underbrace{\mathcal{C}}_{1}\big)$. More generally, one can define the (r, s)-part of the coproduct by the projection of the image of Δ on $\big(I \oplus \underbrace{\mathcal{C}}_{r}\big) \boxtimes_c \big(I \oplus \underbrace{\mathcal{C}}_{s}\big)$.

Since the dual of the notion of coproduct is the notion of product, we have to be careful with coproperad. For instance, the target space of a morphism of coproperads is a direct sum of modules and not a product. (The same problem appears at the level of algebras and coalgebras). We generalize here the notion of *connected* coalgebra, which is the dual notion of Artin rings, introduced by D. Quillen in [Qui69] Appendix B, Section 3 (see also J.-L. Loday and M. Ronco [LR06] Section 1).
Let $(\mathcal{C},\, \Delta,\, \varepsilon,\, u)$ be an coaugmented (dg) coproperad. Denote by $\overline{\mathcal{C}} := \mathrm{Ker}(\mathcal{C} \xrightarrow{\varepsilon} I)$ its *augmentation*. We have $\mathcal{C} = \overline{\mathcal{C}} \oplus I$ and $\Delta(I) = I \boxtimes_c I$. For $X \in \overline{\mathcal{C}}$, denote by $\overline{\Delta}$ the non-primitive part of the coproduct, that is $\Delta(X) = I \boxtimes_c X + X \boxtimes_c I + \overline{\Delta}(X)$. The *coradical filtration* of $\mathcal{C}$ is defined inductively as follows

$$\begin{aligned} &F_0 := \mathbb{K}\, I \\ &F_r := \{X \in \mathcal{C} \,|\, \overline{\Delta}(X) \in F_{r-1} \boxtimes_c F_{r-1}\}. \end{aligned}$$

An augmented coproperad is *connected* if the coradical filtration is exhaustive $\mathcal{C} = \bigcup_{r \geq 0} F_r$. This condition implies that $\mathcal{C}$ is *conilpotent* which means that for every $X \in \mathcal{C}$, there is an integer n such that $\overline{\Delta}^n(X) = 0$. This assumption is always required to construct morphisms or coderivations between coproperads (see next paragraph and Lemma 15 for instance).

For the same reason, we will sometimes work with the invariant version of the composition product denoted $\mathcal{P} \boxtimes_c^{\mathbb{S}} \mathcal{Q}$ when working with coproperads. It is defined by the same formula as for $\boxtimes_c$ but where we consider the invariant elements under the actions of the symmetric groups instead of the coinvariants (see Lemma 2 for instance). When we want to emphasize the difference between invariants and coinvariants, we use the notations $\boxtimes^{\mathbb{S}}$ and $\boxtimes_{\mathbb{S}}$. Otherwise, we use only $\boxtimes$ since the two are isomorphic in characteristic 0.

1.4. **Free properad and cofree connected coproperad.** Recall from [Val04] the construction of the free properad. Let V be an $\mathbb{S}$-bimodule. Denote by $V^+ := V \oplus I$ its augmentation and by $V_n := (V^+)^{\boxtimes_c n}$ the n-fold "tensor" power of V^+. This last module can be thought of as n-levelled graphs with vertices indexed by V and I. We define on V_n an equivalence relation $\sim$ by identifying two graphs when one is obtained from the other by moving an operation from a level to an upper or lower level. (Note that this permutation of the place of the operations induces signs). We consider the quotient $\widetilde{V}_n := V_n/\sim$ by this relation. Finally, the free properad $\mathcal{F}(V)$ is given by a particular colimit of the $\widetilde{V}_n$. The dg $\mathbb{S}$-bimodule $\mathcal{F}(V)$ is generated by graphs without levels with vertices indexed by elements of V. We denote such graphs by $\mathcal{G}(v_1, \ldots, v_n)$, with $v_1, \ldots, v_n \in V$. Since $\mathcal{G}(v_1, \ldots, v_n)$ represents an equivalence class of levelled graphs, we can chose, up to signs, an order for the vertices. (Any graph $\mathcal{G}$ with n vertices is the quotient by the relation $\sim$ of a graph with n levels and only one non-trivial vertex on each level). The composition product of $\mathcal{F}(V)$ is given by the grafting. It is naturally graded by the number of vertices. This grading is called the *weight*. The part of weight n is denoted by $\mathcal{F}(V)^{(n)}$.

Since we are working over a field of characteristic 0, the cofree connected coproperad on a dg $\mathbb{S}$-bimodule V has the same underlying space as the free properad, that is $\mathcal{F}^c(V) = \mathcal{F}(V)$. The coproduct is given by pruning the graphs into two parts. This coproperad verifies the universal property only among connected coproperads (see Proposition 2.7 of [Val07a])

1.5. **Props.** We would like to define the notion of *prop* as a monoid in the category of $\mathbb{S}$-bimodules with the composition product $\boxtimes$. Since this last one has no unit and is not a monoidal product, strictly speaking, we have to make this definition explicit.

Definition (Prop)**.** A *prop* $\mathcal{P}$ is an $\mathbb{S}$-bimodule endowed with two maps $\mathcal{P} \boxtimes \mathcal{P} \xrightarrow{\mu} \mathcal{P}$ and $I \xrightarrow{\eta} \mathcal{P}$ such that the first is associative and the second one verifies

$$\begin{tikzcd}
I \boxtimes_c \mathcal{P} \arrow[r, tail] \arrow[drr, "\sim"'] & I \boxtimes \mathcal{P} \arrow[r, tail, "\eta \boxtimes \mathcal{P}"] & \mathcal{P} \boxtimes \mathcal{P} \arrow[d, "\mu"] & \mathcal{P} \boxtimes I \arrow[l, tail, "\mathcal{P} \boxtimes \eta"'] & \mathcal{P} \boxtimes_c I \arrow[l, tail] \arrow[dll, "\sim"] \\
 & & \mathcal{P}. & &
\end{tikzcd}$$

This definition is equivalent to the original definition of Adams and MacLane [Ada78, ML65]. The original definition consists of two coherent bilinear products, the vertical and horizontal compositions of operations. The definition of the composition product given here includes these two previous compositions at the same time. The partial product $\mathcal{P} \boxtimes_{(1,1)} \mathcal{P} \xrightarrow{\mu_{(1,1)}} \mathcal{P}$ composes two operations. If they are connected by at least one edge, this composition is the vertical composition, otherwise this composition can be seen as the horizontal composition of operations. This presentation will allow us later to define the bar construction, resolutions and minimal models for props.

It is straightforward to extend the results of the preceding subsections to props. There exists notions of augmented props, free prop, coprop and connected cofree coprop. We refer the reader to [Val07a] Section 2 for a complete treatment.

1.6. **(Co)triple interpretation.** The free prop(erad) functor induces a triple $\mathcal{F} : \mathbb{S}-\mathrm{biMod} \to \mathbb{S}-\mathrm{biMod}$ such that an algebra over it is a prop(erad) (see D. Borisov and Y.I. Manin [BM06]). When $(\mathcal{P}, \mu)$ is a prop(erad), we will denote by $\widetilde{\mu}_{\mathcal{P}} : \mathcal{F}(\mathcal{P})^{(\geq 2)} \to \mathcal{P}$ the associated map. Dually, the notion of coprop(erad) is equivalent to the notion of coalgebra over the cotriple $\mathcal{F}^c : \mathbb{S}-\mathrm{biMod} \to \mathbb{S}-\mathrm{biMod}$. When $(\mathcal{C}, \Delta)$ is a coprop(erad), we will denote by $\widetilde{\Delta}_{\mathcal{C}} : \mathcal{C} \to \mathcal{F}^c(\mathcal{C})^{(\geq 2)}$ the associated map.

1.7. **Non-symmetric prop(erad).** In the sequel, we will have to work with algebraic structures endowed with operations having no symmetries. One can model them with properads but the action of the symmetric group gives no real information. Therefore, we introduction the notion of *non-symmetric properad* which will be enough. Since this structure is the direct generalization of the notion of non-symmetric operad, we call it *non-symmetric properad.* All the definitions and propositions of this section can be generalized directly to props. For simplicity, we only make them explicit in the case of properads.

Definition. A *(dg)* $\mathbb{N}$*-bimodule* is a collection $\{\mathsf{P}(m, n)\}_{m,\, n \in \mathbb{N}^*}$ of dg modules.

Definition (Non-symmetric connected composition product)**.** Let P and Q be two $\mathbb{N}$-bimodules, we define their *non-symmetric connected composition product* by the following formula

$$\mathsf{P} \boxtimes_c \mathsf{Q}(m,\, n) := \bigoplus_{N \in \mathbb{N}^*} \left(\bigoplus_{\bar{l},\, \bar{k},\, \bar{\jmath},\, \bar{\imath}} \mathsf{P}(\bar{l},\, \bar{k}) \otimes \mathbb{K}[\mathbb{S}^c_{\bar{k},\, \bar{\jmath}}] \otimes \mathsf{Q}(\bar{\jmath},\, \bar{\imath}) \right)_{\mathbb{S}_b^{\mathrm{op}} \times \mathbb{S}_a},$$

where the second direct sum runs over the b-tuples $\bar{l}$, $\bar{k}$ and the a-tuples $\bar{\jmath}$, $\bar{\imath}$ such that $|\bar{l}| = m$, $|\bar{k}| = |\bar{\jmath}| = N$, $|\bar{\imath}| = n$ and where the coinvariants correspond to the following action of $\mathbb{S}_b^{\mathrm{op}} \times \mathbb{S}_a$:

$$p_1 \otimes \cdots \otimes p_b \otimes \sigma \otimes q_1 \otimes \cdots \otimes q_a \sim p_{\tau^{-1}(1)} \otimes \cdots \otimes p_{\tau^{-1}(b)} \otimes \tau_{\bar{k}}\, \sigma\, \nu_{\bar{\jmath}} \otimes q_{\nu(1)} \otimes \cdots \otimes q_{\nu(a)},$$

for $\sigma \in \mathbb{S}^c_{\bar{k},\, \bar{\jmath}}$ and for $\tau \in \mathbb{S}_b$ with $\tau_{\bar{k}}$ the corresponding block permutation, $\nu \in \mathbb{S}_a$ and $\nu_{\bar{\jmath}}$ the corresponding block permutation. Since the context is obvious, we still denote it by $\boxtimes_c$.

The definition of the monoidal product for $\mathbb{S}$-bimodule is based on 2-levelled graphs with leaves, inputs and outputs labelled by integers. This definition is based on non-labelled 2-levelled graphs. We define the *non-symmetric composition product* $\boxtimes$ by the same formula with the set of all permutations of $\mathbb{S}_N$ instead of connected permutations.

Proposition 1. *The category* $(\mathbb{N}\text{-}biMod, \boxtimes_c, I)$ *of* $\mathbb{N}$*-bimodules with the product* $\boxtimes_c$ *and the unit* I *is a monoidal category.*

Proof. The proof is similar to the one for $\mathbb{S}$-bimodules (see [Val07a] Proposition 1.6). $\square$

Definition (Non-symmetric properad)**.** A *non-symmetric properad* (P, μ, η) is a monoid in the monoidal category $(\mathbb{N}\text{-biMod}, \boxtimes_c, I)$.

Example. A non-symmetric properad P concentrated in arity $(1, n)$, with $n \geq 1$, is the same as a non-symmetric operad.

1.8. **Representations of prop(erad)s, gebras.** Let $\mathcal{P}$ and $\mathcal{Q}$ be two prop(erad)s. A morphism $\mathcal{P} \xrightarrow{\Phi} \mathcal{Q}$ of $\mathbb{S}$-bimodules is a *morphism of prop(erad)s* if it commutes with the products and the units of $\mathcal{P}$ and $\mathcal{Q}$. In this case, we say that $\mathcal{Q}$ is a *representation* of $\mathcal{P}$.

We will be mainly interested in representations of the following form. Let X be a dg vector space. We consider the $\mathbb{S}$-bimodule End_X defined by $\mathrm{End}_X(m,n) := \mathrm{Hom}_{\mathbb{K}}(X^{\otimes n}, X^{\otimes m})$. The natural composition of maps provides this $\mathbb{S}$-bimodule with a structure of prop and properad. It is called the *endomorphism prop(erad)* of the space X.

Props and properads are meant to model the operations acting on types of algebras or bialgebras in a generalized sense. When $\mathcal{P}$ is a prop(erad), we call $\mathcal{P}$*-gebra* a dg vector space X with a morphism of prop(erad)s $\mathcal{P} \to \mathrm{End}_X$, that is a representation of $\mathcal{P}$ of the form End_X. When $\mathcal{P}$ is an operad, a $\mathcal{P}$-gebra is an algebra over $\mathcal{P}$. To code operations with multiple inputs and multiple outputs acting on an algebraic structure, we cannot use operads anymore and we need to use prop(erad)s. The categories of (involutive) Lie bialgebras and (involutive) Frobenius bialgebras are categories of gebras over a properad (see Section 9). The categories of (classical) associative bialgebras and infinitesimal Hopf algebras (see [Agu00]) are governed by non-symmetric properads. In these cases, the associated prop is freely obtained from a properad. Therefore, the prop does not model more relations than the properad and the two categories of gebras over the prop and the properad are equal. For more details, see the beginning of Section 5.5.

2. Convolution prop(erad)

When A is an associative algebra and C a coassociative coalgebra, the space of morphisms $\mathrm{Hom}_{\mathbb{K}}(C, A)$ from C to A is naturally an associative algebra with the convolution product. We generalize this property to prop(erad)s, that is the space of morphisms $\mathrm{Hom}(\mathcal{C}, \mathcal{P})$ from a coprop(erad) $\mathcal{C}$ and a prop(erad) $\mathcal{P}$ is a prop(erad). From this rich structure, we get general operations, the main one being the *intrinsic* Lie bracket used to study the deformation theory of algebraic structures later in 8.2.

2.1. **Convolution prop(erad).** For two $\mathbb{S}$-bimodules $M = \{M(m,n)\}_{m,n}$ and $N = \{N(m,n)\}_{m,n}$, we denote by $\mathrm{Hom}(M,N)$ the collection $\{\mathrm{Hom}_{\mathbb{K}}\big(M(m,n), N(m,n)\big)\}_{m,n}$ of morphisms of $\mathbb{K}$-modules. It is an $\mathbb{S}$-bimodule with the action by conjugation, that is

$$(\sigma.f.\tau)(x) := \sigma.(f(\sigma^{-1}.x.\tau^{-1})).\tau,$$

for $\sigma \in \mathbb{S}_m$, $\tau \in \mathbb{S}_n$ and $f \in \mathrm{Hom}\big(M,N\big)(m,n)$. An invariant element under this action is an equivariant map from M to N, that is $\mathrm{Hom}(M,N)^{\mathbb{S}} = \mathrm{Hom}^{\mathbb{S}}(M,N)$.

When $\mathcal{C}$ is a coassociative coalgebra and $\mathcal{P}$ is an associative algebra, $\mathrm{Hom}(\mathcal{C},\mathcal{P})$ is an associative algebra known as the *convolution algebra.* When $\mathcal{C}$ is a cooperad and $\mathcal{P}$ is an operad, $\mathrm{Hom}(\mathcal{C},\mathcal{P})$ is an operad called the *convolution operad* by C. Berger and I. Moerdijk in [BM03] Section 1. We extend this construction to properads and props.

Lemma 2. *Let* $\mathcal{C}$ *be a coprop(erad) and* $\mathcal{P}$ *be a prop(erad). The space of morphisms* $\mathrm{Hom}(\mathcal{C},\mathcal{P}) = \mathcal{P}^{\mathcal{C}}$ *is a prop(erad).*

Proof. We use the notations of Section 1.1 (see also Section 1.2 of [Val07a]). We define an associative and unital map $\mu_{\mathcal{P}^{\mathcal{C}}} : \mathcal{P}^{\mathcal{C}} \boxtimes_{\mathbb{S}} \mathcal{P}^{\mathcal{C}} \to \mathcal{P}^{\mathcal{C}}$ as follows. Let $\mathcal{G}^2(f_1, \ldots, f_r; g_1, \ldots, g_s) \in \mathcal{P}^{\mathcal{C}} \boxtimes \mathcal{P}^{\mathcal{C}}(m,n)$ be a 2-levelled graph whose vertices of the first level are labelled by $f_1, \ldots, f_r$ and whose vertices of the second level are labelled by $g_1, \ldots, g_s$. The image of $\mathcal{G}^2(f_1, \ldots, f_r;\, g_1, \ldots, g_s)$ under $\mu_{\mathcal{P}^\mathcal{C}}$ is the composite

$$\mathcal{C} \xrightarrow{\Delta_\mathcal{C}} \mathcal{C}\boxtimes^{\mathbb{S}}\mathcal{C} \rightarrowtail \mathcal{C}\boxtimes\mathcal{C} \xrightarrow{\widetilde{\mathcal{G}}^2(f_1,\ldots,f_r;\,g_1,\ldots,g_s)} \mathcal{P}\boxtimes\mathcal{P} \twoheadrightarrow \mathcal{P}\boxtimes_{\mathbb{S}}\mathcal{P} \xrightarrow{\mu_\mathcal{P}} \mathcal{P},$$

where $\widetilde{\mathcal{G}}^2(f_1, \ldots, f_r;\, g_1, \ldots, g_s)$ applies f_i on the element of $\mathcal{C}$ indexing the i^{th} vertex of the first level and g_j on the element of $\mathcal{C}$ indexing the j^{th} vertex of the second level of an element of $\mathcal{C}\boxtimes\mathcal{C}$. Since the action of the symmetric groups on $\mathcal{P}^\mathcal{C}$ is defined by conjugation and since the image of the coproduct lives in the space of invariants, this map factors through the coinvariants, that is $\mathcal{P}^\mathcal{C}\boxtimes_{\mathbb{S}}\mathcal{P}^\mathcal{C} \to \mathcal{P}^\mathcal{C}$.
The unit is given by the map $\mathcal{C} \xrightarrow{\varepsilon} I \xrightarrow{\eta} \mathcal{P}$. The associativity of $\mu_{\mathcal{P}^\mathcal{C}}$ comes directly from the coassociativity of $\Delta_\mathcal{C}$ and the associativity of $\mu_\mathcal{P}$. $\square$

Definition. The properad $\mathrm{Hom}(\mathcal{C}, \mathcal{P})$ is called the *convolution prop(erad)* and is denoted by $\mathcal{P}^\mathcal{C}$.

Assume now that $(\mathcal{C}, d_\mathcal{C})$ is a dg coprop(erad) and $(\mathcal{P}, d_\mathcal{P})$ is a dg prop(erad). The *derivative* of a graded linear map f from $\mathcal{C}$ to $\mathcal{P}$ is defined as follows

$$D(f) := d_\mathcal{P} \circ f - (-1)^{|f|} f \circ d_\mathcal{C}.$$

A 0-cycle for this differential is a morphism of chain complexes, that is it commutes with the differentials. In Section 7.4, we give a geometric interpretation of this derivative. The derivative is a derivation for the product of the prop(erad) $\mathrm{Hom}(\mathcal{C}, \mathcal{P})$ that verifies $D^2 = 0$. We sum up these relations in the following proposition. The same result holds in the non-symmetric case.

Proposition 3. *When* $(\mathcal{C}, d_\mathcal{C})$ *is a dg coprop(erad) and* $(\mathcal{P}, d_\mathcal{P})$ *is a dg prop(erad),* $(\mathrm{Hom}(\mathcal{C}, \mathcal{P}), D)$ *is a dg prop(erad).*
When $(\mathcal{C}, d_\mathcal{C})$ *is a dg non-symmetric coprop(erad) and* $(\mathcal{P}, d_\mathcal{P})$ *is a dg non-symmetric prop(erad),* $(\mathrm{Hom}(\mathcal{C}, \mathcal{P}), D)$ *is a dg non-symmetric prop(erad).*

2.2. **Lie-admissible products and Lie brackets associated to a properad.** In [KM01], the authors proved that the total space $\oplus_n \mathcal{P}(n)$, as well as the space of coinvariants $\oplus_n \mathcal{P}(n)_{\mathbb{S}_n}$, of an operad is endowed with a natural Lie bracket. This Lie bracket is the anti-symmetrization of the preLie product $p \circ q = \sum_i p \circ_i q$ defined by the sum on all possible ways of composing two operations p and q. Notice that this result was implicitly stated by Gerstenhaber in [Ger63]. We generalize this results to properads.

For any pair of elements, μ and ν, in a (non-symmetric) properad $\mathcal{P}$, we denote by $\mu \circ \nu$ the sum of all the possible compositions of μ by ν along any 2-levelled graph with two vertices in $\mathcal{P}$. For another element η in $\mathcal{P}$, the composition $(\mu \circ \nu) \circ \eta$ is spanned by graphs of the form



,


or


.

Let us denote by $\mu \circ (\nu,\, \eta)$ the summand spanned by graphs the first type.
In the same way, $\mu \circ (\nu \circ \eta)$ is spanned by graphs of the form



,


or


.

and we denote by $(\mu,\, \nu) \circ \eta$ the summand of $\mu \circ (\nu \circ \eta)$ spanned by graphs of the first (from the left) type. With these notations, we have in $\mathcal{P}$ the following formula

$$(\mu \circ \nu) \circ \eta - \mu \circ (\nu \circ \eta) = \mu \circ (\nu,\, \eta) - (\mu,\, \nu) \circ \eta.$$

When $\mathcal{P} = A$ is concentrated in arity $(1, 1)$, it is an associative algebra. In this case, the product $\circ$ is the associative product of A. When $\mathcal{P}$ is an operad, the operation $(\mu,\, \nu) \circ \eta$ vanishes and the product $\mu \circ \nu$ is right symmetric, that is $(\mu \circ \nu) \circ \eta - \mu \circ (\nu \circ \eta) = (\mu \circ \eta) \circ \nu - \mu \circ (\eta \circ \nu)$. Such a product is called *preLie*. In the general case of properads, this product verifies a weaker relation called *Lie-admissible* because its anti-symmetrized bracket verifies the Jacobi identity. Denote by $\mathrm{As}(\mu,\, \nu,\, \eta) := (\mu \circ \nu) \circ \eta - \mu \circ (\nu \circ \eta)$ the associator of μ, ν and η.

Definition (Lie-admissible algebra)**.** A graded vector space A with a binary product $\circ$ is called a (graded) *Lie-admissible* algebra if one has $\sum_{\sigma \in \mathbb{S}_3} \mathrm{sgn}(\sigma)\mathrm{As}(-,\, -,\, -)^{\sigma} = 0$, where, for instance, $\mathrm{As}(-,\, -,\, -)^{(23)}$ applied to μ, ν and η is equal to $(-1)^{|\nu||\eta|}((\mu \circ \eta) \circ \nu - \mu \circ (\eta \circ \nu))$. A *differential graded Lie-admissible algebra* (or dg Lie-admissible algebra for short) is a dg module (A, d_A) endowed with a Lie-admissible product $\circ$ such that the d_A is a derivation.

Proposition 4. *Let $\mathcal{P}$ be a dg properad or a non-symmetric dg properad, the space $\bigoplus_{m,n} \mathcal{P}(m,n)$, endowed with the product $\circ$, is a dg Lie-admissible algebra.*

Proof. Let $H = \{id,\, (23)\}$ and $K = \{id,\, (12)\}$ be two subgroups of $\mathbb{S}_3$. We have

$$\begin{aligned}
\sum_{\sigma \in \mathbb{S}_3} \mathrm{sgn}(\sigma)\mathrm{As}(-,\, -,\, -)^{\sigma} &= \sum_{\sigma \in \mathbb{S}_3} \mathrm{sgn}(\sigma)\big((- \circ (- \circ -))^{\sigma} - ((- \circ -) \circ -)^{\sigma}\big) \\
&= \sum_{\tau H \in \mathbb{S}_3 \backslash H} \mathrm{sgn}(\tau) \underbrace{\Big((- \circ (-,\, -))^{\tau} - (- \circ (-,\, -))^{\tau(23)}\Big)}_{=0} - \\
&\quad \sum_{\omega K \in \mathbb{S}_3 \backslash K} \mathrm{sgn}(\omega) \underbrace{\Big(((-,\, -) \circ -)^{\omega} - ((-,\, -) \circ -)^{\omega(12)}\Big)}_{=0} \\
&= 0.
\end{aligned}$$

☐

Actually on the direct sum $\bigoplus_{m,n} \mathcal{P}(m,n)$ of the components of a properad, there are higher operations with $r+s$ inputs which turns it into a "non-differential B_∞-algebra". We refer to the next section for more details.

For a prop $\mathcal{P}$, we still define the product $p \circ q$ on $\bigoplus_{m,n} \mathcal{P}(m,n)$ by all the possible ways of composing the operations p and q, that is all vertical composites and the horizontal one.

Proposition 5. [1] *Let $\mathcal{P}$ be a dg prop or a non-symmetric dg prop, the space $\bigoplus_{m,n} \mathcal{P}(m,n)$, endowed with the product $\circ$, is a dg associative algebra.*

Proof. We denote by $p \circ_v q$ the sum of all vertical (connected) composites of p and q and by $p \circ_h q$ the horizontal composite. We continue to use the notation $p \circ_v (q,r)$ to represent the composite of an operation p connected to two operations q and r above. We have (in degree 0)

$$\begin{aligned}
&(p \circ q) \circ r = (p \circ_v q + p \circ_h q) \circ r = \\
&p \circ_v q \circ_v r + p \circ_v (q,r) + (p \circ_v q) \circ_h r + (p \circ_v r) \circ_h q + p \circ_h (q \circ_v r) + (p,q) \circ_v r + p \circ_h q \circ_h r,
\end{aligned}$$

and

$$\begin{aligned}
&p \circ (q \circ r) = p \circ (q \circ_v r + q \circ_h r) = \\
&p \circ_v q \circ_v r + (p,q) \circ_v r + p \circ_h (q \circ_v r) + (p \circ_v q) \circ_h r + q \circ_h (p \circ_v r) + p \circ_v (q,r) + p \circ_h q \circ_h r.
\end{aligned}$$

Since the horizontal product is commutative, $(p \circ_v r) \circ_h q$ is equal to $q \circ_h (p \circ_v r)$, which finally implies $(p \circ q) \circ r = p \circ (q \circ r)$. ☐

These structures pass to coinvariants $\bigoplus \mathcal{P}_{\mathbb{S}} := \bigoplus_{m,n} \mathcal{P}(m,n)_{\mathbb{S}_m^{\mathrm{op}} \times \mathbb{S}_n}$ as follows.

Proposition 6. *Let $\mathcal{P}$ be a dg properad (respectively dg prop), the dg Lie-admissible (associative) product $\circ$ on $\bigoplus \mathcal{P}$ induces a dg Lie-admissible (associative) product on the space of coinvariants $\bigoplus \mathcal{P}_{\mathbb{S}}$.*

---

[1] This result was mentioned to the second author by M.M. Kapranov (long time ago).

Proof. It is enough to prove that the space $C := \{p - \tau.p.\nu\, ;\, p \in \mathcal{P}(m,n),\ \tau \in \mathbb{S}_m,\ \nu \in \mathbb{S}_n\}$ is a two-sided ideal for the Lie-admissible product $\circ$. Let us denote $p \circ q$ by $\sum_\sigma \mu(p,\, \sigma,\, q)$, where μ is the composition map of the properad $\mathcal{P}$ and where σ runs thought connected permutations. For any $\tau \in \mathbb{S}_m$, we have

$$(p - \tau.p) \circ q = \sum_\sigma \big(\mu(p,\, \sigma,\, q) - \mu(\tau.p,\, \sigma,\, q)\big) = \sum_\sigma \big(\mu(p,\, \sigma,\, q) - \tau_\sigma.\mu(p,\, \sigma,\, q)\big) \in C,$$

where τ_σ is a permutation which depends on σ. For any $\nu \in \mathbb{S}_n$, we have

$$\begin{aligned}(p - p.\nu) \circ q &= \sum_\sigma \mu(p,\, \sigma,\, q) - \sum_\sigma \mu(p,\, \nu.\sigma,\, q) = \sum_\sigma \mu(p,\, \sigma,\, q) - \sum_\sigma \mu(p,\, \sigma',\, q).\nu_\sigma \\ &= \sum_\sigma \big(\mu(p,\, \sigma,\, q) - \mu(p,\, \sigma,\, q).\nu_{\sigma''}\big) \in C,\end{aligned}$$

since the connected permutations σ' obtained runs thought the same set of connected permutations as σ. Therefore, C is a right ideal. The same arguments prove that C is a left ideal. ☐

In the sequel, we will have to work with the space of invariants $\bigoplus \mathcal{P}^{\mathbb{S}} := \bigoplus_{m,n} \mathcal{P}(m,n)^{\mathbb{S}_m^{\mathrm{op}} \times \mathbb{S}_n}$, and not coinvariants, of a properad. Since we work over a field of characteristic zero, both are canonically isomorphic. Let V be a vector space with an action of a finite group G. The subspace of invariants is defined by $V^G := \{v \in V \mid v.g = v\ ,\ \forall g \in G\}$ and the quotient space of coinvariants is defined by $V_G := V/ < v - v.g,\ \forall (v,g) \in V \times G >$. The map from V^G to V_G is the composite of the inclusion $V^G \rightarrowtail V$ followed by the projection $V \twoheadrightarrow V_G$. The inverse map $V_G \to V^G$ is given by $[v] \mapsto \dfrac{1}{|G|} \sum_{g \in G} v.g$, where $[v]$ denotes the class of v in V_G.

Corollary 7. *Let $\mathcal{P}$ be a dg properad (respectively dg prop), its total space of invariant elements $\bigoplus \mathcal{P}^{\mathbb{S}}$ is a dg Lie-admissible algebra (dg associative algebra).*

The Lie-admissible relation of a product $\circ$ is equivalent to the Jacobi identity $[[-,\, -],\, -] + [[-,\, -],\, -]^{(123)} + [[-,\, -],\, -]^{(132)} = 0$ for its induced bracket $[\mu,\, \nu] := \mu \circ \nu - (-1)^{|\mu||\nu|} \nu \circ \mu$.

Theorem 8. *Let $\mathcal{P}$ be a dg properad (respectively dg prop), its total space $\bigoplus \mathcal{P}$, the total space of coinvariant elements $\bigoplus \mathcal{P}_{\mathbb{S}}$ and the total space of invariant elements $\bigoplus \mathcal{P}^{\mathbb{S}}$ are dg Lie algebras.*

The first of this statement is also true for non-symmetric dg prop(erad)s.

2.3. LR-algebra associated to a properad.

On the total space of a properad, we have constructed a binary product $\circ$ in the previous section. We now define more general operations with multiple inputs.

Definition (LR-operations)**.** Let $(\mathcal{P},\, \mu)$ be a properad and $p_1, \ldots,\, p_r$ and $q_1, \ldots,\, q_s$ be elements of $\mathcal{P}$. Their *LR-operation* $\{p_1, \ldots,\, p_r\}\{q_1, \ldots,\, q_s\}$ is defined by

$$\sum_\sigma \mu(p_1, \ldots,\, p_r;\, \sigma;\, q_1, \ldots,\, q_s),$$

where σ runs through connected permutations.

In order words, the LR-product is the sum over all possible ways to compose the elements of $\mathcal{P}$. These operations are obviously graded symmetric with respect to Koszul-Quillen sign convention, that is

$$\{p_1, \ldots, p_r\}\{q_1, \ldots, q_s\} = \varepsilon(\sigma, p_1, \ldots, p_r).\varepsilon(\tau, q_1, \ldots, q_s)\{p_{\sigma(1)}, \ldots, p_{\sigma(r)}\}\{q_{\tau(1)}, \ldots, q_{\tau(s)}\},$$

for $\sigma \in \mathbb{S}_r$ and $\tau \in \mathbb{S}_s$. The element $\varepsilon(\sigma, p_1, \ldots, p_r) \in \{+1, -1\}$ stands for the Koszul-Quillen signs induced by the permutations of the graded elements $p_1, \ldots, p_r$ under σ. Notice that the Lie-admissible product is equal to $p \circ q := \{p\}\{q\}$. By convention, we set $\{\,\}\{\,\} = 0$, $\{\,\}\{q\} = q$, $\{p\}\{\,\} = p$ and $\{\,\}\{q_1, \ldots, q_s\} = 0$ for $s > 1$, $\{p_1, \ldots, p_r\}\{\,\} = 0$ for $r > 1$. The name *LR-operations* stands for Left-Right operations as well as for Loday-Ronco operations since such operations are used in [LR06] to extend Cartier-Milnor-Moore Theorem to non-cocommutative Hopf algebras.

Proposition 9. *The LR-operations satisfy the relations of a "non-differential B_∞-algebra", that is, for all $o_1, \dots, o_r, p_1, \dots, p_s, q_1, \dots, q_t$ in $\mathcal{P}$.*

$$\sum_{\Theta} \varepsilon\{\{o_1,\dots,o_{i_1}\}\{p_1,\dots,p_{j_1}\},\dots,\{o_{i_1+\cdots+i_{a-1}+1},\dots,o_r\}\{p_{j_1+\cdots+j_{a-1}+1},\dots,p_r\}\}\{q_1,\dots,q_t\}$$

$$=\sum_{\Theta'} \varepsilon'\{o_1,\dots,o_s\}\{\{p_1,\dots,p_{k_1}\}\{q_1,\dots,q_{l_1}\},\dots,\{p_{k_1+\cdots+k_{b-1}+1},\dots p_s\}\{q_{l_1+\cdots+l_{b-1}+1},\dots,q_t\}\},$$

where Θ runs over $1 \leqslant a \leqslant Max(r,s)$, $i_1,\dots,i_a \geqslant 0$ such that $i_1+\cdots+i_a=r$, $j_1,\dots,j_a \geqslant 0$ such that $j_1+\cdots+j_a=s$ and where Θ' runs over $1 \leqslant b \leqslant Max(s,t)$, $k_1,\dots,k_b \geqslant 0$ such that $k_1+\cdots+k_b=s$, $l_1,\dots,l_b \geqslant 0$ such that $l_1+\cdots+l_b=t$. The sign ε comes from the permutations of the o and the p and the sign ε' comes from the permutations of the p and the q.

Proof. It is a direct translation to LR-operations of the associativity of the operad $\mathcal{P}$. See also, Example 1.7 (d) of [LR06] and Lemma 2. ☐

Therefore, the total space $\oplus\mathcal{P}$ of a properad $\mathcal{P}$, with the LR-operations, forms a "non-differential B_∞", structure that we call a *LR-algebra*. The same result also holds for non-symmetric prop(erad)s.

Proposition 10. *The $\mathcal{P}$ be a dg prop(erad), its total space $\oplus\mathcal{P}$, the total space of coinvariants elements $\oplus\mathcal{P}_{\mathbb{S}}$ and the total space of invariants elements $\oplus\mathcal{P}^{\mathbb{S}}$ form a LR-algebra.*

Proof. The structure of LR-algebra of $\oplus\mathcal{P}$ factors through the coinvariant elements $\oplus\mathcal{P}_{\mathbb{S}}$ by the same arguments as in Proposition 6. Since the space of coinvariant and invariant elements are isomorphic, we can transfer this structure to invariant elements. ☐

2.4. **Lie-admissible bracket and LR-algebra of a convolution properad.** Since $\mathrm{Hom}(\mathcal{C},\mathcal{P})$ is an properad, it has a Lie-admissible bracket and more generally it enjoys a structure of LR-algebra by the preceding sections. We make these structures explicit here. We will use them later on in our study of deformation theory (see Section 8.2).

Definition (Convolution product)**.** Let f and g be two elements of $\mathrm{Hom}(\mathcal{C},\,\mathcal{P})$. Their *convolution product* $f \star g$ is defined by the following composite

$$\begin{tikzcd} \mathcal{C} \arrow[r, "\Delta_{(1,\,1)}"] & \mathcal{C} \boxtimes_{(1,1)} \mathcal{C} \arrow[r, "f \boxtimes_{(1,1)} g"] & \mathcal{P} \boxtimes_{(1,1)} \mathcal{P} \arrow[r, "\mu"] & \mathcal{P}. \end{tikzcd}$$

Since the partial coproduct of a coproperad (or a cooperad) is not coassociative in general, the convolution product is not associative.

Proposition 11. *Let $\mathcal{P}$ be a dg prop(erad) and $\mathcal{C}$ be a dg coprop(erad). The convolution product $\star$ on $\oplus\mathrm{Hom}(\mathcal{C},\,\mathcal{P})$ is equal to the product $\circ$ associated to the convolution dg prop(erad). In the case of dg (co)properads, it is dg Lie-admissible and for dg (co)props, it is dg associative.*
This convolution product is stable on the space of invariant elements $\oplus\mathrm{Hom}^{\mathbb{S}}(\mathcal{C},\,\mathcal{P})$ with respect to the action of the symmetric groups.

Proof. The image of the map $\Delta_{(1,1)}$ is a sum over all possible 2-levelled graphs with two vertices indexed by some elements of $\mathcal{C}$. Therefore, the map $\star$ is equal to the sum of all possible compositions of f and g.
Saying that f and g are invariant elements in $\mathrm{Hom}(\mathcal{C},\,\mathcal{P})$ means that they are equivariant maps. Since the composition map μ of $\mathcal{P}$ and the partial coproduct $\Delta_{(1,\,1)}$ are also equivariant maps, we have

$$\begin{aligned} (\sigma.f \star g.\tau)(c) &= \sigma.(f \star g(\sigma^{-1}.c.\tau^{-1})).\tau = \sigma.(\mu \circ (f \otimes g) \circ \Delta_{(1,\,1)}(\sigma^{-1}.c.\tau^{-1})).\tau \\ &= \sigma.\sigma^{-1}.(f \star g)(c).\tau^{-1}.\tau = f \star g(c). \end{aligned}$$

☐

Using the projections $\Delta_{(r,\,s)}$ of the coproduct of $\mathcal{C}$, we make explicit the LR-operations with r and s inputs of $\mathrm{Hom}(\mathcal{C},\,\mathcal{P})$ as follows.

Proposition 12. *Let* $f_1, \ldots, f_r$ *and* $g_1, \ldots, g_s$ *be elements of* $\mathrm{Hom}(\mathcal{C}, \mathcal{P})$. *Their* LR-operation $\{f_1, \ldots, f_r\}\{g_1, \ldots, g_s\}$ *is equal to*

$$\mathcal{C} \xrightarrow{\Delta_{(r,\, s)}} (I \oplus \underbrace{\mathcal{C}}_{r}) \boxtimes^{\mathbb{S}} (I \oplus \underbrace{\mathcal{C}}_{s}) \rightarrowtail$$

$$\underbrace{\mathcal{C} \otimes \cdots \otimes \mathcal{C}}_{r} \boxtimes \underbrace{\mathcal{C} \otimes \cdots \otimes \mathcal{C}}_{s} \xrightarrow{\{f_1,\ldots,f_r\}\boxtimes\{g_1,\ldots,g_r\}} \mathcal{P} \boxtimes \mathcal{P} \twoheadrightarrow \mathcal{P} \boxtimes_{\mathbb{S}} \mathcal{P} \xrightarrow{\mu} \mathcal{P},$$

where $\{f_1, \ldots, f_r\} = \sum_{\sigma \in \mathbb{S}_r} \varepsilon(\sigma, f_1, \ldots, f_r) f_{\sigma(1)} \otimes \cdots \otimes f_{\sigma(r)}$. *The element* $\varepsilon(\sigma, f_1, \ldots, f_r) \in \{+1, -1\}$ *stands for the Koszul-Quillen signs induced by the permutations of the graded elements* $f_1, \ldots, f_r$ *under* σ. *This means that we apply* $\{f_1, \ldots, f_r\}$ *and* $\{g_1, \ldots, g_s\}$ *everywhere we can.*

Proof. The proof is similar to the previous one. ☐

Theorem 13. *Let* $\mathcal{C}$ *be a dg coprop(erad) and* $\mathcal{P}$ *be a dg prop(erad), the space* $\oplus\mathrm{Hom}(\mathcal{C}, \mathcal{P})$ *is a dg LR-algebra and thus a dg Lie algebra, structures that are stable on the space of equivariant maps* $\oplus\mathrm{Hom}^{\mathbb{S}}(\mathcal{C}, \mathcal{P})$.

Proof. Since the $\Delta_{(r,\, s)}$ and μ are equivariant maps, the LR-operations are stable on the space of equivariant maps $\mathrm{Hom}^{\mathbb{S}}(\mathcal{C}, \mathcal{P})$ by their explicit form given in the previous proposition. ☐

Remark. In the case of the convolution properad, we do not have to transfer the structure of LR-algebra or Lie algebra from $\mathrm{Hom}(\mathcal{C}, \mathcal{P})$ to $\mathrm{Hom}^{\mathbb{S}}(\mathcal{C}, \mathcal{P})$ through the coinvariant-invariant isomorphism. These structures on directly stable on the space of invariant elements.

When $\mathcal{C} = C$ is a coassociative coalgebra and $\mathcal{P} = A$ an associative algebra, the product $\boxtimes$ is equal to $\otimes$ and is bilinear. In this case, the partial coproduct of C is equal to the coproduct of C and is coassociative. (All the $\Delta_{(r,\, s)}$ are null for $r > 1$ or $s > 1$). In this case, the product $\star$ is the classical convolution product on $\mathrm{Hom}(C, A)$, which is associative.

When $\mathcal{C}$ is a cooperad and $\mathcal{P}$ is an operad we have $\Delta_{(r,\, s)} = 0$ for $r > 1$ as the operations $\{f_1, \ldots, f_r\}\{g_1, \ldots, g_s\}$ are null unless $r = 1$. The remaining operations $\{f\}\{g_1, \ldots, g_s\}$ are graded symmetric *brace* operations coming from the brace-type relations verified by the operadic product (see [GO04, LM05]). Remark that when $\mathcal{C}$ is a non-symmetric cooperad and $\mathcal{P}$ a non-symmetric operad, we can define non-symmetric braces on $\mathrm{Hom}(\mathcal{C},\mathcal{P})$ without the sum over all permutations. In this case, we find the classical non-symmetric braces of [Ger63], see also [GV95, Val06b]. The convolution product verifies the relation $(f \star g) \star h - f \star (g \star h) = \{f\}\{g, h\}$. Therefore, in the operadic case, the (graded) symmetry of the brace products implies that the associator $(f \star g) \star h - f \star (g \star h)$ is symmetric in g and h. Hence the convolution product $\star$ on $\mathrm{Hom}(\mathcal{C}, \mathcal{P})$ is a graded preLie product. For an interpretation of the LR-operations (or braces operations) on cohomology theories, we refer the reader to Section 8.

3. Bar and cobar construction

In this section, we recall the definitions of the bar and cobar constructions for (co)properads and extend it to (co)props. We prove adjunction between these two constructions using the notion of *twisting morphism*, that is Maurer-Cartan elements in the convolution prop(erad). Finally, we show that the bar-cobar construction provides us with a canonical cofibrant resolution.

3.1. Infinitesimal bimodule over a prop(erad).

The notion of bimodule M over a prop(erad) $\mathcal{P}$, defined in a categorical way, is given by two compatible actions, left $\mathcal{P} \boxtimes M \to M$ and right $M \boxtimes \mathcal{P} \to M$. For our purposes, we need a *linearized or infinitesimal* version of bimodules. Such a phenomenon cannot be seen on the level of associative algebras since the monoidal product $\otimes$ defining them is bilinear.

The $\mathbb{S}$-bimodule $(M \oplus N) \boxtimes O$ is the sum over 2-levelled graphs with vertices on the top level labelled by elements of O and with vertices on the bottom level labelled by elements of M or N.

We denote by $(\underbrace{M}_{r} \oplus N) \boxtimes O$ the sub-$\mathbb{S}$-module of $(M \oplus N) \boxtimes O$ with exactly r elements of M on the bottom level.

Definition (Infinitesimal bimodule)**.** Let $(\mathcal{P}, \mu)$ be a prop(erad). An *infinitesimal $\mathcal{P}$-bimodule* is an $\mathbb{S}$-bimodule M endowed with two action maps of degree 0

$$\lambda \ : \ \mathcal{P} \boxtimes (\mathcal{P} \oplus \underbrace{M}_{1}) \to M \quad \text{and} \quad \rho \ : \ (\mathcal{P} \oplus \underbrace{M}_{1}) \boxtimes \mathcal{P} \to M,$$

such that the following diagrams commute

- Compatibility between the left action λ and the composition product μ of $\mathcal{P}$:

$$\begin{tikzcd}
\mathcal{P} \boxtimes \mathcal{P} \boxtimes (\mathcal{P} \oplus \underbrace{M}_{1}) \arrow[r, "\mathcal{P}\boxtimes(\lambda+\mu)"] \arrow[d, "\mu\boxtimes(\mathcal{P}\oplus M)"] & \mathcal{P} \boxtimes (\mathcal{P} \oplus \underbrace{M}_{1}) \arrow[d, "\lambda"] \\
\mathcal{P} \boxtimes (\mathcal{P} \oplus \underbrace{M}_{1}) \arrow[r, "\lambda"] & M,
\end{tikzcd}$$

- Compatibility between the right action ρ and the composition product μ of $\mathcal{P}$:

$$\begin{tikzcd}
(\mathcal{P} \oplus \underbrace{M}_{1}) \boxtimes \mathcal{P} \boxtimes \mathcal{P} \arrow[r, "(\rho+\mu)\boxtimes\mathcal{P}"] \arrow[d, "(\mathcal{P}\oplus M)\boxtimes\mu"] & (\mathcal{P} \oplus \underbrace{M}_{1}) \boxtimes \mathcal{P} \arrow[d, "\rho"] \\
(\mathcal{P} \oplus \underbrace{M}_{1}) \boxtimes \mathcal{P} \arrow[r, "\rho"] & M,
\end{tikzcd}$$

- Compatibility between the left and the right action :

$$\begin{tikzcd}
\mathcal{P} \boxtimes (\mathcal{P} \oplus \underbrace{M}_{1}) \boxtimes \mathcal{P} \arrow[r, "(\lambda+\mu)\boxtimes\mathcal{P}"] \arrow[d, "\mathcal{P}\boxtimes(\rho+\mu)"] & (\mathcal{P} \oplus \underbrace{M}_{1}) \boxtimes \mathcal{P} \arrow[d, "\rho"] \\
\mathcal{P} \boxtimes (\mathcal{P} \oplus \underbrace{M}_{1}) \arrow[r, "\lambda"] & M.
\end{tikzcd}$$

The notation $\mathcal{P} \boxtimes \mathcal{P} \boxtimes (\mathcal{P} \oplus \underbrace{M}_{1})$ stands for the sub-$\mathbb{S}$-bimodule of $\mathcal{P} \boxtimes \mathcal{P} \boxtimes (\mathcal{P} \oplus M)$ with only one M on the upper level. It is represented by linear combinations of 3-levelled graphes whose vertices are indexed by elements of $\mathcal{P}$ and just one of M on the first level. The other $\mathbb{S}$-bimodules with just one element coming from M are denoted in the same way, $\mathcal{P} \boxtimes (\mathcal{P} \oplus \underbrace{M}_{1}) \boxtimes \mathcal{P}$ has one element of M on the second level and $(\mathcal{P} \oplus \underbrace{M}_{1}) \boxtimes \mathcal{P} \boxtimes \mathcal{P}$ has one element of M on the third level.

One purpose of this notion is to define the notion of *abelian or infinitesimal* extension of a prop(erad) $\mathcal{P}$. It is defined by a prop(erad) structure on $\mathcal{P} \oplus M$, when M is an infinitesimal bimodule over $\mathcal{P}$ (see Section 8.4 Lemma 73). Another important property is that, for any sub-$\mathbb{S}$-bimodule J of $\mathcal{P}$, the ideal generated by J in $\mathcal{P}$ is equal to the free infinitesimal $\mathcal{P}$-bimodule on J.

Since the prop(erad) $\mathcal{P}$ has a unit, it is equivalent to have two actions $\lambda \ : \ \mathcal{P} \boxtimes_{(1,1)} M \to M$ and $\rho \ : \ M \boxtimes_{(1,1)} \mathcal{P} \to M$ that are compatible with the composition product of prop(erad) $\mathcal{P}$. Notice that the category of infinitesimal bimodules over a prop(erad) forms an abelian category.

Example. Any morphism of prop(erad)s $f : \mathcal{P} \to \mathcal{Q}$ defines an infinitesimal $\mathcal{P}$-bimodule structure on $\mathcal{Q}$:

$$\mathcal{P} \boxtimes_{(1,1)} \mathcal{Q} \xrightarrow{f \boxtimes \mathcal{Q}} \mathcal{Q} \boxtimes_{(1,1)} \mathcal{Q} \xrightarrow{\mu_{\mathcal{Q}}} \mathcal{Q} \quad \text{and} \quad \mathcal{Q} \boxtimes_{(1,1)} \mathcal{P} \xrightarrow{\mathcal{Q} \boxtimes f} \mathcal{Q} \boxtimes_{(1,1)} \mathcal{Q} \xrightarrow{\mu_{\mathcal{Q}}} \mathcal{Q}.$$

3.2. **(Co)Derivations.** Let $(\mathcal{P}, \mu)$ be a dg prop(erad) and (M, λ, ρ) be an infinitesimal $\mathcal{P}$-bimodule

Definition (Derivation)**.** A homogenous morphism $\partial : \mathcal{P} \to M$ is a *homogenous derivation* if

$$\partial \circ \mu_{(1,1)}(-,-) = \rho(\partial(-), -) + \lambda(-, \partial(-)).$$

This formula, applied to elements $p_1 \boxtimes_{(1,1)} p_2$ of $\mathcal{P} \boxtimes_{(1,1)} \mathcal{P}$, where p_1 and p_2 are homogenous elements of $\mathcal{P}$, gives

$$\partial \circ \mu(p_1 \boxtimes_{(1,1)} p_2) \quad = \quad \rho\big(\partial(p_1) \boxtimes_{(1,1)} p_2\big) + (-1)^{|\partial||p_1|} \lambda\big(p_1 \boxtimes_{(1,1)} \partial(p_2)\big).$$

A *derivation* is a sum of homogenous derivations. The set of homogenous derivations of degree n is denoted by $\mathrm{Der}^n(\mathcal{P},\, M)$ and the set of derivations is denoted $\mathrm{Der}^\bullet(\mathcal{P},\, M)$

Example. The differential of a dg prop(erad) $\mathcal{P}$ is a derivation of degree -1, that is an element of $\mathrm{Der}^{-1}(\mathcal{P},\, \mathcal{P})$.

In this section, we only consider derivations $\mathrm{Der}(\mathcal{P},\, \mathcal{Q})$, where the infinitesimal $\mathcal{P}$-bimodule structure on $\mathcal{Q}$ is given by a morphism of prop(erad)s $\mathcal{P} \to \mathcal{Q}$. In the rest of the text, we need the following lemma which makes explicit the derivations on a free prop(erad). For a prop(erad) $(\mathcal{Q}, \mu_{\mathcal{Q}})$, any graph $\mathcal{G}$ of $\mathcal{F}(\mathcal{Q})^{(n)}$ represents a class $\overline{\mathcal{G}}$ of levelled graphs of $\mathcal{Q}^{\boxtimes n}$. We recall that there is a morphism $\widetilde{\mu}_{\mathcal{Q}} : \mathcal{F}(\mathcal{Q}) \to \mathcal{Q}$, the counit of adjunction, equal to $\widetilde{\mu}_{\mathcal{Q}}(\mathcal{G}) := \mu_{\mathcal{Q}}^{\circ(n-1)}(\overline{\mathcal{G}})$. The morphism $\widetilde{\mu}_{\mathcal{Q}}$ is the only morphism of prop(erad)s extending the map $\mathcal{Q} \xrightarrow{\mathrm{Id}} \mathcal{Q}$.

Lemma 14. *Let $\rho : \mathcal{F}(V) \to \mathcal{Q}$ be a morphism of prop(erad)s of degree 0. Every derivation from the free dg prop(erad) $\mathcal{F}(V)$ to $\mathcal{Q}$ is characterized by its restriction on V, that is there is a canonical one-to-one correspondence $\mathrm{Der}^n_\rho(\mathcal{F}(V),\, \mathcal{Q}) \cong \mathrm{Hom}^{\mathbb{S}}_n(V,\, \mathcal{Q})$.*
For every morphism of dg $\mathbb{S}$-bimodules $\theta : V \to \mathcal{Q}$, we denote the unique derivation which extends θ by ∂_θ. The image of an element $\mathcal{G}(v_1, \ldots, v_n)$ of $\mathcal{F}(V)^{(n)}$ under ∂_θ is

$$\partial_\theta(\mathcal{G}(v_1, \ldots, v_n)) = \sum_{i=1}^{n} (-1)^{|\theta|.(|v_1|+\cdots+|v_{i-1}|)} \widetilde{\mu}_{\mathcal{Q}}\big(\mathcal{G}(\rho(v_1), \ldots, \rho(v_{i-1}), \theta(v_i), \rho(v_{i+1}), \ldots, \rho(v_n))\big).$$

Proof. Let us denote by θ the restriction of the derivation ∂ on V, that is $\theta = \partial_V : V \to \overline{\mathcal{Q}}$. From θ, we can construct the whole derivation ∂ by induction on the weight n of the free prop(erad) $\mathcal{F}(V)$ as follows.
For $n = 1$, we have $\partial^1_\theta = \theta : V \to \mathcal{Q}$. Suppose now that $\partial^n_\theta : \mathcal{F}(V)^{(n)} \to \mathcal{Q}$ is constructed and it is given by the formula of the Lemma. Any simple element of $\mathcal{F}(V)^{(n+1)}$ represented by a graph with $n+1$ vertices indexed by elements of V is the concatenation of a graph with n vertices with an extra vertex from the top or the bottom. In the last case, ∂^{n+1}_θ is given the commutative diagram

$$\begin{tikzcd}
\mathcal{F}(V)^{(n+1)} \arrow[rr, "\partial^{n+1}_\theta"] & & \mathcal{Q} \\
V \boxtimes_{(1,1)} \mathcal{F}(V)^{(n)} \arrow[u, "\mu_{\mathcal{F}(V)}"'] \arrow[rr, "\rho \boxtimes \partial^n_\theta + \partial^n_\theta \boxtimes \rho"] & & \mathcal{Q} \boxtimes_{(1,1)} \mathcal{Q}. \arrow[u, "\mu_{\mathcal{Q}}"']
\end{tikzcd}$$

The other case is dual. It is easy to check that the formula is still true for elements of $\mathcal{F}(V)^{(n+1)}$, that is graphs with $n+1$ vertices. Finally, since ρ is a morphism of prop(erad)s, ∂_θ is well defined and is a derivation. □

Example. A differential ∂ on a free prop(erad) $\mathcal{F}(V)$ is a derivation of $\mathrm{Der}^{-1}_{\mathrm{Id}}(\mathcal{F}(V),\, \mathcal{F}(V))$ such that $\partial^2 = 0$.

Definition (quasi-free prop(erad))**.** A dg prop(erad) $(\mathcal{F}(V),\,\partial)$ such that the underlying prop(erad) is free is called a *quasi-free* prop(erad).

Notice that in a quasi-free prop(erad), the differential is not freely generated and is a derivation of the form given above.

Dually, let $(\mathcal{C},\Delta^{\mathcal{C}})$ and $(\mathcal{D},\Delta^{\mathcal{D}})$ be two coaugmented dg coprop(erad)s and let $\rho\,:\,\mathcal{C}\to\mathcal{D}$ be a morphism of coaugmented dg coprop(erad)s of degree 0. One can define the dual notion of infinitesimal comodule over a coprop(erad) and general coderivations. Since we only need coderivations between two coprop(erad)s, we do not go into such details here.

Definition (Coderivation)**.** A homogeneous morphism $d\,:\,\mathcal{C}\to\mathcal{D}$ is a *homogeneous coderivation* of ρ if the following diagram is commutative

$$\begin{tikzcd}
\mathcal{C} \arrow[rr, "d"] \arrow[d, "\Delta^{\mathcal{C}}_{(1,1)}"] & & \mathcal{D} \arrow[d, "\Delta^{\mathcal{D}}_{(1,1)}"] \\
\mathcal{C}\boxtimes\mathcal{C} \arrow[rr, "d\boxtimes\rho+\rho\boxtimes d"] & & \mathcal{D}\boxtimes\mathcal{D}.
\end{tikzcd}$$

A *coderivation* is a sum of homogenous coderivations. The space of coderivations is denoted by $\mathrm{Coder}^{\bullet}_{\rho}(\mathcal{C},\,\mathcal{D})$.

Example. The differential of a dg coprop(erad) $\mathcal{C}$ is a coderivation of degree -1.

Remark. For a cooperad $\mathcal{D}$, we can define a more general notion of coderivation form a $\mathcal{D}$-cobimodule to $\mathcal{D}$ by a similar formula. The definition given here is a particular case. Since $\rho\,:\,\mathcal{C}\to\mathcal{D}$ is a morphism of coprop(erad)s, it provides $\mathcal{C}$ with a natural structure of $\mathcal{D}$-cobimodule.

As explained in the first section, the dual statement of Lemma 14 holds only for connected coprop(erad)s.

Lemma 15. *Let $\mathcal{C}$ be a connected coprop(erad) and let $\rho\,:\,\mathcal{C}\to\mathcal{F}^c(W)$ be a morphism of augmented coprop(erad)s. Every coderivation from $\mathcal{C}$ to the cofree connected coprop(erad) $\mathcal{F}^c(W)$ is characterized by its projection on W, that is there is a canonical one-to-one correspondence $Coder^n_{\rho}(\mathcal{C},\,\mathcal{F}^c(W))\cong\mathrm{Hom}^{\mathbb{S}}_n(\overline{\mathcal{C}},\,W)$.*

Proof. The proof is similar to the one of Lemma 14 and goes by induction on r, where F_r stands for the coradical filtration of $\mathcal{C}$. The assumption that the coprop(erad) $\mathcal{C}$ is connected ensures that the image of an element X of F_r under d lives in $\bigoplus_{n\le r}\mathcal{F}^c(W)^{(n)}$. Therefore, $d(X)$ is finite and d is well defined. □

We denote by d_ν the unique coderivation which extends a map $\nu\,:\,\overline{\mathcal{C}}\to W$.

Example. A differential d on a cofree coprop(erad) $\mathcal{F}^c(W)$ is a coderivation of $\mathrm{Der}^{-1}_{\mathrm{Id}}(\mathcal{F}^c(W),\,\mathcal{F}^c(W))$ such that $d^2=0$. By the preceding lemma, it is characterized by the composite $\mathcal{F}^c(W)\xrightarrow{d}\mathcal{F}^c(W)\twoheadrightarrow W$. Its explicit formula can be found in Lemma 22.

Definition (quasi-cofree coprop(erad))**.** A dg coprop(erad) $(\mathcal{F}^c(W),\,d)$ such that the underlying coprop(erad) is connected cofree is called a *quasi-cofree* coprop(erad).

3.3. **(De)Suspension.** The homological *suspension* of a dg $\mathbb{S}$-bimodule M is denoted by $sM:=\mathbb{K}s\otimes M$ with $|s|=1$, that is $(sM)_i\cong M_{i-1}$. Dually, the homological *desuspension* of M is denoted by $s^{-1}M:=\mathbb{K}s^{-1}\otimes M$ with $|s^{-1}|=-1$, that is $(s^{-1}M)_i\cong M_{i+1}$.

Let $(\mathcal{P},\,d)$ be an augmented dg $\mathbb{S}$-bimodule, that is $\mathcal{P}=\bar{\mathcal{P}}\oplus I$. A map of augmented $\mathbb{S}$-bimodules $\mu\,:\,\mathcal{F}^c(\bar{\mathcal{P}})\to\mathcal{P}$ consists of a family of morphisms of dg $\mathbb{S}$-bimodules $\mu_n\,:\,\mathcal{F}^c(\mathcal{P})^{(n)}\to\mathcal{P}$ for each integer $n\ge 1$. (For $n=0$, the map μ is the identity $I\to I$.) There is a one-to-one correspondence between maps $\{\mathcal{F}^c(\bar{\mathcal{P}})\to\mathcal{P}\}$ and maps $\{\mathcal{F}^c(s\bar{\mathcal{P}})\to s\mathcal{P}\}$. To each map $\mu\,:\,\mathcal{F}^c(\bar{\mathcal{P}})\to\mathcal{P}$, we associate the map $s\mu\,:\,\mathcal{F}^c(s\bar{\mathcal{P}})\to s\mathcal{P}$ defined as follows for $n\ge 1$,

$$(s\mu)_n\,:\,\mathcal{F}^c(s\bar{\mathcal{P}})^{(n)}\xrightarrow{\tau_n}s^n\mathcal{F}^c(\bar{\mathcal{P}})^{(n)}\xrightarrow{s^{-(n-1)}}s\mathcal{F}^c(\bar{\mathcal{P}})^{(n)}\xrightarrow{s\otimes\mu_n}s\mathcal{P},$$

where the map τ_n moves the place of the suspension elements from the vertices outside the graph. Since it involves permutations between suspensions s and elements of $\mathcal{P}$, the map τ_n yields signs by Koszul-Quillen rule. Using the fact that an element of $\mathcal{F}^c(\bar{\mathcal{P}})$ is an equivalent class of graphs with levels (see 1.4), one can make these signs explicit. The exact formula relating $(s\mu)$ to μ is

$$\mu\big(\mathcal{G}(p_1, \ldots, p_n)\big) = (-1)^{\varepsilon(p_1,\ldots,p_n)} s^{-1}(s\mu)\big(\mathcal{G}(sp_1, \ldots, sp_n)\big),$$

where $\varepsilon(p_1, \ldots, p_n) = (n-1)|p_1| + (n-2)|p_2| + \cdots + |p_{n-1}|$.
The degrees of μ and $s\mu$ are related by the formula $|(s\mu)_n| = |\mu_n| - (n-1)$. Therefore, the degree of μ_n is $n-2$ if and only if the degree of $(s\mu)_n$ is -1.

Dually, for any map of augmented $\mathbb{S}$-bimodules $\delta : \mathcal{C} \to \mathcal{F}(\overline{\mathcal{C}})$, we denote by δ_n the composite $\mathcal{C} \xrightarrow{\delta} \mathcal{F}(\overline{\mathcal{C}}) \twoheadrightarrow \mathcal{F}(\overline{\mathcal{C}})^{(n)}$. There is a one-to-one correspondence between maps $\{\mathcal{C} \to \mathcal{F}(\overline{\mathcal{C}})\}$ and maps $\{s^{-1}\mathcal{C} \to \mathcal{F}(s^{-1}\overline{\mathcal{C}})\}$. To each map $\delta : \mathcal{C} \to \mathcal{F}(\overline{\mathcal{C}})$, we associate the map $s^{-1}\delta : s^{-1}\mathcal{C} \to \mathcal{F}^c(s^{-1}\overline{\mathcal{C}})$ defined as follows, for $n \geq 1$,

$$(s^{-1}\delta)_n \ : \ s^{-1}\mathcal{C} \xrightarrow{s^{-(n-1)} \otimes \delta_n} s^{-n}\mathcal{F}(\overline{\mathcal{C}})^{(n)} \xrightarrow{\sigma_n} \mathcal{F}(s^{(-1)}\overline{\mathcal{C}})^{(n)}.$$

We have $|(s^{-1}\delta)_n| = |\delta_n| - (n-1)$. The degree of δ_n is $n-2$ if and only if the degree of $(s^{-1}\delta)_n$ is -1.

3.4. **Twisting morphism.** We generalize the notion of *twisting morphism* (or twisting cochains) of associative algebras (see E. Brown [Bro59] and J.C. [Moo71]) to prop(erad)s.

Let $\mathcal{C}$ be a dg coprop(erad) and $\mathcal{P}$ be a dg prop(erad). We proved in Theorem 13 that $\mathrm{Hom}^{\mathbb{S}}(\mathcal{C}, \mathcal{P})$ is a dg Lie-admissible algebra with the convolution product.

Definition. A morphism $\mathcal{C} \xrightarrow{\alpha} \mathcal{P}$, of degree -1, is called a *twisting morphism* if it is a solution of the *Maurer-Cartan* equation

$$D(\alpha) + \alpha \star \alpha = 0.$$

Denote by $\mathrm{Tw}(\mathcal{C}, \mathcal{P})$ the set of twisting morphisms in $\mathrm{Hom}^{\mathbb{S}}(\mathcal{C}, \mathcal{P})$, that is Maurer-Cartan elements in the convolution prop(erad). Since twisting morphisms have degree -1, it is equivalent for them to be solution of the classical Maurer-Cartan equation in the associated dg Lie algebra, that is $D(\alpha) + \frac{1}{2}[\alpha, \alpha] = 0$.

When $\mathcal{P}$ is augmented and $\mathcal{C}$ coaugmented, we will consider either a twisting morphism between $\mathcal{C}$ and $\mathcal{P}$, which sends I to 0, or the associated morphism which sends I to I and $\overline{\mathcal{C}}$ to $\bar{\mathcal{P}}$.

The following constructions show that the bifunctor $\mathrm{Tw}(-, -)$ can be represented on the left and on the right.

3.5. **Bar construction.** We recall from [Val07a] Section 4, the definition of the *bar construction* for properads and extend it to props. It is a functor

$$B \ : \ \{\text{aug. dg prop(erad)s}\} \longrightarrow \{\text{coaug. dg coprop(erad)s}\}.$$

Let $(\mathcal{P},\, \mu,\, \eta,\, \epsilon)$ be an augmented prop(erad). Denote by $\bar{\mathcal{P}}$ its augmentation ideal $\mathrm{Ker}(\mathcal{P} \xrightarrow{\epsilon} I)$. The prop(erad) $\mathcal{P}$ is naturally isomorphic to $\mathcal{P} = I \oplus \bar{\mathcal{P}}$. The bar construction $B(\mathcal{P})$ of $\mathcal{P}$ is a dg coprop(erad) whose underlying space is the cofree coprop(erad) $\mathcal{F}^c(s\bar{\mathcal{P}})$ on the suspension of $\bar{\mathcal{P}}$. The partial product of $\mathcal{P}$ induces a map of augmented $\mathbb{S}$-bimodules defined by the composite

$$\mu_2 \ : \ \overline{\mathcal{F}}^c(\bar{\mathcal{P}}) \twoheadrightarrow \mathcal{F}^c(\bar{\mathcal{P}})^{(2)} \cong \bar{\mathcal{P}} \boxtimes_{(1,1)} \bar{\mathcal{P}} \xrightarrow{\mu_{(1,1)}} \bar{\mathcal{P}}.$$

We have seen in the previous section that μ_2 induces a map $s\mu_2$. Consider the map $\mathbb{K}s \otimes \mathbb{K}s \xrightarrow{\Pi_s} \mathbb{K}s$ of degree -1 defined by $\Pi_s(s \otimes s) := s$. The map $s\mu_2$ is equal to the composite

$$\begin{aligned} s\mu_2 \quad : \quad & \overline{\mathcal{F}}^c(s\bar{\mathcal{P}}) \twoheadrightarrow \mathcal{F}^c(s\bar{\mathcal{P}})^{(2)} \cong (\mathbb{K}s \otimes \bar{\mathcal{P}}) \boxtimes_{(1,1)} (\mathbb{K}s \otimes \bar{\mathcal{P}}) \\ & \xrightarrow{\mathrm{Id} \otimes \tau \otimes \mathrm{Id}} (\mathbb{K}s \otimes \mathbb{K}s) \otimes (\bar{\mathcal{P}} \boxtimes_{(1,1)} \bar{\mathcal{P}}) \xrightarrow{\Pi_s \otimes \mu_{(1,1)}} \mathbb{K}s \otimes \bar{\mathcal{P}}. \end{aligned}$$

Since $\mathcal{F}^c(s\bar{\mathcal{P}})$ is a cofree connected coprop(erad), by Lemma 15 there exists a unique coderivation $d_2 := d_{s\mu_2} : \mathcal{F}^c(s\bar{\mathcal{P}}) \to \mathcal{F}^c(s\bar{\mathcal{P}})$ which extends $s\mu_2$. When $(\mathcal{P},\, d_{\mathcal{P}})$ is an augmented dg prop(erad), the differential $d_{\mathcal{P}}$ on $\mathcal{P}$ induces an internal differential d_1 on $\mathcal{F}^c(s\bar{\mathcal{P}})$. The total complex of this bicomplex is the *bar construction* $B(\mathcal{P},\, d_{\mathcal{P}}) := (\mathcal{F}^c(s\bar{\mathcal{P}}),\, d = d_1 + d_2)$ of the augmented dg prop(erad) $(\mathcal{P},\, d_{\mathcal{P}})$.

Notice that the relation $d^2 = 0$ can be understood conceptually from the Lie-admissible relations verified by the partial product of the prop(erad) $\mathcal{P}$.

3.6. **Cobar construction.** Dually, the *cobar construction* ([Val07a] Section 4) for coprop(eard)s is a functor

$$\Omega \ : \ \{\text{coaug. dg coprop(erad)s}\} \longrightarrow \{\text{aug. dg prop(erad)s}\}.$$

Let $(\mathcal{C},\, \Delta,\, \varepsilon,\, u)$ be a coaugmented coprop(erad). Denote by $\overline{\mathcal{C}}$ its augmentation $\mathrm{Ker}(\mathcal{C} \xrightarrow{\varepsilon} I)$. In this case, $\mathcal{C}$ splits naturally as $\mathcal{C} = I \oplus \overline{\mathcal{C}}$. The cobar construction $\Omega(\overline{\mathcal{C}})$ of $\overline{\mathcal{C}}$ is a dg prop(erad) whose underlying space is the free prop(erad) $\mathcal{F}(s^{-1}\overline{\mathcal{C}})$ on the desuspension of $\overline{\mathcal{C}}$.
The partial coproduct of $\mathcal{C}$ induces a natural map of augmented $\mathbb{S}$-bimodules defined by

$$\Delta_2 \ : \ \overline{\mathcal{C}} \xrightarrow{\Delta_{(1,1)}} \overline{\mathcal{C}} \boxtimes_{(1,1)} \overline{\mathcal{C}} \cong \mathcal{F}(\overline{\mathcal{C}})^{(2)} \rightarrowtail \overline{\mathcal{F}}(\overline{\mathcal{C}}).$$

This map gives a map $s^{-1}\Delta_2 \ : \ s^{-1}\overline{\mathcal{C}} \to \mathcal{F}(s^{-1}\overline{\mathcal{C}})$. Consider $\mathbb{K}s^{-1}$ equipped with the diagonal map $\mathbb{K}s^{-1} \xrightarrow{\Delta_s} \mathbb{K}s^{-1} \otimes \mathbb{K}s^{-1}$ of degree -1 defined by the formula $\Delta_s(s^{-1}) := s^{-1} \otimes s^{-1}$. The map $s^{-1}\Delta_2$ is equal to

$$\begin{aligned} s^{-1}\Delta_2 \quad : \quad & \mathbb{K}s^{-1} \otimes \overline{\mathcal{C}} \xrightarrow{\Delta_s \otimes \Delta_{(1,1)}} \mathbb{K}s^{-1} \otimes \mathbb{K}s^{-1} \otimes \overline{\mathcal{C}} \boxtimes_{(1,1)} \overline{\mathcal{C}} \xrightarrow{\mathrm{Id} \otimes \tau \otimes \mathrm{Id}} \\ & (\mathbb{K}s^{-1} \otimes \mathcal{C}) \boxtimes_{(1,1)} (\mathbb{K}s^{-1} \otimes \mathcal{C}) \cong \mathcal{F}(s^{-1}\overline{\mathcal{C}})^{(2)} \rightarrowtail \mathcal{F}(s^{-1}\overline{\mathcal{C}}). \end{aligned}$$

Since $\mathcal{F}(s^{-1}\overline{\mathcal{C}})$ is a free prop(erad), by Lemma 14 there exists a unique derivation $\partial_2 := \partial_{s^{-1}\Delta_2}$: $\mathcal{F}(s^{-1}\overline{\mathcal{C}}) \to \mathcal{F}(s^{-1}\overline{\mathcal{C}})$ which extends $s^{-1}\Delta_2$. When $(\mathcal{C},\, d_{\mathcal{C}})$ is an augmented dg coprop(erad), the differential $d_{\mathcal{C}}$ on $\mathcal{C}$ induces an internal differential ∂_1 on $\mathcal{F}(s^{-1}\overline{\mathcal{C}})$. The total complex of this bicomplex is the *cobar construction* $\Omega(\mathcal{C},\, d_{\mathcal{C}}) := (\mathcal{F}(s^{-1}\overline{\mathcal{C}}),\, \partial = \partial_1 + \partial_2)$ of the augmented dg coprop(erad) $(\mathcal{C},\, d_{\mathcal{C}})$.

3.7. **Bar-Cobar Adjunction.** As for derivations, a morphism of prop(erad)s is characterized by the image of the indecomposable elements. We recall this fact and the dual statement in the following lemma.

Lemma 16. *Let V be an $\mathbb{S}$-bimodule and let $\mathcal{Q}$ be a prop(erad), there is a canonical one-to-one correspondence* $\mathrm{Mor}_{\mathrm{prop(erad)s}}\big(\mathcal{F}(V), \mathcal{Q}\big) \cong \mathrm{Hom}^{\mathbb{S}}(V, \mathcal{Q})$.
Dually, let W be an $\mathbb{S}$-bimodule and let $\mathcal{C}$ be a coprop(erad), there is a canonical one-to-one correspondence $\mathrm{Mor}_{\mathrm{coprop(erad)s}}\big(\mathcal{C}, \mathcal{F}^c(W)\big) \cong \mathrm{Hom}^{\mathbb{S}}(\mathcal{C}, W)$.

Let $(\mathcal{C}, d_{\mathcal{C}})$ be a dg coprop(erad) and $(\mathcal{P}, d_{\mathcal{P}})$ be a dg prop(erad). We will apply this result to the bar and the cobar construction of $\mathcal{P}$ and $\mathcal{C}$ respectively, that is we want to describe the space of morphisms of **dg**-prop(erad)s $\mathrm{Mor}_{\mathrm{dg\ prop(erad)s}}\,(\Omega(\mathcal{C}),\, \mathcal{P})$ for instance. By the preceding lemma, this space is isomorphic to the space of morphisms of $\mathbb{S}$-bimodules $\mathrm{Hom}^{\mathbb{S}}_0(s^{-1}\mathcal{C}, \mathcal{P})$ of degree 0 whose unique extension commutes with the differentials. Therefore, this space of morphisms is the subspace of $\mathrm{Hom}^{\mathbb{S}}_{-1}(\mathcal{C}, \mathcal{P})$ whose elements satisfy a certain relation, which is exactly the Maurer-Cartan equation.

Proposition 17. *For every augmented dg prop(erad) $\mathcal{P}$ and every coaugmented dg coprop(erad) $\mathcal{C}$, there are canonical one-to-one correspondences*

$$\mathrm{Mor}_{dg\ prop(erad)s}\,(\Omega(\mathcal{C}),\, \mathcal{P}) \cong \mathrm{Tw}(\mathcal{C}, \mathcal{P}) \cong \mathrm{Mor}_{dg\ coprop(erad)s}\,(\mathcal{C},\, \mathcal{B}(\mathcal{P}))\,.$$

Proof. Since $\Omega(\mathcal{C}) = \mathcal{F}(s^{-1}(\overline{\mathcal{C}}))$, by Lemma 16 every morphism φ of $\mathbb{S}$-bimodules in $\mathrm{Hom}_0^{\mathbb{S}}(s^{-1}\mathcal{C}, \mathcal{P})$ extends to a unique morphism of prop(erad)s between $\Omega(\mathcal{C})$ and $\mathcal{P}$. The latter one commutes with the differentials if and only if the following diagram commutes

$$\begin{tikzcd}
s^{-1}\overline{\mathcal{C}} \arrow[r, "\varphi"] \arrow[d, "\partial"] & \mathcal{P} \arrow[dr, "d_{\mathcal{P}}"] & \\
\mathcal{F}(s^{-1}\overline{\mathcal{C}})^{(\leqslant 2)} \arrow[r, "\mathcal{F}(\varphi)"] & \mathcal{F}(\mathcal{P}) \arrow[r, "\widetilde{\mu}^{\mathcal{P}}"] & \mathcal{P}.
\end{tikzcd}$$

For an element $c \in \overline{\mathcal{C}}$, we use Sweedler's notation to denote the image of c under Δ_2, that is $\Delta_2(c) = \sum c' \boxtimes_{(1,1)} c''$. The diagram above corresponds to the relation

$$d_{\mathcal{P}} \circ \varphi(s^{-1}c) = \varphi \circ \partial_1(s^{-1}c) + \mu^{\mathcal{P}} \circ (\varphi \boxtimes_{(1,1)} \varphi) \circ s^{-1}\Delta_2(s^{-1}c).$$

Denote by α the desuspension of φ, that is $\alpha(c) = -\varphi(s^{-1}c)$. Since $\partial_1(s^{-1}c) = -s^{-1}\partial_{\mathcal{C}}(c)$, the relation becomes

$$-d_{\mathcal{P}} \circ \alpha(c) = \alpha \circ \partial_{\mathcal{C}}(c) + \mu^{\mathcal{P}} \circ (\alpha \boxtimes_{(1,1)} \alpha) \circ \Delta_2(c),$$

which is the Maurer-Cartan equation. ☐

Therefore, the bar and cobar constructions form a pair of adjoint functors

$$\Omega \ : \ \{\text{coaug. dg coprop(erad)s}\} \rightleftharpoons \{\text{aug. dg prop(erad)s}\} \ : \ B.$$

If we apply the isomorphisms of Proposition 17 to $\mathcal{C} = B(\mathcal{P})$, the morphism associated to the identity on $B(\mathcal{P})$ is the *counit* of the adjunction $\epsilon \,:\, \Omega(B(\mathcal{P})) \to \mathcal{P}$. In this case, we get a universal twisting morphism $B(\mathcal{P}) \to \mathcal{P}$.

The morphism associated to the identity of $\Omega(\mathcal{C})$ when $\mathcal{P} = \Omega(\mathcal{C})$ is the *counit* of the adjunction $\mathcal{C} \to B(\Omega(\mathcal{C}))$. In this case, we get a universal twisting morphism $\mathcal{C} \to \Omega(\mathcal{C})$.

Proposition 18. *Any twisting morphism $\alpha \,:\, \mathcal{C} \to \mathcal{P}$ factors through $B(\mathcal{P}) \to \mathcal{P}$ and $\mathcal{C} \to \Omega(\mathcal{C})$.*

$$\begin{tikzcd}
 & \Omega(\mathcal{C}) \arrow[dr] & \\
\mathcal{C} \arrow[ur] \arrow[rr, "\alpha"] \arrow[dr] & & \mathcal{P} \\
 & B(\mathcal{P}) \arrow[ur] &
\end{tikzcd}$$

Proof. It is a corollary of Proposition 17. ☐

3.8. **Props vs properads.** The main difference for (co)bar construction between props and properads lies on the type of graphs and compositions. The underlying module of the bar construction of a prop $\mathcal{P}$ is spanned by not necessarily connected graphs whose vertices are labelled with elements of $\mathcal{P}$. The boundary map is the unique coderivation which extends the partial product. It is given explicitly by the sum of the compositions of pair of vertices that are either adjacent (see Section 4.2) or belong to two different connected graphs. Whereas for a properad, the underlying module is spanned by connected labelled graphs and the boundary map just composes adjacent pairs of operations.

3.9. **Bar-cobar resolution.** In [Val07a] Theorem 5.8, we proved that the unit of adjunction ϵ is a canonical resolution in the weight graded case. We extend this result to any dg properad here.

Theorem 19. *For every augmented dg properad $\mathcal{P}$, the bar-cobar construction is a resolution of $\mathcal{P}$*

$$\epsilon \ : \ \Omega(B(\mathcal{P})) \xrightarrow{\sim} \mathcal{P}.$$

Proof. The bar-cobar construction of $\mathcal{P}$ is the chain complex defined on the underlying $\mathbb{S}$-bimodule $\mathcal{F}\big(s^{-1}\overline{\mathcal{F}}^{c}(s\bar{\mathcal{P}})\big)$. The differential d is the sum of three terms $d = \partial_2 + d_2 + d_{\mathcal{P}}$, where $d_{\mathcal{P}}$ is induced by the differential on $\mathcal{P}$, d_2 is induced by the differential of the bar construction $B(\mathcal{P})$ and ∂_2 is the unique derivation which extends the partial coproduct of $\mathcal{F}^c(s\bar{\mathcal{P}})$.

Define the filtration $F_s := \bigoplus_{r\leq s} \mathcal{F}\big(s^{-1}\overline{\mathcal{F}}^c(s\bar{\mathcal{P}})\big)_r$, where r denotes the total number of elements of $\bar{\mathcal{P}}$. Let $E^\bullet_{st}$ be the associated spectral sequence.
This filtration is bounded below and exhaustive. Therefore, we can apply the classical convergence theorem for spectral sequences (see [Wei94]) and prove that $E^\bullet$ converges to the homology of the bar-cobar construction.
We have that $E^0_{st} = \mathcal{F}_{s+t}\big(s^{-1}\mathcal{F}^c(s\bar{\mathcal{P}})\big)_s$, where $s+t$ is the total homological degree. From $d_2(F_s) \subset F_{s-1}$, $d_{\mathcal{P}}(F_s) \subset F_s$ and $\partial_2(F_s) \subset F_s$, we get that $d^0 = \partial_2 + d_{\mathcal{P}}$. The problem is now reduced to the computation of the homology of the cobar construction of the dg cofree connected coproperad $\mathcal{F}^c(s\bar{\mathcal{P}})$ on the dg $\mathbb{S}$-bimodule $s\bar{\mathcal{P}}$. This complex is equal to the bar-cobar construction of the weight graded properad $(\mathcal{P},\, \mu')$, where $\mathcal{P}^{(0)} = I$ and $\mathcal{P}^{(1)} = \bar{\mathcal{P}}$, such that the composition μ' is null. We conclude using Theorem 5.8 of [Val07a]. ☐

Proposition 20. *The bar-cobar resolution provides a canonical cofibrant resolution to any non-negatively graded dg properad.*

We refer the reader to the Appendix A for the model category structure on dg prop(erads)
Proof. The bar-cobar resolution is quasi-free. We conclude by Corollary 97. ☐

4. Homotopy (Co)prop(erad)s

An associative algebra is a vector space endowed with a binary product that verifies the strict associative relation. J. Stasheff defined in [Sta63] a lax version of this notion. It is the notion of an associative algebra up to homotopy or (strong) homotopy algebra. Such an algebra is a vector space equipped with a binary product that is associative only up to an infinite sequence of homotopies. In this section, we recall the generalization of this notion, that is the notion of *(strong) homotopy properad* due to J. Granåker [Gra07]. We extend it to props and we also define in details the dual notion of *(strong) homotopy coprop(erad)*, which will be essential to deal with minimal models in the next section . The notions of *homotopy non-symmetric (co)properad* and *homotopy non-symmetric (co)prop* are obtained in the same way.

4.1. **Definitions.** Following the same ideas as for algebras (associative or Lie, for instance), we define the notion of *homotopy (co)prop(erad)* via (co)derivations and (co)free (co)prop(erad)s.

Definition (Homotopy prop(erad))**.** A structure of *homotopy prop(erad)* on an augmented dg $\mathbb{S}$-bimodule $(\mathcal{P}, d_{\mathcal{P}})$ is a coderivation d of degree -1 on $\mathcal{F}^c(s\bar{\mathcal{P}})$ such that $d^2 = 0$.

A structure of homotopy prop(erad) is equivalent to a structure of quasi-cofree coprop(erad) on $s\bar{\mathcal{P}}$. We call the latter the *(generalized) bar construction of* $\mathcal{P}$ and we still denote it by $B(\mathcal{P})$. Since $\mathcal{F}^c(s\bar{\mathcal{P}})$ is a cofree connected coprop(erad), by Lemma 15 the coderivation d is characterized by the composite

$$s\mu \ :\ \mathcal{F}^c(s\bar{\mathcal{P}}) \xrightarrow{d} \mathcal{F}^c(s\bar{\mathcal{P}}) \twoheadrightarrow s\mathcal{P},$$

that is $d = d_{s\mu}$. The map $s\mu$ of degree -1 is equivalent to a unique map $\mu\,:\,\mathcal{F}^c(\bar{\mathcal{P}}) \to \mathcal{P}$, such that $\mu_n\,:\,\mathcal{F}^c(\bar{\mathcal{P}})^{(n)} \to \mathcal{P}$ has degree $n-2$. The condition $d^2 = 0$ written with the $\{\mu_n\}_n$ is made explicit in Proposition 23.

Example. A dg prop(erad) is a homotopy prop(erad) such that every map $\mu_n = 0$ for $n \geq 3$. In this case, $(\mathcal{F}^c(s\bar{\mathcal{P}}), d)$ is the bar construction of $\mathcal{P}$.

We define the notion of *homotopy coprop(erad)* by a direct dualization of the previous arguments.

Definition (Homotopy coprop(erad))**.** A structure of *homotopy coprop(erad)* on an augmented dg $\mathbb{S}$-bimodule $(\mathcal{C},\, d_{\mathcal{C}})$ is a derivation ∂ of degree -1 on $\mathcal{F}(s^{-1}\overline{\mathcal{C}})$ such that $\partial^2 = 0$.

A structure of homotopy coprop(erad) is equivalent to a structure of quasi-free prop(erad) on $s^{-1}\overline{\mathcal{C}}$. We call the latter the *(generalized) cobar construction of* $\mathcal{C}$ and we still denote it by $\Omega(\mathcal{C})$. By Lemma 14, the derivation ∂ is characterized by its restriction on $s^{-1}\overline{\mathcal{C}}$

$$s^{-1}\Delta \ :\ s^{-1}\overline{\mathcal{C}} \rightarrowtail \overline{\mathcal{F}}(s^{-1}\overline{\mathcal{C}}) \xrightarrow{\partial} \overline{\mathcal{F}}(s^{-1}\overline{\mathcal{C}}),$$

that is $\partial = \partial_{s^{-1}\Delta}$. The map $s^{-1}\Delta$ of degree -1 is equivalent a map $\Delta : \mathcal{C} \to \mathcal{F}(\overline{\mathcal{C}})$, such that the component $\Delta_n : \mathcal{C} \to \mathcal{F}(\overline{\mathcal{C}})^{(n)}$ has degree $n-2$. The condition $\partial^2 = 0$ is equivalent to relations for the $\{\Delta_n\}_n$ that we make explicit in Proposition 24.

Example. A dg coprop(erad) is a homotopy coprop(erad) such that every map $\Delta_n = 0$ for $n \geq 3$. In this case, $(\mathcal{F}(s^{-1}\overline{\mathcal{C}}), \partial)$ is the cobar construction of $\mathcal{C}$.

When $\mathcal{P}$ is concentrated in arity $(1,1)$, the definition of a homotopy properad on $\mathcal{P}$ is exactly the same than the definition of an strong homotopy algebra given by J. Stasheff in [Sta63]. Dually, when $\mathcal{C}$ is concentrated in arity (1, 1), we get the notion of strong homotopy coassociative algebra. When $\mathcal{P}$ is concentrated in arity $(1,n)$ for $n \geq 1$, we have the notion of *strong homotopy operad* (see [VdL02]). The dual notion gives the definition of a *strong homotopy cooperad*.

REMARK. By abstract nonsense, the notion of homotopy prop(erad) should also come from Koszul duality for colored operads (see [VdL03]). There exists a colored operad whose"algebras" are (partial) prop(erad)s. Such a colored operad is quadratic (the associativity relation of the partial product of a prop(erad) is an equation between compositions of two elements.) It should be a Koszul colored operad. An "algebra" over the Koszul resolution of this colored operad is exactly a homotopy prop(erad).

4.2. **Admissible subgraph.** Let $\mathcal{G}$ be a connected graph directed by a flow and denote by $\mathcal{V}$ its set of vertices. We define a partial order on $\mathcal{V}$ by the following covering relation : $i \prec j$ if i is below j according to the flow and if there is no vertex between them. In this case, we say that i and j are *adjacent* (see also [Val07a] p. 34). Examples of adjacent and non-adjacent vertices can be found in Figure 2.



FIGURE 2. The vertices 1, 2 and 2, 3 are adjacent. The vertices 1 and 3 are not adjacent.

Denote this poset by $\Pi_{\mathcal{G}}$ and consider its Hasse diagram $\mathcal{H}(\mathcal{G})$, that is the diagram composed by the elements of the poset with one edge between two of them, when they are related by a covering relation. See Figure 3 for an example.



FIGURE 3. The Hasse diagramm associated to the graph of Figure 2

Actually, $\mathcal{H}(\mathcal{G})$ is obtained from $\mathcal{G}$ be removing the external edges and by replacing several edges between two vertices by only one edge. Since $\mathcal{G}$ is connected and directed by a flow, the Hasse diagram $\mathcal{H}(\mathcal{G})$ has the same properties. A *convex subset* $\mathcal{V}'$ of $\mathcal{V}$ is a set of vertices of $\mathcal{G}$ such that for every pair $i \leq j$ in $\mathcal{V}'$ the interval $[i,j]$ of $\Pi_{\mathcal{G}}$ is included in $\mathcal{V}'$. If $\mathcal{G}$ is a connected graph of genus 0, the set of vertices of any connected subgraph of $\mathcal{G}$ is convex. This property does not hold any more for connected graphs of higher genus.

Lemma 21. *Let $\mathcal{G}$ be a connected directed graph without oriented loops and let $\mathcal{G}'$ be a connected subgraph of $\mathcal{G}$. The set of vertices of $\mathcal{G}'$ is convex if and only if the contraction of $\mathcal{G}'$ inside of $\mathcal{G}$ gives a graph without oriented loops.*

A connected subgraph $\mathcal{G}'$ with this property is called *admissible* in [Gra07]. We denote by $\mathcal{G}/\mathcal{G}'$ the graph obtained by the contraction of $\mathcal{G}'$ inside $\mathcal{G}$. See Figure 4 for an example of a admissible subgraph and an example of a non-admissible subgraph of $\mathcal{G}$. By extension, an admissible subgraph of a non-necessarily connected graph is a union of admissible subgraphs (eventually empty) of each connected component.



FIGURE 4. Example of a admissible subgraph $\mathcal{G}'$ of $\mathcal{G}$ and an example of a non-admissible subgraph of $\mathcal{G}$.

4.3. **Interpretation in terms of graphs.** Let $\mu : \mathcal{F}^c(\bar{\mathcal{P}}) \to \mathcal{P}$ be a morphism of augmented dg $\mathbb{S}$-bimodules. We denote by $\mu\big(\mathcal{G}(p_1, \ldots, p_n)\big)$ the image of an element $\mathcal{G}(p_1, \ldots, p_n)$ of $\mathcal{F}^c(\bar{\mathcal{P}})^{(n)}$ under μ. Let $\mathcal{G}'$ be a admissible subgraph of $\mathcal{G}$ with k vertices. Denote by $\mathcal{G}/\mu\mathcal{G}'(p_1, \ldots, p_n)$ the element of $\mathcal{F}^c(\bar{\mathcal{P}})^{(n-k+1)}$ obtained by composing $\mathcal{G}'(p_{i_1}, \ldots, p_{i_k})$ in $\mathcal{G}(p_1, \ldots, p_n)$ under μ. When the p_i and μ are not of degree zero, this composition induces natural signs that we make explicit in the sequel. Let start with a representative element of a class of graph $\mathcal{G}(p_1, \ldots, p_n)$ whose vertices are indexed by elements p_i, that is to say we have chosen an order between the p_i (see Section 1.4). The vertices of $\mathcal{G}'$ are indexed by elements $p_{i_1}, \ldots, p_{i_k}$. We denote by $J = (i_1, \ldots, i_k)$ the associated ordered subset of $[n] = \{1, \ldots, n\}$ and $p_J = p_{i_1}, \ldots, p_{i_k}$. Since $\mathcal{G}'$ is an admissible subgraph, its set of vertices forms a convex subset of the set of vertices of $\mathcal{G}$ (or a disjoint union of convex subsets if $\mathcal{G}$ is not connected). Therefore, it is possible to change the order of the vertices of $\mathcal{G}$ such that the vertices of $\mathcal{G}'$ are next to each others. That is there exists two ordered subsets I_1 and I_2 of $[n]$ such that the underlying subsets I_1, I_2 and J without order form a partition of $[n]$ and such that $\mathcal{G}(p_1, \ldots, p_n) = (-1)^{\varepsilon_1}\mathcal{G}(P_{I_1}, P_J, P_{I_2})$. The sign $(-1)^{\varepsilon_1}$ is given by the Koszul-Quillen sign rule from the permutation of the p_i. Now we can apply μ to get

$$\mathcal{G}/\mu\mathcal{G}'(p_1, \ldots, p_n) = (-1)^{\varepsilon_1+\varepsilon_2}\mathcal{G}/\mathcal{G}'\big(P_{I_1}, \mu(\mathcal{G}'(P_J)), P_{I_2}\big),$$

where $\varepsilon_2 = |P_{I_1}|.|\mu|$. It is an easy exercise to prove that this definition of the signs does not depend on the different choices.

Lemma 22. *Let ν be a map $\mathcal{F}^c(W) \to W$ of degree -1. The unique coderivation $d_\nu \in \mathrm{CoDer}^{-1}_{\mathrm{Id}}(\mathcal{F}^c(W), \mathcal{F}^c(W))$ which extends ν is given by*

$$d_\nu\big(\mathcal{G}(w_1, \ldots, w_n)\big) = \sum_{\mathcal{G}' \subset \mathcal{G}} \mathcal{G}/\nu\mathcal{G}'(w_1, \ldots, w_n),$$

where the sum runs over admissible subgraphs $\mathcal{G}'$ of $\mathcal{G}$.

PROOF. This formula defines a coderivation. Since the composite of d_ν with the projection on W is equal to ν, we conclude by the uniqueness property of coderivations of Lemma 15. ☐

Proposition 23. *A map $\mu : \mathcal{F}^c(\bar{\mathcal{P}}) \to \mathcal{P}$ defines a structure of homotopy prop(erad) on the augmented dg $\mathbb{S}$-bimodule $\mathcal{P}$ if and only if, for every $\mathcal{G}(p_1, \dots, p_n)$ in $\mathcal{F}^c(\bar{\mathcal{P}})$, we have*

$$\sum_{\mathcal{G}' \subset \mathcal{G}} (-1)^{\varepsilon(\mathcal{G}', p_1, \dots, p_n)} \mu\big(\mathcal{G}/\mu\mathcal{G}'(p_1, \dots, p_n)\big) = 0,$$

where the sum runs over admissible subgraphs $\mathcal{G}'$ of $\mathcal{G}$.

Proof. By definition, μ induces a structure of homotopy prop(erad) if and only if $d_{s\mu}^2 = 0$. This last condition holds if and only if the composite $\mathrm{proj}_{s\mathcal{P}} \circ d_{s\mu}^2 = (s\mu) \circ d_{s\mu}$ is zero, where $\mathrm{proj}_{s\mathcal{P}}$ is the projection on $s\mathcal{P}$. From Lemma 22, this is equivalent to

$$\sum_{\mathcal{G}' \subset \mathcal{G}} (s\mu)\big(\mathcal{G}/(s\mu)\mathcal{G}'(sp_1, \dots, sp_n)\big) = 0,$$

where the sum runs over admissible subgraphs $\mathcal{G}'$ of $\mathcal{G}$. Recall from Section 3.3 that the signs between $(s\mu)$ and μ are

$$\mu\big(\mathcal{G}(p_1, \dots, p_n)\big) = (-1)^{\varepsilon(p_1, \dots, p_n)} s^{-1}(s\mu)\big(\mathcal{G}(sp_1, \dots, sp_n)\big),$$

where $\varepsilon(p_1, \dots, p_n) = (n-1)|p_1| + (n-2)|p_2| + \cdots + |p_{n-1}|$. Therefore, μ induces a structure of homotopy prop(erad) if and only if

$$\sum_{\mathcal{G}' \subset \mathcal{G}} (-1)^{\varepsilon(\mathcal{G}', p_1, \dots, p_n)} \mu\big(\mathcal{G}/\mu\mathcal{G}'(p_1, \dots, p_n)\big) = 0,$$

where $(-1)^{\varepsilon(\mathcal{G}', p_1, \dots, p_n)}$ is product of the sign coming the composition with $s\mu$ and the sign coming from the formula between μ and $s\mu$. ☐

Remark. In the case of associative algebras, the graphs involved are ladders (branches, directed graphs just one incoming edge and one outgoing edge for each vertex) and we recover exactly the original definition of J. Stasheff [Sta63].

Dually, we have the following characterization of homotopy coprop(erad)s. Let $\mathcal{G}$ be a graph whose i^{th} vertex has n inputs and m outputs. For every graph $\mathcal{G}'$ with n inputs and m outputs, denote by $\mathcal{G} \circ_i \mathcal{G}'$ the graph obtained by inserting $\mathcal{G}'$ in $\mathcal{G}$ at the place of the i^{th} vertex.

Proposition 24. *A map $\Delta : \mathcal{C} \to \mathcal{F}(\overline{\mathcal{C}})$ defines a structure of homotopy coprop(erad) on the augmented dg $\mathbb{S}$-bimodule $\mathcal{C}$ if and only if, for every $c \in \overline{\mathcal{C}}$, we have*

$$\sum (-1)^{\rho(\mathcal{G}_i^2, c_1, \dots, c_l)} \mathcal{G}^1 \circ_i \mathcal{G}_i^2(c_1, \dots, c_{i-1}, c_1', \dots, c_k', c_{i+1}, \dots, c_l) = 0,$$

where the sum runs over elements $\mathcal{G}^1(c_1, \dots, c_l)$ and $\mathcal{G}_i^2(c_1', \dots, c_k')$ such that $\Delta(c) = \sum \mathcal{G}^1(c_1, \dots, c_l)$ and $\Delta(c_i) = \sum \mathcal{G}_i^2(c_1', \dots, c_k')$.

Proof. By definition, Δ induces a structure of homotopy coprop(erad) if and only if $\partial_{s^{-1}\Delta}^2 = 0$. Since $\partial_{s^{-1}\Delta}$ is a derivation, $\partial_{s^{-1}\Delta}^2 = 0$ is equivalent to $\partial_{s^{-1}\Delta} \circ (s^{-1}\Delta)(s^{-1}c) = 0$, for every $c \in \overline{\mathcal{C}}$. Denote $(s^{-1}\Delta)(s^{-1}c) = \sum \mathcal{G}^1(s^{-1}c_1, \dots, s^{-1}c_l)$ and $(s^{-1}\Delta)(s^{-1}c_i) = \sum \mathcal{G}_i^2(s^{-1}c_1', \dots, s^{-1}c_k')$. By the explicit formula for $\partial_{s^{-1}\Delta}$ given in Lemma 14 applied to $\rho = \mathrm{Id}_{\mathcal{F}(s^{-1}\overline{\mathcal{C}})}$, we have

$$\begin{aligned} \partial_{s^{-1}\Delta} \circ (s^{-1}\Delta)(s^{-1}c) &= \partial_{s^{-1}\Delta}\big(\sum \mathcal{G}^1(s^{-1}c_1, \dots, s^{-1}c_l)\big) \\ &= \sum \mathcal{G}^1 \circ_i \mathcal{G}_i^2(s^{-1}c_1, \dots, s^{-1}c_{i-1}, s^{-1}c_1', \dots, s^{-1}c_k', s^{-1}c_{i+1}, \dots, s^{-1}c_l) = 0 \end{aligned}$$

We get back to the map Δ with the formula

$$\Delta(c) = (-1)^{\varepsilon(c_1, \dots, c_l)} \sum \mathcal{G}^1(c_1, \dots, c_l),$$

where $\varepsilon(c_1, \dots, c_l) = (l-1)|c_1| + (l-2)|c_2| + \cdots + |c_{l-1}|$. We conclude as in proof of Proposition 23. ☐

4.4. **Homotopy non-symmetric (co)prop(erad).** It is straightforward to generalize the two previous subsections to non-symmetric (co)prop(erad)s. One has just to consider non-labelled graphs instead of graphs with leaves, inputs and outputs labelled by integers. Therefore, there is a bar and a cobar construction between non-symmetric dg prop(erad)s and non-symmetric dg coprop(erad)s. The notion that will be used in the sequel is the notion of *homotopy non-symmetric prop(erad)*. It is defined by a coderivation on the non-symmetric cofree (connected) coprop(erad). Equivalently, we can describe it in terms of non-labelled graphs like in Proposition 23. The chain complex defining the cohomology of a gebra over a non-symmetric prop(erad) has always such a structure (see Section 8.2).

4.5. **Homotopy properads and associated homotopy Lie algebras.** It was proven in [KM01] that for any operad, $\mathcal{P} = \{\mathcal{P}(n)\}$, the vector space $\bigoplus_n \mathcal{P}(n)$ has a natural structure of Lie algebra which descends to the space of coinvariants $\bigoplus_n \mathcal{P}(n)_{\mathbb{S}_n}$, which is isomorphic to the space of invariants $\bigoplus_n \mathcal{P}(n)^{\mathbb{S}_n}$. In [VdL02] this result was generalized to homotopy operads and the associated L_∞-algebras. In this section, we further extend the results of [KM01, VdL02] from homotopy operads to arbitrary homotopy prop(erad)s: $\mathcal{P} = \{\mathcal{P}(m,n)\}$.

Recall that a structure of L_∞-algebra on $\mathfrak{g}$ is given by a square-zero coderivation on $\mathcal{S}^c(s\mathfrak{g})$, where $\mathcal{S}^c(s\mathfrak{g})$ stands for the cofree cocommutative coalgebra on the suspension of $\mathfrak{g}$. Hence, such a structure is completely characterized by the image of the coderivation on $s\mathfrak{g}$, $\mathcal{S}^c(s\mathfrak{g}) \to s\mathfrak{g}$. Equivalently, an L_∞-algebra is an algebra over the minimal (Koszul) resolution of the operad $\mathcal{L}ie$. We refer the reader to Section 7.1 for more details on L_∞-algebras.

Let $\mathcal{P}$ be an $\mathbb{S}$-bimodule. We denote by $\oplus\mathcal{P}$ the direct sum of all the components of $\mathcal{P}$, that is $\bigoplus_{m,n} \mathcal{P}(m,n)$. We consider the map $\Theta : \mathcal{S}^c(\oplus\mathcal{P}) \to \mathcal{F}^c(\mathcal{P})$ defined by $\Theta(p_1 \odot \cdots \odot p_n) := \sum \mathcal{G}(p_1, \ldots, p_n)$, where the sum runs over the classes of graphs under the action of the automorphism group of the graph. This sum is finite and since a graph is a quotient of a levelled graph (see Section 1.4), the signs are well defined.

Theorem 25. *Let $\mathcal{P}$ be a homotopy properad, the direct sum $\oplus\mathcal{P}$ of its components has an induced L_∞-structure.*

Proof. We define the partial cotriple coproduct of a cofree coprop(erad) by the composite :

$$\Delta' \ : \ \mathcal{F}^c(V) \xrightarrow{\widetilde{\Delta}} \mathcal{F}^c(\mathcal{F}^c(V)) \twoheadrightarrow \mathcal{F}^c(V, \underbrace{\mathcal{F}^c(V)}_{1}),$$

where $\mathcal{F}^c(V, \underbrace{\mathcal{F}^c(V)}_{1})$ represents graphs indexed by elements of V and one element of $\mathcal{F}^c(V)$. Similarly, we define the partial cotriple coproduct of the cofree cocommutative coalgebra by :

$$\delta' \ : \ \mathcal{S}^c(V) \xrightarrow{\widetilde{\delta}} \mathcal{S}^c(\mathcal{S}^c(V)) \twoheadrightarrow \mathcal{S}^c(V, \underbrace{\mathcal{S}^c(V)}_{1}).$$

Let $s\mu : \mathcal{F}^c(s\bar{\mathcal{P}}) \to s\bar{\mathcal{P}}$ be a map of degree -1 defining a homotopy properad structure on $\mathcal{P}$, that is the following composite

$$\mathcal{F}^c(s\bar{\mathcal{P}}) \xrightarrow{\Delta'} \mathcal{F}^c(s\bar{\mathcal{P}}, \underbrace{\mathcal{F}^c(s\bar{\mathcal{P}})}_{1}) \xrightarrow{\mathcal{F}^c(s\bar{\mathcal{P}}, s\mu)} \mathcal{F}^c(s\bar{\mathcal{P}}) \xrightarrow{s\mu} s\bar{\mathcal{P}}$$

is zero. A map $l : \mathcal{S}^c(s\mathfrak{g}) \to s\mathfrak{g}$ induces a square-zero coderivation on $\mathcal{S}^c(s\mathfrak{g})$ means that the following composite is equal to zero

$$\mathcal{S}^c(s\mathfrak{g}) \xrightarrow{\delta'} \mathcal{S}^c(s\mathfrak{g}, \underbrace{\mathcal{S}^c(s\mathfrak{g})}_{1}) \xrightarrow{\mathcal{S}^c(s\mathfrak{g}, l)} \mathcal{S}^c(s\mathfrak{g}) \xrightarrow{l} s\mathfrak{g}.$$

We define the induced L_∞ structure by

$$l \ : \ \mathcal{S}^c(s(\oplus\bar{\mathcal{P}})) \xrightarrow{\Theta} \mathcal{F}^c(s\bar{\mathcal{P}}) \xrightarrow{s\mu} s\mathcal{P}.$$

The relation of the L_∞ structure for l lifts to the relation of the homotopy prop(erad) by the following commutative diagram :

$$\begin{tikzcd}
\mathcal{S}^c(s(\oplus\bar{\mathcal{P}})) \arrow[r, "\delta'"] \arrow[d, "\Theta"] & \mathcal{S}^c(s(\oplus\bar{\mathcal{P}}), \underbrace{\mathcal{S}^c(s(\oplus\bar{\mathcal{P}}))}_{1}) \arrow[r, "\mathcal{S}^c(s\bar{\mathcal{P}},l)"] & \mathcal{S}^c(s(\oplus\bar{\mathcal{P}})) \arrow[r, "l"] & s(\oplus\bar{\mathcal{P}}) \\
\mathcal{F}^c(s\bar{\mathcal{P}}) \arrow[r, "\Delta'"] & \mathcal{F}^c(s\bar{\mathcal{P}}, \underbrace{\mathcal{F}^c(s\bar{\mathcal{P}})}_{1}) \arrow[r, "\mathcal{F}^c(s\bar{\mathcal{P}},s\mu)"] & \mathcal{F}^c(s\bar{\mathcal{P}})\ , \arrow[ur, "s\mu"] &
\end{tikzcd}$$

which concludes the proof. ☐

When $\mathcal{P}$ is a (strict) prop(erad), the induced structure is the (strict) Lie algebra coming from the anti-symmetrization of the Lie-admissible algebra of Proposition 4. Theorem 25 generalizes the well-know fact that a homotopy (associative) algebra is a homotopy Lie algebra by anti-symmetrization of the structure maps.

The same statement holds for the space of coinvariants elements and the space of invariant elements.

Theorem 26. *Let $\mathcal{P}$ be a homotopy properad, the total space of coinvariant elements $\oplus\mathcal{P}_{\mathbb{S}}$ and the the total space of invariant elements $\oplus\mathcal{P}^{\mathbb{S}}$ have an induced L_∞-structure.*

PROOF. We apply the same arguments as in the proof of Proposition 6. ☐

We prove below that the maps $\mathcal{P} \to \oplus\mathcal{P}$ and $\mathcal{P} \to \oplus\mathcal{P}^{\mathbb{S}}$ are functors for the category of the homotopy prop(erad)s to one of homotopy Lie algebras (see Proposition 34). The same result holds for non-symmetric homotopy properads as well.

4.6. **Homotopy convolution prop(erad).** In this section, we extend the definition of the convolution prop(erad) to the homotopy case.

Theorem 27. *When $(\mathcal{C}, \Delta)$ is a (non-symmetric) homotopy coprop(erad) and $(\mathcal{P}, \mu)$ is a (non-symmetric) prop(erad), the convolution prop(erad) $\mathcal{P}^{\mathcal{C}} = \mathrm{Hom}(\mathcal{C}, \mathcal{P})$ is a homotopy (non-symmetric) prop(erad).*
The same result holds when $\mathcal{C}$ is a (non-symmetric) coprop(erad) and $\mathcal{P}$ a homotopy (non-symmetric) prop(erad).

PROOF. To an element $\mathcal{G}(f_1, \ldots, f_n)$ of $\mathcal{F}^c(\bar{\mathcal{P}}^{\overline{\mathcal{C}}})^{(n)}$, we consider the map $\widetilde{\mathcal{G}}(f_1, \ldots, f_n) : \mathcal{F}(\overline{\mathcal{C}})^{(n)} \to \mathcal{F}^c(\bar{\mathcal{P}})^{(n)}$ defined by $\mathcal{G}'(c_1, \ldots, c_n) \mapsto (-1)^{\xi}\mathcal{G}(f_1(c_1), \ldots, f_n(c_n))$ if $\mathcal{G}' \cong \mathcal{G}$ and 0 otherwise, where $\xi = \sum_{i=2}^{n} |f_i|(|c_1| + \cdots + |c_{i-1}|)$. We define maps $\mu_n : \mathcal{F}^c(\bar{\mathcal{P}}^{\overline{\mathcal{C}}})^{(n)} \to \mathcal{P}^{\mathcal{C}}$ by the formula

$$\mu_n\big(\mathcal{G}(f_1, \ldots, f_n)\big) := \widetilde{\mu}_{\mathcal{P}} \circ \widetilde{\mathcal{G}}(f_1, \ldots, f_n) \circ \Delta_n.$$

The degree of Δ_n is $n-2$ and the degree of $\widetilde{\mu}_{\mathcal{P}}$ is zero. Therefore, the degree of μ_n is $n-2$.
The map μ verifies the relation of Proposition 23

$$\begin{aligned}
\sum_{\mathcal{G}' \subset \mathcal{G}} \pm\, \mu\big(\mathcal{G}/\mu\mathcal{G}'(f_1, \ldots, f_n)\big) \quad &= \quad \sum \pm\, \widetilde{\mu}_{\mathcal{P}} \circ \widetilde{\mathcal{G}/\mathcal{G}'}(f_1, \ldots, \mu_k(\mathcal{G}'(f_{i_1}, \ldots, f_{i_k})), \ldots, f_n) \circ \Delta_l \\
&= \quad \sum \pm\, \widetilde{\mu}_{\mathcal{P}} \circ \widetilde{\mathcal{G}/\mathcal{G}'}(f_1, \ldots, \widetilde{\mu}_{\mathcal{P}} \circ \widetilde{\mathcal{G}}'(f_{i_1}, \ldots, f_{i_k}) \circ \delta_k, \ldots, f_n) \circ \Delta_l,
\end{aligned}$$

where the sum runs over admissible subgraphs $\mathcal{G}'$ of $\mathcal{G}$. We denote by k the number of vertices of $\mathcal{G}'$ and $l = n-k+1$.We use the generic notation i for the new vertex of $\mathcal{G}/\mathcal{G}'$ obtained after contracting $\mathcal{G}'$. For every element $c \in \overline{\mathcal{C}}$, we denote by $\Delta(c) = \sum \mathcal{G}^1(c_1, \ldots, c_l)$ and $\Delta(c_i) = \sum \mathcal{G}_i^2(c'_1, \ldots, c'_k)$. The associativity of the product of $\mathcal{P}$ gives

$$\begin{aligned}
&\sum_{\mathcal{G}' \subset \mathcal{G}} (-1)^{\varepsilon(\mathcal{G}', f_1, \ldots, f_n)}\, \mu\big(\mathcal{G}/\mu\mathcal{G}'(f_1, \ldots, f_n)\big)(c) = \\
&\widetilde{\mu}_{\mathcal{P}} \circ \widetilde{\mathcal{G}}(f_1, \ldots, f_n) \circ \Big(\sum (-1)^{\rho(\mathcal{G}_i^2, c_1, \ldots, c_l)}\, \mathcal{G}^1 \circ_i \mathcal{G}_i^2(c_1, \ldots, c'_1, \ldots, c'_k, \ldots, c_l)\Big).
\end{aligned}$$

Since $(\mathcal{C}, \Delta)$ is a homotopy coprop(erad), the last term vanishes by Proposition 24.
The same statement in the non-symmetric case is proven in the same way and the dual statement also. ☐

REMARK. In the particular case when $\mathcal{C}$ is a homotopy coalgebra and $\mathcal{P}$ an associative algebra, $\mathrm{Hom}(\mathcal{C}, \mathcal{P})$ is a homotopy algebra. In the same way, when $\mathcal{C}$ is a homotopy operad and $\mathcal{P}$ an operad, $\mathrm{Hom}(\mathcal{C}, \mathcal{P})$ is an homotopy operad (see Lemma 5.10 of [VdL02]).

Theorem 28. *When $(\mathcal{C}, \Delta)$ is a homotopy coprop(erad) and $(\mathcal{P}, \mu)$ is a prop(erad) (or when $(\mathcal{C}, \Delta)$ is a coprop(erad) and $(\mathcal{P}, \mu)$ is a homotopy prop(erad)), the total space of the convolution prop(erad) $\mathcal{P}^{\mathcal{C}} = \mathrm{Hom}(\mathcal{C}, \mathcal{P})$ is a homotopy Lie algebras.*
The total subspace $\mathrm{Hom}^{\mathbb{S}}(\mathcal{C}, \mathcal{P})$ of invariant elements is a sub-L_∞-algebra.

PROOF. The first part is a direct corollary of Theorem27 and Theorem25. Since the structure maps of this Ł$_\infty$-algebra are composite of equivariant maps $(\Delta_n,\ \widetilde{\mu}_{\mathcal{P}})$, the induce an L_∞-algebra structure on the total space of $\mathrm{Hom}^{\mathbb{S}}(\mathcal{C}, \mathcal{P})$. (This is similar to the one used in the proof of Proposition 11). ☐

In the latter case, the L_∞-'operations' or homotopies are explicitly given by the following formula. The image of $f_1, \ldots, f_n \in \mathrm{Hom}^{\mathbb{S}}(\mathcal{C}, \mathcal{P})$ under l_n, for $n > 1$, is given by

$$l_n(f_1, \ldots, f_n) = \sum_{\sigma \in \mathbb{S}_n} (-1)^{\mathrm{sgn}(\sigma, f_1, \ldots, f_n)} \widetilde{\mu}_{\mathcal{P}} \circ (f_{\sigma(1)} \otimes \cdots \otimes f_{\sigma(n)}) \circ \Delta_n,$$

where $(-1)^{\mathrm{sgn}(\sigma, f_1, \ldots, f_n)}$ is the Koszul-Quillen sign appearing after permutating the f_i with σ. The first 'operation' l_1 is the differential, that is $l_1(f) := D(f) = d_{\mathcal{P}} \circ f - (-1)^{|f|} f \circ d_{\mathcal{C}}$.

In this homotopy Lie algebra, the generalized Maurer-Cartan equation is well defined since the formal infinite sum $Q(\alpha) := \sum_{n \geq 1} \frac{1}{n!} l_n(\alpha, \ldots, \alpha)$ is in fact equal to the composite $D + \widetilde{\mu}_{\mathcal{P}} \circ \mathcal{F}(\alpha) \circ \Delta$ in $\mathrm{Hom}(\mathcal{C}, \mathcal{P})$, when $\mathcal{C}$ is a homotopy coprop(erad) and to $D + \mu \circ \mathcal{F}(\alpha) \circ \widetilde{\Delta}_{\mathcal{C}}$ when $\mathcal{P}$ is a homotopy prop(erad). (See 7.3 for the general definition of filtered L_∞-algebras).

Definition. Let $(\mathcal{C}, \Delta)$ be a homotopy coprop(erad) and $(\mathcal{P}, \mu)$ be a prop(erad) (or $(\mathcal{C}, \Delta)$ a coprop(erad) and $(\mathcal{P}, \mu)$ a homotopy properad). A morphism $\mathcal{C} \xrightarrow{\alpha} \mathcal{P}$, of degree -1, is called a *twisting morphism* if it is a solution of the (generalized) Maurer-Cartan equation

$$Q(\alpha) := \sum_{n \geq 1} \frac{1}{n!} l_n(\alpha, \ldots, \alpha) = 0,$$

in the homotopy Lie algebra $\mathrm{Hom}^{\mathbb{S}}(\mathcal{C}, \mathcal{P})$. We denote this set by $\mathrm{Tw}(\mathcal{C}, \mathcal{P})$.

We can represent the bifunctor $\mathrm{Tw}(-, -)$ in the same as in the strict case (see Proposition 17).

Proposition 29. *Let $(\mathcal{C}, \Delta)$ be a homotopy coprop(erad) and $(\mathcal{P}, \mu)$ be a prop(erad). There is a natural bijection*

$$\mathrm{Mor}_{dg\ prop(erad)s}\left(\Omega(\mathcal{C}),\ \mathcal{P}\right) \cong \mathrm{Tw}(\mathcal{C}, \mathcal{P}).$$

Let $(\mathcal{C}, \Delta)$ be a coprop(erad) and $(\mathcal{P}, \mu)$ be a homotopy prop(erad). There is a natural bijection

$$\mathrm{Tw}(\mathcal{C}, \mathcal{P}) \cong \mathrm{Mor}_{dg\ coprop(erad)s}\left(\mathcal{C},\ \mathcal{B}(\mathcal{P})\right).$$

PROOF. The proof is a direct generalization of the proof of Proposition 17. ☐

4.7. **Morphisms of homotopy (co)prop(erad)s.** In this section, we recall the notion of morphism between two homotopy properads due to [Gra07]. We extend it to homotopy (co)props and make them explicit in terms of Maurer-Cartan elements in some convolution L_∞-algebra.

Since a homotopy properad is equivalent to its associated (generalized) bar construction, the notion of *morphism of homotopy properads* (or *weak morphism*) is defined as follows.

Definition. [Gra07] Let $\mathcal{P}_1$ and $\mathcal{P}_2$ be two homotopy prop(erad)s. A morphism between $\mathcal{P}_1$ and $\mathcal{P}_2$ is a morphism of dg coprop(erad)s between their bar constructions : $B(\mathcal{P}_1) \to B(\mathcal{P}_2)$.

A morphism of dg coprop(erad)s $\Phi : B(\mathcal{P}_1) = \mathcal{F}^c(s\bar{\mathcal{P}}_1) \to B(\mathcal{P}_2) = \mathcal{F}^c(s\bar{\mathcal{P}}_2)$ is characterized by its image on $s\bar{\mathcal{P}}_2$. We denote by $s^{-1}\varphi : B(\mathcal{P}_1) \to \bar{\mathcal{P}}_2$ the composite of Φ with the projection on $s\bar{\mathcal{P}}_2$ followed by the desuspension. Notice that the degree of $s^{-1}\varphi$ is -1. By Proposition 29, Φ is a morphism of dg coprop(erad)s if and only if $s^{-1}\varphi$ is a Maurer-Cartan element in $\mathrm{Hom}^{\mathbb{S}}(B(\mathcal{P}_1), \mathcal{P}_2)$, that is

$$Q(s^{-1}\varphi) = \sum_{n\geq 1} \frac{1}{n!}\, l_n(s^{-1}\varphi, \ldots, s^{-1}\varphi) = D(s^{-1}\varphi) + \mu_{\mathcal{P}_2} \circ \mathcal{F}^c(s^{-1}\varphi) \circ \widetilde{\Delta} = 0,$$

where $\widetilde{\Delta}$ is the coproduct map $B(\mathcal{P}_1) = \mathcal{F}^c(s\bar{\mathcal{P}}_1) \to \mathcal{F}^c(\mathcal{F}^c(s\bar{\mathcal{P}}_1))$.

Proposition 30. *A morphism of $\mathbb{S}$-bimodules $\varphi : B(\mathcal{P}_1) \to s\bar{\mathcal{P}}_2$ induces a morphism of homotopy properads between $\mathcal{P}_1$ and $\mathcal{P}_2$ if and only if $s^{-1}\varphi$ is a Maurer-Cartan element in the L_∞-algebra* $\mathrm{Hom}^{\mathbb{S}}(B(\mathcal{P}_1), \mathcal{P}_2)$*, that is $Q(s^{-1}\varphi) = 0$.*

Like in Section 4.3, we make explicit the above definition in terms of graphs.

Proposition 31. *A map $s^{-1}\varphi : B(\mathcal{P}_1) \to \bar{\mathcal{P}}_2$ is a morphism of homotopy prop(erad)s if and only if, for every class of graphs $\mathcal{G}$ under the action of the automorphism group, the following relation holds*

$$\sum s\mu_k^{\mathcal{P}_2}\big(\mathcal{G}/\varphi\mathcal{G}_1 \sqcup \ldots \sqcup \varphi\mathcal{G}_k\big) = \sum \varphi\big(\mathcal{G}/(s\mu^{\mathcal{P}_1})\mathcal{G}'\big),$$

where the first sum runs over all partition of the graph $\mathcal{G}$ into admissible subgraphs $\mathcal{G}_1 \sqcup \ldots \sqcup \mathcal{G}_k$ and where the second sum runs over all admissible subgraph $\mathcal{G}'$ of $\mathcal{G}$. Once again, the signs are induced by Koszul-Quillen rule, when apply to elements $sp_1, \ldots, sp_n$, such that n is the number of vertices of $\mathcal{G}$.

Proof. The map $s^{-1}\varphi : B(\mathcal{P}_1) \to \bar{\mathcal{P}}_2$ induces a unique morphism of coprop(erad)s $\Phi : B(\mathcal{P}_1) \to B(\mathcal{P}_2)$ which commutes with the differentials if and only if the above relation is verified. (The left hand term is the projection on $\bar{\mathcal{P}}_2$ of the composite $d_{B(\mathcal{P}_2)} \circ \Phi$ and the right hand term is the projection on the same space of the composite $\Phi \circ d_{B(\mathcal{P}_1)}$, that is $\varphi \circ d_{B(\mathcal{P}_1)}$.)

□

When applied to A_∞-algebras, the underlying graphs are ladders and this proposition gives the classical notion of weak morphisms, that is morphisms between A_∞-algebras.

Dually, we define the notion of morphism between homotopy coprop(erad)s.

Definition. Let $\mathcal{C}_1$ and $\mathcal{C}_2$ be two homotopy prop(erad)s. A morphism between $\mathcal{C}_1$ and $\mathcal{C}_2$ is a morphism of dg prop(erad)s between their cobar constructions : $\Omega(\mathcal{C}_1) \to \Omega(\mathcal{C}_2)$.

A morphism of dg prop(erad)s $\Psi : \Omega(\mathcal{C}_1) = \mathcal{F}(s^{-1}\bar{\mathcal{C}}_1) \to \Omega(\mathcal{C}_2) = \mathcal{F}(s^{-1}\bar{\mathcal{C}}_2)$ is characterized by the image of $s^{-1}\bar{\mathcal{C}}_1$. We denote by $s^{-1}\psi : \bar{\mathcal{C}}_1 \to \Omega(\mathcal{C}_2)$ the desuspension of the restriction of Ψ on $s^{-1}\bar{\mathcal{C}}_1$. By Proposition 29, Ψ is a morphism of dg prop(erad)s if and only if $s^{-1}\psi$ is a twisting morphism in $\mathrm{Hom}^{\mathbb{S}}(\bar{\mathcal{C}}_1, \Omega(\mathcal{C}_2))$, that is

$$Q(s^{-1}\psi) = \sum_{n\geq 1} \frac{1}{n!}\, l_n(s^{-1}\psi, \ldots, s^{-1}\psi) = D(s^{-1}\psi) + \widetilde{\mu} \circ \mathcal{F}(s^{-1}\psi) \circ \Delta_{\mathcal{C}_1} = 0,$$

where $\widetilde{\mu}$ is the composition map $\mathcal{F}(\Omega(\mathcal{C}_2)) = \mathcal{F}(\mathcal{F}(s^{-1}\bar{\mathcal{C}}_2)) \to \mathcal{F}(s^{-1}\bar{\mathcal{C}}_2) = \Omega(\mathcal{C}_2)$.

Proposition 32. *A morphism of $\mathbb{S}$-bimodules $\psi : s^{-1}\bar{\mathcal{C}}_1 \to \Omega(\mathcal{C}_2)$ induces a morphism of homotopy coproperads between $\mathcal{C}_1$ and $\mathcal{C}_2$ if and only if $s^{-1}\psi$ is a Maurer-Cartan element in the L_∞-algebra* $\mathrm{Hom}^{\mathbb{S}}(\mathcal{C}_1, \Omega(\mathcal{C}_2))$*, that is $Q(s^{-1}\psi) = 0$.*

We now prove that the convolution prop(erad) is a construction functorial with respect to the first argument.

Theorem 33. *Let Ψ be a morphism of homotopy coprop(erad)s between $\mathcal{C}_1$ and $\mathcal{C}_2$. Let $\mathcal{P}$ be a prop(erad). There exists a natural morphism of homotopy prop(erad)s between* $\mathrm{Hom}(\mathcal{C}_2, \mathcal{P})$ *and* $\mathrm{Hom}(\mathcal{C}_1, \mathcal{P})$ *induced by Ψ.*

The same statement holds in the non-symmetric case.

Proof. Let Ψ denote the morphism of dg prop(erads) $\Omega(\mathcal{C}_1) \to \Omega(\mathcal{C}_1)$ and $s^{-1}\psi$ the induced twisting morphism $\bar{\mathcal{C}}_1 \to \Omega(\mathcal{C}_2)$, that is $Q(s^{-1}\psi) = 0$. We define the morphism of coprop(erad)s $\Phi \ : \ B(\mathrm{Hom}(\mathcal{C}_2, \mathcal{P})) \to B(\mathrm{Hom}(\mathcal{C}_1, \mathcal{P}))$ by its image φ on $s\,\overline{\mathrm{Hom}}(\mathcal{C}_1, \mathcal{P}) = \mathrm{Hom}(s^{-1}\bar{\mathcal{C}}_1, \bar{\mathcal{P}})$ as follows. Let $\mathcal{G}(f_1, \ldots, f_n) \in B(\mathrm{Hom}(\mathcal{C}_2, \mathcal{P})) = \mathcal{F}^c(s\,\overline{\mathrm{Hom}}(\mathcal{C}_1, \mathcal{P})) = \mathcal{F}^c(\mathrm{Hom}(s^{-1}\bar{\mathcal{C}}_1, \bar{\mathcal{P}}))$. The image of $\mathcal{G}(f_1, \ldots, f_n)$ under φ is equal to the composite

$$\varphi(\mathcal{G}(f_1, \ldots, f_n)) \ : \ s^{-1}\bar{\mathcal{C}}_1 \xrightarrow{\psi} \mathcal{F}(s^{-1}\bar{\mathcal{C}}_2) \xrightarrow{\widetilde{\mathcal{G}}(f_1, \ldots, f_n)} \mathcal{F}(\bar{\mathcal{P}}) \xrightarrow{\widetilde{\mu}_{\mathcal{P}}} \mathcal{P}.$$

It remains to prove that $s^{-1}\varphi$ is a twisting element in $\mathrm{Hom}(B(\mathrm{Hom}(\mathcal{C}_2, \mathcal{P})), \mathrm{Hom}(\mathcal{C}_1, \mathcal{P}))$, that is $Q(s^{-1}\varphi) = 0$. By the definition of Q in this homotopy prop(erad) and by the 'associativity' of $\widetilde{\mu}_{\mathcal{P}}$, $Q(s^{-1}\varphi)(\mathcal{G}(f_1, \ldots, f_n))$ is equal to the composite

$$\bar{\mathcal{C}}_1 \xrightarrow{\Delta_{\mathcal{C}_1}} \mathcal{F}(\bar{\mathcal{C}}_1) \xrightarrow{\mathcal{F}(s^{-1}\psi)} \mathcal{F}(\mathcal{F}(s^{-1}\bar{\mathcal{C}}_2)) \xrightarrow{\widetilde{\mu}} \mathcal{F}(s^{-1}\bar{\mathcal{C}}_2) \xrightarrow{\widetilde{\mathcal{G}}(f_1, \ldots, f_n)} \mathcal{F}(\bar{\mathcal{P}}) \xrightarrow{\widetilde{\mu}_{\mathcal{P}}} \mathcal{P},$$

where $\widetilde{\mu}$ is the 'triple' map associated to the free prop(erad) $\mathcal{F}(s^{-1}\bar{\mathcal{C}}_2)$. Therefore $Q(s^{-1}\varphi)(\mathcal{G}(f_1, \ldots, f_n)) = \widetilde{\mu}_{\mathcal{P}} \circ \widetilde{\mathcal{G}}(f_1, \ldots, f_n) \circ Q(s^{-1}\psi)$ which vanishes since $Q(s^{-1}\psi) = 0$. ☐

The dual statement is also true and can be proved in the same way. It will appear in a future work of the second author in relation with the the transfer of algebraic structures up to homotopy through a deformation-retract (homological perturbation lemma).

Proposition 34. *The constructions given in Theorem 25 and Theorem 26 provide us with three functors,*

$$\textsf{Category of homotopy properads} \longrightarrow \textsf{Category of homotopy Lie algebras.}$$

Proof. Let $\Phi \ : \ B(\mathcal{P}_1) \to B(\mathcal{P}_2)$ be a morphism of coprop(erad)s defining a morphism of homotopy prop(erad)s between $\mathcal{P}_1$ and $\mathcal{P}_2$. The associated projection φ verifies $Q(s^{-1}\varphi) = 0$, that is

$$\mathcal{F}^c(s\bar{\mathcal{P}}_1) \xrightarrow{\widetilde{\Delta}} \mathcal{F}^c(\mathcal{F}^c(s\bar{\mathcal{P}}_1)) \xrightarrow{\mathcal{F}^c(s^{-1}\varphi)} \mathcal{F}^c(\bar{\mathcal{P}}_2) \xrightarrow{\mu_{\mathcal{P}_2}} \mathcal{P}_2$$

equals 0. We define the following map

$$f \ : \ \mathcal{S}^c(s(\oplus\bar{\mathcal{P}}_1)) \xrightarrow{\Theta} \mathcal{F}^c(s\bar{\mathcal{P}}_1) \xrightarrow{\varphi} s(\oplus\bar{\mathcal{P}}_2).$$

The map f is a morphism of L_∞-algebras. Its desuspension $s^{-1}f$ verifies the Maurer-Cartan equation in the L_∞-algebra $\mathrm{Hom}(\mathcal{S}^c(s(\oplus\mathcal{P}_1)), \oplus\mathcal{P}_2)$ (see [Dol07]). The Maurer-Cartan equation for $s^{-1}f$ lifts to the Maurer-Cartan equation for $s^{-1}\varphi$ via Θ, that is the following diagram is commutative

$$\begin{tikzcd}
\mathcal{S}^c(s(\oplus\bar{\mathcal{P}}_1)) \arrow[r, "\widetilde{\delta}"] \arrow[d, "\Theta"] & \mathcal{S}^c(\mathcal{S}^c(s(\oplus\bar{\mathcal{P}}_1))) \arrow[r, "\mathcal{S}^c(s^{-1}f)"] & \mathcal{S}^c(\oplus\bar{\mathcal{P}}_2) \arrow[r, "l_{\oplus\mathcal{P}_2}"] & \oplus\bar{\mathcal{P}}_2 \\
\mathcal{F}^c(s\bar{\mathcal{P}}_1) \arrow[r, "\widetilde{\Delta}"] & \mathcal{F}^c(\mathcal{F}^c(s\bar{\mathcal{P}}_1)) \arrow[r, "\mathcal{F}^c(s^{-1}\varphi)"] & \mathcal{F}^c(\bar{\mathcal{P}}_2) \, , \arrow[ur, "\mu_{\mathcal{P}_2}"] &
\end{tikzcd}$$

which concludes the proof. ☐

Corollary 35. *Let Ψ be a morphism of homotopy coprop(erad)s between $\mathcal{C}_1$ and $\mathcal{C}_2$. Let $\mathcal{P}$ be a prop(erad). There exists a natural morphism of L_∞-algebras between* $\mathrm{Hom}(\mathcal{C}_2, \mathcal{P})$ *and* $\mathrm{Hom}(\mathcal{C}_1, \mathcal{P})$ *induced by Ψ. Its restriction to* $\mathrm{Hom}^{\mathbb{S}}(\mathcal{C}_2, \mathcal{P})$ *gives a natural morphism of L_∞-algebras between* $\mathrm{Hom}^{\mathbb{S}}(\mathcal{C}_2, \mathcal{P})$ *and* $\mathrm{Hom}^{\mathbb{S}}(\mathcal{C}_1, \mathcal{P})$.

Proof. The first part is a direct corollary of Theorem 33 and Proposition 34. Since these constructions are composite of equivariant maps, they are stable on the space of invariant elements $\mathrm{Hom}^{\mathbb{S}}(\mathcal{C}_2, \mathcal{P})$ and $\mathrm{Hom}^{\mathbb{S}}(\mathcal{C}_1, \mathcal{P})$. ☐

5. Models

In this section, we recall the definitions of *minimal* and *quadratic model* for properads and we formally extend them to props. Recall that a model is a quasi-free resolution. Our viewpoint here is to classify properads according to the form of their minimal model, when it exists. For instance, a properad is *Koszul* if and only if it admits a quadratic model. To clarify the genus of some resolutions, we introduce the notion of *contractible prop(erad)s.* Such properads have genus 0 quadratic models.

5.1. **Minimal models.** Recall that a *quasi-free* prop(erad) is a (dg) prop(erad) whose underlying $\mathbb{S}$-bimodule, that is forgetting the differential map, is a free prop(erad) $\mathcal{F}(M)$. It is not necessarily a free dg prop(erad) since the differential ∂ may not be freely generated by the differential of M.

Definition (Model)**.** Let $\mathcal{P}$ be a prop(erad). A *model* of $\mathcal{P}$ is a quasi-free prop(erad) $(\mathcal{F}(M),\, \partial)$ equipped with a quasi-isomorphism $\mathcal{F}(M) \xrightarrow{\sim} \mathcal{P}$.

Theorem 19 proves that every augmented prop(erad) has a canonical model given by the bar-cobar construction. Some prop(erad)s admit more simple models. The differential ∂ of a quasi-free prop(erad) $\mathcal{F}(M)$ is by definition a derivation. Lemma 14 shows that it is characterized by its restriction $\partial_M \,:\, M \to \mathcal{F}(M)$ on M.

Definition (Decomposable differential)**.** The differential ∂ of a quasi-free prop(erad) is called *decomposable* if the image of its restriction to M, $\partial_M \,:\, M \to \mathcal{F}(M)$, is composed by decomposable elements, that is $\mathrm{Im}(\partial_M) \subset \bigoplus_{n\geq 2} \mathcal{F}(M)^{(n)}$.

Definition (Minimal model)**.** A model $(\mathcal{F}(M),\, \partial)$ is called *minimal* if its differential ∂ is decomposable.

5.2. **Form of minimal models.** From Theorem 19, we know that every augmented (dg) properad admits a resolution of the form $\Omega(B(P))$. A natural way to get a minimal model from this would be to consider the homology of the bar construction, try to endow it with a structure of homotopy coproperad and then take the generalized cobar construction of it. In this section, we prove that when minimal models exist, they are of this form.

Proposition 36. *Let $(\mathcal{F}(M),\, \partial)$ be a quasi-free properad with a decomposable differential generated by a non-negatively graded $\mathbb{S}$-module M. Then the homology of the bar construction $B(\mathcal{F}(M))$ of $(\mathcal{F}(M),\, \partial)$ is equal to the suspension of M.*

Proof. The bar construction of the dg-properad $\mathcal{P} := \mathcal{F}(M)$ is defined by the underlying $\mathbb{S}$-bimodule $B(\mathcal{P}) := \mathcal{F}^c(s\bar{\mathcal{P}}) = \mathcal{F}^c(s\,\bar{\mathcal{F}}(M))$. The differential d is the sum of two terms $d_0 + \tilde{\partial}$. The component $\tilde{\partial}$ comes from ∂ and d_0 is the unique coderivation which extends the partial product of $\mathcal{F}(M)$.
Consider the filtration $F_s := \bigoplus_{r\leq s} \mathcal{F}^c\big(s\bar{\mathcal{F}}(M)\big)_r$, where r is the sum of the degrees of the elements of M. Let's denote by $E^\bullet_{st}$ the associated spectral sequence.
Since the chain complex M is bounded below, this filtration is bounded below $F_{-1} = 0$. It is obviously exhaustive, therefore the classical theorem of convergence of spectral sequences shows that $E^\bullet$ converges to the homology of $B(\mathcal{F}(M))$.
We have $\tilde{\partial}(F_s) \subset F_{s-1}$ and $d_0(F_s) \subset F_s$. Hence, the first term of the spectral sequence is $E^0_{st} = \mathcal{F}^c_{s+t}\big(s\bar{\mathcal{F}}(M)\big)_s$, where $s+t$ is the total homological degree, and $d^0 = d_0$. We have reduced the problem to computing the homology of the bar construction of the free properad on M, which is equal to ΣM by Corollary 5.10 of [Val07a] (where we choose to put each element of M in weight 1). ☐

The next proposition shows that, when a minimal model of a properad $\mathcal{P}$ exists, it is necessarily given by a quasi-free properad on the homology of the bar construction of $\mathcal{P}$.

Theorem 37. *Let $\mathcal{P}$ be an augmented dg properad and let $(\mathcal{F}(M),\, \partial)$ be a minimal model of $\mathcal{P}$. The $\mathbb{S}$-bimodule sM is isomorphic to the homology of the bar construction of $\mathcal{P}$.*

Proof. In [Val07a], we proved in Proposition 4.9 that the bar construction preserves quasi-isomorphisms. Therefore, the bar construction of $\mathcal{F}(M)$ is quasi-isomorphic to the bar construction of $\mathcal{P}$. We conclude by Proposition 36. $\square$

We denote by $\mathcal{P}^{¡} := H_\bullet(B(\mathcal{P}))$ the homology of the bar construction of $\mathcal{P}$. When $(\mathcal{F}(s^{-1}\mathcal{P}^{¡}),\, \partial)$ is a minimal model of $\mathcal{P}$, the derivation ∂ is equivalent to a structure of homotopy coproperad on $\mathcal{P}^{¡}$ such that $\delta_1 = 0$. That is $(\mathcal{F}(s^{-1}\mathcal{P}^{¡}),\, \partial)$ is the generalized cobar construction $\Omega(\mathcal{P}^{¡})$ of the homotopy coproperad $\mathcal{P}^{¡}$. As a conclusion, we have that following corollary which gives the form of minimal models.

Corollary 38. *A minimal model of an augmented dg-properad $\mathcal{P}$ is always the cobar construction $\Omega(\mathcal{P}^{i})$ on the homology of $B(\mathcal{P})$ endowed with a structure of homotopy coproperad.*

In the sequel, we will only consider props freely generated by a properad, in the sense of the horizontal (concatenation) product. The minimal model of such props is given by the generalized cobar construction of the associated homotopy coproperad, viewed as a homotopy coprop. And the result of the preceding lemma still holds.

5.3. **Quadratic models and Koszul duality theory.** In general, it is a difficult problem to find the minimal model of a prop(erad). One can first compute the homology of the bar construction and then provide a structure of homotopy coproperad on it, that is with higher homotopy cooperations. For some weight graded properads, there exist simple minimal models which are given by the Koszul duality theory. These properads are called Koszul.

Definition (Quadratic differential)**.** The differential ∂ of a quasi-free prop(erad) is called *quadratic* if the image of $\partial_M\, :\, M \to \mathcal{F}(M)$ is in $\mathcal{F}(M)^{(2)}$.

Definition (Quadratic model)**.** A model $(\mathcal{F}(M),\, \partial)$ is called *quadratic* if its differential ∂ is quadratic.

When $\mathcal{P}$ is a weight graded properad, its bar construction splits as a direct sum of finite chain complexes indexed by the weight (cf. [Val07a] Section 7.1.1). In this case, we can talk about top dimensional homology groups.

Theorem 39. *Let $\mathcal{P}$ be a weight graded properad concentrated in homological degree* 0*. The following assertions are equivalent.*

1. *The homology of $B(\mathcal{P})$ is concentrated in top dimension.*
2. *The $\mathbb{S}$-bimodule $\mathcal{P}^{i}$ is a strict coproperad.*
3. *The properad $\mathcal{P}$ admits a quadratic model : $\Omega(\mathcal{P}^{i}) \xrightarrow{\sim} \mathcal{P}$.*

Proof. (1) ⇒ (2) is given by Proposition 7.2 of [Val07a].
(2) ⇒ (3) is given by Theorem 5.9 of [Val07a]. When $\mathcal{P}^{¡}$ has a structure of strict coproperad, its cobar construction is a resolution of $\mathcal{P}$ and the differential of it is quadratic.
(3) ⇒ (1) Since $\mathcal{P}$ is isomorphic to $\mathcal{F}(M_0)/(\partial(M_1))$, which ∂ quadratic, this presentation is quadratic. Define an extra weight on M by the formula $\omega(M_n) := n+1$. With this weight, the quasi-isomorphism $\mathcal{F}(M) \xrightarrow{\rho} \mathcal{P}$ is a morphism of weight graded dg-properads. The induced morphism $B(\rho)$ on the bar construction preserves this grading. Therefore we have $H_n(B(\mathcal{P})^{(n)}) = H_n(B(\mathcal{F}(M))^{(n)}) = (sM)_n$ and the homology of the bar construction of $\mathcal{P}$ is concentrated in top dimension. $\square$

In this case, the properad $\mathcal{P}$ is called a *Koszul* properad. The coproperad $\mathcal{P}^{¡}$ is its *Koszul dual* and $\mathcal{P}$ has a quadratic model which is the cobar construction on $\mathcal{P}^{¡}$. In other words, a properad is Koszul when its bar construction is *formal*, that is when $B(\mathcal{P})$ is quasi-isomorphic to its homology $\mathcal{P}^{¡}$ as a dg-coproperad. This case is simple and particularly efficient. When $\mathcal{P} = F(V)/(R)$ has a quadratic presentation with a finite dimensional space of generators V, then the linear dual (twisted by the signature representation) of the coproperad $\mathcal{P}^{¡}$ is a properad equal, up to suspension, to $\mathcal{P}^! = F(V^\vee)/(\mathcal{R}^\perp)$ where $V^\vee$ is the linear dual of V twisted by the signature representation. This relation provides a concrete method to compute the minimal model of Koszul properads. The next step is to be able to prove that it is Koszul. Koszul duality theory provides a smaller chain complex $\mathcal{P}^{¡} \boxtimes \mathcal{P}$ which is acyclic if and only if the properad $\mathcal{P}$ is Koszul. Therefore, there are simple methods to show that a properad is Koszul. When a properad is defined by two Koszul properads with a distributive law, Proposition 8.4 of [Val07a] shows that it is Koszul. In the operadic case, there are basically two other efficient methods. First if the homology of the free $\mathcal{P}$-algebra is acyclic then the operad $\mathcal{P}$ is Koszul (see Proposition 5.3.5 of [Fre04]). Finally, when the operad is set theoretic, we can use the associated poset to prove that it is Koszul (see [Val06a]).

5.4. **Homotopy Koszul properads.** If a properad is Koszul, then we have clearly cut means to construct its minimal model. However, the ordinary notion of Koszulness does not cover many important examples. For example, the properad of associative bialgebras is not Koszul since it is not quadratic and any Koszul properad has a quadratic presentation by Corollary 7.5 of [Val07a]. So we are left in such cases with no concrete methods of proving that a particular properad $\mathcal{P}$ admits a minimal model, and, if so, constructing it explicitly. It is already a highly non-trivial problem in general to find the set of generators for a minimal model, not speaking about the differential. In this section we extend the notion of Koszulness in such a way that some of the above problems become effectively solvable.

Definition. Let $\mathcal{P} = \mathcal{F}(V)/(\mathcal{R})$ be a properad generated by an $\mathbb{S}$-bimodule $V = \{V(m,n)\}$ concentrated in degree zero, and with an ideal generated by $\mathcal{R} \subset \mathcal{F}(V)^{(\geq 2)}$. Let $\pi_k : \mathcal{F}(V) \to \mathcal{F}(V)^{(k)}$ be the natural projection, and let us set,

$$\mathcal{R}_k := \pi_k(\mathcal{R}), \quad \text{for } k = 2, 3, \ldots.$$

Let us also denote by $\mathcal{P}^{(\geq k)}$ the image of $\mathcal{F}(V)^{(\geq k)}$ under the natural epimorphism $\mathcal{F}(V) \twoheadrightarrow \mathcal{P}$. The properad $\mathcal{P}$ is called *homotopy Koszul* if

(i) the quadratic properad $\mathcal{P}_2 := \mathcal{F}(V)/(\mathcal{R}_2)$ is Koszul,
(ii) $\mathcal{P}$ and $\mathcal{P}_2$ are isomorphic as $\mathbb{S}$-bimodules,
(iii) there is an extra grading on the properad $\mathcal{P} = \oplus_\lambda \mathcal{P}(\lambda)$, with $\mathcal{P}(\lambda)$ being a collection of finite-dimensional $\mathbb{S}$-bimodules.

In practice the conditions (i)-(iii) above are often not hard to check (see examples below). As an extra grading one can use, for example, the path grading of a free properad introduced by Kontsevich and studied in [MV03]. The main motivation behind the definition is the following.

Theorem 40. *If a properad $\mathcal{P}$ is homotopy Koszul, then it admits a minimal model of the form $(\mathcal{F}(s\bar{\mathcal{P}}_2^{¡}), \delta)$, where $\mathcal{P}_2^{¡}$ is the coproperad Koszul dual to $\mathcal{P}_2$.*

Proof. Consider the bounded above increasing filtration $F_{-p}\mathcal{P} := \mathcal{P}^{(\geq p)}$ of the properad $\mathcal{P}$. As $F_{-p}\mathcal{P} \cap P(\lambda)$ are finite-dimensional vector spaces, the spectral sequences associated with this filtration (see below) have good convergence properties. Since $\mathcal{P}$ is isomorphic to $\mathcal{P}_2$ as an $\mathbb{S}$-bimodule, the associated graded properad,

$$\bigoplus_{p \geq 0} \frac{\mathcal{P}^{(\geq p)}}{\mathcal{P}^{(\geq p+1)}},$$

is isomorphic to $\mathcal{P}_2$ as a properad. Then we have,

Claim 1. *The homologies of the bar constructions, $B(\mathcal{P})$ and $B(\mathcal{P}_2)$, are isomorphic as $\mathbb{S}$-bimodules, i.e. $H_\bullet(B(\mathcal{P})) \simeq \mathcal{P}_2^{¡}$ as $\mathbb{S}$-bimodules.*

Indeed, the filtration $F_{-p}\mathcal{P} := \mathcal{P}^{(\geq p)}$ induces an associated filtration of the complex $B(\mathcal{P})$ (as differential in $B(\mathcal{P})$ is built from compositions in $\mathcal{P}$ which respect the filtration $F_{-p}\mathcal{P}$). By the above observation, the 0th term, E^0, of the associated spectral sequence, $\{E^r, d^r\}$, is exactly the complex $B(\mathcal{P}_2)$, $E^0_{pq} = B(\mathcal{P}_2)^{(-p)}_{p+q}$ and $d^0 = d_{B(\mathcal{P}_2)}$. As $\mathcal{P}_2$ is Koszul, $E^1 = H_\bullet(B(\mathcal{P}_2))$ is exactly the Koszul dual coproperad $\mathcal{P}_2^{¡}$, that is $E^1_{pq} = 0$ for $q \neq -2p$ and $E^1_{pq} = H_{-p}\big(B(\mathcal{P}_2)^{(-p)}\big) = (\mathcal{P}_2^{¡})^{(-p)}$ when $q = -2p$. The induced differentials, d^r for $r \geq 1$, are zero because of the homological degree 0 assumption on $\mathcal{P}$. Thus the spectral sequence $\{E^r, d^r\}$ degenerates at the first term. The extra grading on the properad $\mathcal{P}$ induces an extra grading λ on $B(\mathcal{P})$ which makes $F_{-p}(B(\mathcal{P})) \cap$

$B(\mathcal{P})(\lambda)$ into a bounded filtration of $B(\mathcal{P})(\lambda)$. Hence it converges to $H_\bullet(B(\mathcal{P}))(\lambda)$ by the Classical Convergence Theorem 5.5.1 of [Wei94], thereby proving Claim 1.

Choosing a homological splitting of the complex $B(\mathcal{P})$,

$$\begin{tikzcd} H_\bullet(B(\mathcal{P})) \arrow[r, shift left, "i"] & B(\mathcal{P}) \arrow[l, shift left, "p"] \arrow[loop right, "h"] \end{tikzcd},$$

one can use dual transfer formulae of [Gra07] for homotopy coproperads to induce on the $\mathbb{S}$-bimodule $H_\bullet(B(\mathcal{P})) \simeq \mathcal{P}_2^{\text{!`}}$ the associated strongly homotopy coproperad structure, that is a differential, δ, in the free properad [2] $\mathcal{F}\big(s^{-1}H_\bullet(\bar{B}(\mathcal{P}))\big) = \Omega\big(H_\bullet(B(\mathcal{P}))\big)$ generated by $H_\bullet(B(\mathcal{P}))$. In general, this differential is *not* quadratic, i.e. the induced homotopy coproperad structure on $H_\bullet(B(\mathcal{P}))$ is *not* equal to the coproperad structure on $\mathcal{P}_2^{\text{!`}}$. Moreover, the chosen homological splitting provides us canonically with a morphism of homotopy coproperads which extends i,

$$H_\bullet(B(\mathcal{P})) \longrightarrow B(\mathcal{P}),$$

i.e. with a morphism of dg properads,

$$\phi : (\mathcal{F}(s^{-1}H_\bullet(\bar{B}(\mathcal{P}))), \delta) \longrightarrow \Omega(B(\mathcal{P})).$$

As $\Omega(B(\mathcal{P})) \xrightarrow{\simeq} \mathcal{P}$ is a resolution of $\mathcal{P}$ by Theorem 19, the required Theorem 40 follows immediately from the following

Claim 2. *Under the assumption on the properad $\mathcal{P}$ the morphism ϕ is a quasi-isomorphism.*

Indeed, the introduced above filtration of the bar construction, $B(\mathcal{P})$, induces a filtration, $F_{-p}H_\bullet(B(\mathcal{P}))$, of its homology with the associated graded coproperad being exactly $\mathcal{P}_2^{\text{!`}}$. This filtration of $H_\bullet(B(\mathcal{P}))$ induces in turn a filtration of the complex $(\mathcal{F}(s^{-1}H_\bullet(\bar{B}(\mathcal{P}))), \delta)$. The 0th term of the associated spectral sequence is precisely the minimal model, $(\mathcal{F}(s^{-1}\bar{\mathcal{P}}_2^{\text{!`}}), \delta)$, of the properad $\mathcal{P}_2$. As the latter is Koszul by assumption, its homology is equal to $\mathcal{P}_2$. By homological degree assumption on $\mathcal{P}$, the induced differential on the next term of the spectral sequence vanishes so that it degenerates. The extra grading assumption on $\mathcal{P}$ implies that this spectral sequences converges to the homology $(\mathcal{F}(s^{-1}H_\bullet(\bar{B}(\mathcal{P}))), \delta)$ which is equal, therefore, as an $\mathbb{S}$-bimodule to $\mathcal{P}_2 \simeq \mathcal{P}$. This fact completes the proof of Claim 2 and hence of the Theorem. $\square$

The operad $\mathcal{P}_2$ is Koszul means that the differential of the minimal model $(\Omega(\mathcal{P}_2^{\text{!`}}), \delta_2)$ is quadratic, that is $\delta_2 : s^{-1}\bar{\mathcal{P}}_2^{\text{!`}} \to \mathcal{F}(s^{-1}\bar{\mathcal{P}}_2^{\text{!`}})^{(2)}$. Since the transfer of homotopy coproperad structures does not change the map Δ_2 defining the homotopy coproperad structure on $H_\bullet(B(\mathcal{P}))$ but just add extra terms Δ_n, for $n \geq 3$, the final differential δ defining a minimal model of $\mathcal{P}$ is equal to δ_2 plus extra terms δ_n for $n \geq 3$ such that $\delta_n : s^{-1}\bar{\mathcal{P}}_2^{\text{!`}} \to \mathcal{F}(s^{-1}\bar{\mathcal{P}}_2^{\text{!`}})^{(n)}$, that is to say, δ is a perturbation of δ_2.

The coproperad $\mathcal{P}_2^{\text{!`}}$ is computable by Koszul duality theory. Therefore the above Theorem gives an immediate estimate of the set of generators for a minimal model of a homotopy Koszul properad. Moreover, the differential in this quasi-free model can in principle be computed via ordinary homotopy transfer formulae.

---

[2] In fact the Granåker formulae provide us in general with a differential δ in a *completed* (with respect to the number of vertices) free properad: there is no guarantee that such δ applied to a generator is a *finite* sum of terms but we can only be sure that δ is continuous with respect to the topology induced by the number of vertices filtration. However, our assumption on existence of an extra gradation in $\mathcal{P}$ implies that δ is well-defined in the ordinary category of properads: it is *finite* on every generator so that $(\mathcal{F}(s^{-1}H_\bullet(\bar{B}(\mathcal{P}))), \delta)$ makes sense without completion.

It is important to notice that had we chosen to work with topological properads (with topology induced by the number of vertices or genus filtrations), the condition (iii) in the definition of homotopy Koszulness can be safely omitted — Theorem 40 stays true in the category of (completed) topological properads because all the spectral sequences we used in the proof stay convergent by classical Complete Convergence Theorem 5.5.10 (see p.139 in [Wei94]). As an example of the deformation quantization prop [Mer04] shows, working with topological prop(erad)s is unavoidable in application of the theory of prop(erad)s to geometry and mathematical physics.

The class of properads which are homotopy Koszul but not Koszul is non-empty and contains an important example of the properad, $\mathcal{A}ss\mathcal{B}i$, of (co)associative bialgebras which can be defined as a quotient,

$$\mathcal{A}ss\mathcal{B}i := \mathcal{F}(V)/(\mathcal{R})$$

of the free properad, $\mathcal{F}(V)$, generated by the $\mathbb{S}$-bimodule $V = \{V(m,n)\}$,

$V(m,n) :=$

$\mathbb{K}[\mathbb{S}_2] \otimes \mathbb{K}[\mathbb{S}_1] \equiv \mathrm{span}$  if $m = 2, n = 1$,

$\mathbb{K}[\mathbb{S}_1] \otimes \mathbb{K}[\mathbb{S}_2] \equiv \mathrm{span}$  if $m = 1, n = 2$,

0 otherwise,

representing a binary product and a binary coproduct without symmetries, modulo the ideal generated by relations

$\mathcal{R}$: 

These relations stand respectively for the associativity of the product, the coassociativity of the coproduct and the relation between them, that is the coproduct is a morphism of algebras or equivalently the product is a morphism of coalgebras. As the ideal contains 4-vertex graphs, the properad $\mathcal{A}ss\mathcal{B}i$ is not quadratic. Hence $\mathcal{A}ss\mathcal{B}i$ can not be Koszul in the ordinary sense. However, we have the following

Proposition 41. *The properad $\mathcal{A}ss\mathcal{B}i$ is homotopy Koszul.*

Proof. (i) The properad $\mathcal{A}ss\mathcal{B}i_2$ is Koszul as it is generated by the bimodule V with the relations,



which verify the Distributive Law (see Section 5.6 and Proposition 8.5 of [Val07a]).
(ii) The $\mathbb{S}$-bimodule isomorphism $\mathcal{A}ss\mathcal{B}i \simeq \mathcal{A}ss\mathcal{B}i_2$ was established in [EE05].
(iii) The ideal generated by $\mathcal{R}$ preserves the path grading (see [MV03] for its definition and main properties) of the free properad $\mathcal{F}(V)$ and hence induces an associated filtration on $\mathcal{A}ss\mathcal{B}i$ which satisfies the last condition in the definition of a homotopy Koszulness properad. ☐

Corollary 42. *(cf.* [Mar06]*) The properad $\mathcal{A}ss\mathcal{B}i$ admits a minimal resolution, $\mathcal{F}(\mathcal{C})$, generated by the $\mathbb{S}$-bimodule $\mathcal{C} = \{\mathcal{C}(m,n)\}_{m,n\geq 1, m+n\geq 3}$, with*

$\mathcal{C}(m,n) := s^{m+n-3}\mathbb{K}[\mathbb{S}_m] \otimes \mathbb{K}[\mathbb{S}_n] =$ *span*  .

Proof. The Koszul dual properad of $\mathcal{A}ss\mathcal{B}i_2$ is the properad generated by a binary product and a binary coproduct which are associative and coassociative. All the composites with the product and the coproduct vanish except . The only non-vanishing element of this properad are obtained by composing first some products and then coproducts. We conclude that $\mathcal{A}ss\mathcal{B}i_2^{¡}(m,n) = s^{m-2}\mathbb{K}[\mathbb{S}_m] \otimes s^{n-2}\mathbb{K}[\mathbb{S}_n]$ for $m,n \geq 1, m+n \geq 3$ and zero otherwise. Then Theorem 40 implies the claim. ☐

We refer the reader to Section 6.2 for another application of the notion of homotopy Koszulness.

5.5. **Models for associative algebras, non-symmetric operads, operads, properads, props.** There are several different notions of algebraic objects in the literature that are used to model the operations acting on some algebraic category. We briefly recall them in the following table.

Operations		 no symmetry			
Composition		 planar	 non-planar		
Monoidal category	(Vect, $\otimes$)	(gVect, $\circ$)	($\mathbb{S}$-Mod, $\circ$)	($\mathbb{S}$-biMod, $\boxtimes_c$)	($\mathbb{S}$-biMod, $\boxtimes$)
Monoid	Associative algebras	Non-symmetric operads	Operads	Properads	Props
Modules	Modules	Non-symmetric algebras	Algebras	(Bial)gebras	(Bial)gebras
Free monoid	Ladders (Tensor module)	Planar trees	Trees	Connected graphs	Graphs

To each pair of such objects, there is a forgetful functor and a left adjoint :

$$\text{Associative algebras} \rightleftarrows \text{Non-symmetric operads} \rightleftarrows \text{Operads} \rightleftarrows \text{Properads} \rightleftarrows \text{Props}.$$

Let us make them explicit.

- To any prop $\mathcal{P}$, the associated properad $\mathcal{U}^{\text{props}}_{\text{properads}}(\mathcal{P})$ is given by the same underlying $\mathbb{S}$-bimodule where we only consider vertical compositions of operations based on connected graphs. That is we forget the horizontal composition. Its left adjoint $\mathcal{F}^{\text{props}}_{\text{properads}}(\mathcal{P})$ is given by the free symmetric tensor on $\mathcal{P}$ for the horizontal tensor product. (This functor was introduced in [Val07a] at Section 1 where it is denoted by $\mathcal{S}$.) In other words, we freely generate horizontal compositions from a properad to get a prop.

- The operad obtained from a properad $\mathcal{P}$ is the $\mathbb{S}$-module $\mathcal{U}^{\text{properads}}_{\text{operads}}(\mathcal{P})(n) := \mathcal{P}(1, n)$ equipped with the restriction to one rooted trees composition. Its left adjoint functor is $\mathcal{F}^{\text{properads}}_{\text{operads}}(\mathcal{P})(m, n) := \mathcal{P}(n)$ for $m = 1$ and 0 for $m > 1$.
- For any operad $\mathcal{P}$, we consider the non-symmetric operad $\mathcal{U}^{\text{operads}}_{\text{non-symm. operads}}(\mathcal{P}) = \mathcal{P}$ where we forget the action of the symmetric group. The left adjoint is given by
$$\mathcal{F}^{\text{operads}}_{\text{non-symm. operads}}(\mathcal{P})(n) = \mathcal{P}(n) \otimes \mathbb{K}[\mathbb{S}_n].$$
(see M. Aguiar and M. Livernet [AL07])
- The pair of adjoint functors between associative algebras and non-symmetric operads is defined in the same then the pair of functors between operads and properads. In one way, we just consider the unital operation (arity (1)) of a non-symmetric operad. In this other way, for an associative algebra we define a non-symmetric operad concentrated in arity (1).

Proposition 43. *All these functors are exact, that is the image of a quasi-isomorphism is a quasi-isomorphism.*

Proof. It is trivial for the forgetful functors and for the functors $F^{\text{operads}}_{\text{operads}}$ and $F^{\text{non-symm. operads}}_{\text{ass. algebras}}$ because the underlying dg-module does not change. Since the functor $F^{\text{operads}}_{\text{non-symm. operads}}$ is given by tensoring $\mathbb{S}_n$-modules with the flat $\mathbb{K}$ module $\mathbb{K}[\mathbb{S}_n]$ (the characteristic of $\mathbb{K}$ is 0), it is exact. Over a field of characteristic 0, the functor $F^{\text{props}}_{\text{properads}}$ is also exact . ☐

This proposition justifies the following philosophy. To study the deformation theory of elements of an algebraic category, that is a class of gebras (modules, algebras, bialgebras), one should first model this category using the simplest possible object of the previous table. For instance, associative, diassociative, dendriform algebras [LFCG01] are encoded each time by a non-symmetric

operad. Commutative, Lie, PreLie, Gerstenhaber, Poisson algebras are modelled by operads. Lie bialgebras, infinitesimal Hopf algebras [Agu00], (associative) bialgebras (see 9.3) are representations of properads. Non-unital infinitesimal Hopf algebras, Semi Hopf algebras, Lily bialgebras [Lod06] can only be represented by a prop.
Then to study the deformation theory of this algebraic category, that is to define the stable notion up to homotopy (see 6.1) or the deformation complex (see 8.2), one has to find a cofibrant resolution (bar-cobar, minimal model for instance) of the related operad, properad or prop $\mathcal{P}$. This resolution contains all the necessary data since a resolution for the induced prop is "freely" obtained by the free exact functor.

5.6. **Models generated by genus 0 differentials.** Let $\mathcal{A}$ be a category of gebras defined by some products and some coproducts with relations that can be written as linear combinations of connected graphs of genus 0, for example Lie bialgebras, Frobenius bialgebras, Infinitesimal bialgebras (see [Gan03, Val07a]). In this case, the class of gebras can be faithfully modelled with a smaller algebraic object called a dioperad [Gan03].

A dioperad is a properad with only compositions of operations based on genus 0 connected graphs. Hence, there is a natural forgetful functor from properads to dioperads. To any properad $\mathcal{P}$, the associated dioperad $\mathcal{U}^{\text{properads}}_{\text{dioperads}}(\mathcal{P})$ has the same underlying $\mathbb{S}$-bimodule and we only consider vertical compositions of operations based on connected graphs of genus 0. Let us denote by $\square$ the restriction of $\boxtimes$ to genus 0 graphs. With this notation, a dioperad is a monoid $(\mathcal{D}, \mu_{\mathcal{D}})$ in the monoidal category $(\mathbb{S}\text{-biMod}, \square)$. From now on, let us denote the genus in exponent. For instance, $\mathcal{F}^0$ will denote the free dioperad functor $\mathcal{F}^{\text{dioperads}}_{\mathbb{S}\text{-biMod}}$ and $\mathcal{F}$ will simply denote the free properad functor $\mathcal{F}^{\text{properads}}_{\mathbb{S}\text{-biMod}}$.

Proposition 44. *The left adjoint of the forgetful functor* $\mathcal{U}^{\textit{properads}}_{\textit{dioperads}}(\mathcal{P})\ :\ \text{Properads} \to \text{Dioperads}$ *is given by*

$$\mathcal{F}(\mathcal{D})/I,$$

where I is the (properadic) ideal generated by the image under $\mu_{\mathcal{D}} - \mathrm{Id}$ of $\mathcal{F}^0(\mathcal{D})^{(2)}$, that is the connected graphs of genus 0 with two vertices.
In other words, this construction is the quotient of the free properad on $\mathcal{D}$, consider as an $\mathbb{S}$-bimodule, by the (dioperadic) composition of any pair of adjacent vertices with only one edge in between.

Notice that this construction is the same as the universal enveloping algebra of a Lie algebra. Therefore, we will often call it *the universal enveloping properad of a dioperad* and $\mathcal{F}^{\text{properads}}_{\textit{dioperads}}$ the *universal enveloping functor.*

Proof. The proof is the same as the proof of the universal property of the universal enveloping algebra of a Lie algebra. Hence it is left to the reader. $\square$

A direct corollary gives that the universal enveloping properad of a dioperad defined by generators and relations is a properad given the same generators and relations.

Corollary 45. *Let $\mathcal{D}$ be a dioperad defined by generators and relations : $\mathcal{D} = \mathcal{F}^0(V)/(R)$, where (R) is the (dioperadic) ideal generated by R. The universal enveloping properad is equal to*

$$\mathcal{F}^{\textit{properads}}_{\textit{dioperad}}(\mathcal{D}) = \mathcal{F}(V)/(R),$$

where (R) is the (properadic) ideal generated by R.

Even if an algebraic category $\mathcal{A}$ can be modelled by a dioperad, the induced cofibrant resolution of this dioperad does not contains all the data necessary for the study of deformation theory of $\mathcal{A}$ because the universal enveloping functor $\mathcal{F}^{\text{properads}}_{\text{dioperads}}$ is not exact as the following counter-example shows.

Let $\varepsilon\mathcal{B}i$ be the properad which models infinitesimal bialgebras (see [Val07a] Section 2.9). We consider its Koszul dual properad without the relation  $= 0$. Let us denote it by $\mathcal{NC}$-$\mathcal{F}rob$

because it models some kind of non-commutative Frobenius bialgebras. A $\mathcal{NC}$-$\mathcal{F}rob$-bialgebra is a vector space X equipped with a binary associative product $\mu : X \otimes X \to X$ and a binary coassociative coproduct $\Delta : X \to X \otimes X$ such that Δ is a morphism of bimodules. This means

$$\Delta \circ \mu = (\mathrm{Id} \otimes \mu) \circ (\Delta \otimes \mathrm{Id}) = (\mu \otimes \mathrm{Id}) \circ (\mathrm{Id} \otimes \Delta).$$

The graphical picture of all the relations is the following :



Since the relations are linear combinations of connected graphs of genus 0, this category is faithfully modelled by the dioperad $\mathcal{NC}$-$\mathcal{F}rob^0 = \mathcal{F}^0(V)/(R)$. The exponent 0 stands for the restriction to graphs of genus 0. It was proved in [Gan03] that $\mathcal{NC}$-$\mathcal{F}rob^0$ is a Koszul dioperad, since its Koszul dual dioperad $\varepsilon\mathcal{B}i^0$ is Koszul by means of distributive laws. That is the dioperad $\mathcal{NC}$-$\mathcal{F}rob^0$ admits a quadratic dioperadic (genus 0) model $(\mathcal{F}^0(\mathcal{C}), \partial^0) \xrightarrow{\sim} \mathcal{NC}$-$\mathcal{F}rob^0$, where $\mathcal{C}$ is the codioperad $s^{-1}\varepsilon\mathcal{B}i^{0\vee}$. (Notice that there is no direct proof of this fact.) The differential ∂^0 splits each element of $\mathcal{C}$ into two vertices with only one edge in between.

Consider now the properad $\mathcal{NC}$-$\mathcal{F}rob = \mathcal{F}(V)/(R)$, which is the image under the universal enveloping functor $\mathcal{F}_{\text{dioperads}}^{\text{properads}}$ of $\mathcal{F}rob^0$ by Corollary 45. The image of the chain complex $(\mathcal{F}^0(\mathcal{C}), \partial^0)$ under the functor $\mathcal{F}_{\text{dioperads}}^{\text{properads}}$ is the quasi-free properad on $\mathcal{C}$ with the differential ∂^0, that is the cobar construction of $\mathcal{C}$, where this later is considered as a coproperad. The homology of this chain complex is not concentrated in degree 0.

We build a cycle based on graphs of genus 2 from the following picture :

$$\begin{tikzcd}
\mu \circ \Delta \circ \mu \circ \Delta \arrow[r, "-\mu\circ R_{lm}\circ\Delta"] \arrow[d, "\mu\circ R_{rm}\circ\Delta"] & \mu \circ (\mu \otimes \mathrm{Id}) \circ (\mathrm{Id} \otimes \Delta) \circ \Delta \arrow[d, "\mu\circ(\mu\otimes\mathrm{Id})\circ R_c"] \\
\mu \circ (\mathrm{Id} \otimes \mu) \circ (\Delta \otimes \mathrm{Id}) \circ \Delta \arrow[r, "-R_a\circ(\Delta\otimes\mathrm{Id})\circ\Delta"] & \mu \circ (\mu \otimes \mathrm{Id}) \circ (\Delta \otimes \mathrm{Id}) \circ \Delta,
\end{tikzcd}$$

where R_{rm} stands for the "right module" relation $\mu \circ \Delta \to (\mathrm{Id} \otimes \mu) \circ (\Delta \otimes \mathrm{Id})$, R_{lm} for the "left module" relation $\mu \circ \Delta \to (\mu \otimes \mathrm{Id}) \circ (\mathrm{Id} \otimes \Delta)$, R_a the associativity relation $\mu \circ (\mu \otimes \mathrm{Id}) \to \mu \circ (\mathrm{Id} \otimes \mu)$ and R_c the coassociativity relation $(\Delta \otimes \mathrm{Id}) \circ \Delta \to (\mathrm{Id} \otimes \Delta) \circ \Delta$.
The graphical picture is as follows :



Then, the cycle is based upon the following picture :



We denote with the same notation the corresponding homotopies, that is elements of $\mathcal{C}$:

$$\partial_0(R_{rm}) = \mu \circ \Delta - (\mathrm{Id} \otimes \mu) \circ (\Delta \otimes \mathrm{Id}) \quad , \quad \partial_0(R_{lm}) = \mu \circ \Delta - (\mu \otimes \mathrm{Id}) \circ (\mathrm{Id} \otimes \Delta),$$
$$\partial_0(R_a) = \mu \circ (\mu \otimes \mathrm{Id}) - \mu \circ (\mathrm{Id} \otimes \mu) \quad , \quad \partial_0(R_c) = (\Delta \otimes \mathrm{Id}) \circ \Delta - (\mathrm{Id} \otimes \Delta) \circ \Delta.$$

The previous picture proves that

$$\xi := \mu \circ R_{rm} \circ \Delta - \mu \circ R_{lm} \circ \Delta - R_a \circ (\Delta \otimes \mathrm{Id}) \circ \Delta + \mu \circ (\mu \otimes \mathrm{Id}) \circ R_c$$

is a cycle in $(\mathcal{F}(\mathcal{C}), \partial^0)$, that is $\partial^0(\xi) = 0$.

Lemma 46. *The cycle ξ is not a boundary under ∂^0.*

Proof. The degree of ξ is 1. Suppose that there exists an element ζ of degree 2 such that $\partial^0(\zeta) = \xi$. This element belongs to

$$\zeta \in \mathcal{F}(\mathcal{C}_0 \oplus \underbrace{\mathcal{C}_1}_{(2)}) \oplus \mathcal{F}(\mathcal{C}_0 \oplus \underbrace{\mathcal{C}_2}_{(1)}).$$

Let us denote by $\zeta = \zeta_1 + \zeta_2$ each component. The image under the quadratic differential ∂^0 of any element of $\mathcal{F}(\underbrace{\mathcal{C}_0}_{(k)} \oplus \underbrace{\mathcal{C}_2}_{(1)})$ is an element of $\mathcal{F}(\underbrace{\mathcal{C}_0}_{(k+1)} \oplus \underbrace{\mathcal{C}_1}_{(1)})$. And since the genus of the differential ∂^0 is 0, ζ_2 is in $\mathcal{F}^2(\underbrace{\mathcal{C}_0}_{(1)} \oplus \underbrace{\mathcal{C}_2}_{(1)})$, that is the part of genus 2 of $\mathcal{F}(\mathcal{C}_0 \oplus \mathcal{C}_2)$. The $\mathbb{S}$-bimodule $\mathcal{C}_0$ is equal to $V =$ $\oplus$, that is binary. Hence $\mathcal{F}(\underbrace{\mathcal{C}_0}_{(1)} \oplus \underbrace{\mathcal{C}_2}_{(1)})$ is concentrated in genus 0 and 1, which proves $\zeta_2 = 0$.

Since the image of $\mathcal{F}(\underbrace{\mathcal{C}_0}_{(k)} \oplus \underbrace{\mathcal{C}_1}_{(2)})$ under ∂^0 is in $\mathcal{F}(\underbrace{\mathcal{C}_0}_{(k+2)} \oplus \underbrace{\mathcal{C}_1}_{(1)})$, ζ_1 must belong to $\mathcal{F}(\mathcal{C}_1)^{(2)}$. More precisely, ζ_1 is a element of $\mathcal{F}^2(\mathcal{C}_1)^{(2)}$ because the differential ∂^0 preserves the genus. The $\mathbb{S}$-bimodule $\mathcal{C}_1$ is generated by the four elements $R_{rm} \in \mathcal{C}(2,2)$, $R_{lm} \in \mathcal{C}(2,2)$, $R_a \in \mathcal{C}(1,3)$ and $R_c \in \mathcal{C}(3,1)$. The only way to get an element of genus 2 is to graft one element from $\mathcal{C}(1,3)$ to an element from $\mathcal{C}(3,1)$. Finally ζ is linear combination of $R_c \circ \sigma \circ Ra$, with $\sigma \in \mathbb{S}_3$. And in this case, $\partial^0(\zeta)$ cannot contain elements like $\mu \circ R_{rm} \circ \Delta - \mu \circ R_{lm} \circ \Delta$ whence the contradiction. $\square$

This counter-example answers a question raised by [MV03], that is the functor $\mathcal{F}^{\mathrm{props}}_{\mathrm{dioperads}}$ is not exact.

Theorem 47. *The universal enveloping functor $\mathcal{F}^{properads}_{dioperads}$ is not exact.*

For this reason, we are reluctant to include dioperads in the preceding table. It is not enough in general to find a resolution of the genus 0 part of a properad to generate a complete resolution of it. Nevertheless, it is sometimes the case. We have emphasized the class of properads that admits a quadratic model, that is Koszul properad. We do the same thing with properads for which there exists a model with a genus 0 differential.

Definition (Contractible properad)**.** We call *contractible properad* any properad $\mathcal{P}$ that admits a model $(\mathcal{F}(\mathcal{C}), \partial^0) \xrightarrow{\sim} \mathcal{P}$ with $\partial^0|_{\mathcal{C}} : \mathcal{C} \to \mathcal{F}^0(\mathcal{C})$, that is the part of genus 0 of the free properad on $\mathcal{C}$.

It is equivalent to ask that $\mathcal{C}$ is a homotopy coproperad with structure maps $\delta_n : \mathcal{C} \to \mathcal{F}^0(\mathcal{C})^{(n)}$ with image of genus 0. In other words, $\mathcal{C}$ is a *homotopy codioperad*.

Proposition 48. *Let $\mathcal{P} = \mathcal{F}(V)/(R)$ be a properad defined by genus* 0 *relations, $R \subset \mathcal{F}^0(V)$. The properad $\mathcal{P}$ is a contractible properad if and only if the associated dioperad $\mathcal{D} := \mathcal{F}^0(V)/(R)$ admits a quasi-free (dioperadic) resolution $(\mathcal{F}^0(\mathcal{C}), \partial^0) \xrightarrow{\sim} \mathcal{D}$, which is a quasi-isomorphism preserved by the universal enveloping functor $\mathcal{F}^{properads}_{dioperads}$.*

Proof. If $\mathcal{P}$ is contractible, we denote by $(\mathcal{F}(\mathcal{C}), \partial^0) \xrightarrow{\sim} \mathcal{P}$ its genus 0 differential model. Since ∂^0 preserves the genus, the chain complex $(\mathcal{F}(\mathcal{C}), \partial^0)$ is equal to the direct sum of sub-complexes $\bigoplus_{g\geq 0}(\mathcal{F}^g(\mathcal{C}), \partial^0)$. Hence, the genus 0 chain complex is a a resolution of $\mathcal{D}$. And by Corollary 45 the image under the universal enveloping functor $\mathcal{F}^{\text{properads}}_{\text{dioperads}}$ of the quasi-isomorphism $(\mathcal{F}^0(\mathcal{C}), \partial^0) \xrightarrow{\sim} \mathcal{D}$ is the resolution $(\mathcal{F}(\mathcal{C}), \partial^0) \xrightarrow{\sim} \mathcal{P}$. The other way is trivial. □

A Koszul contractible properad $\mathcal{P}$ is a properad with a minimal model $(\mathcal{F}(\mathcal{C}), \partial^0) \xrightarrow{\sim} \mathcal{P}$ whose differential ∂^0 is quadratic and genus 0. It is equivalent to say that $\mathcal{C}$ is a codioperad. If a properad $\mathcal{P} = \mathcal{F}(V)/(R)$ with genus 0 relations is contractible Koszul, then the associated dioperad $\mathcal{D} = \mathcal{F}^0(V)/(R)$ is Koszul in the sense of [Gan03]. But it is not true that any Koszul dioperad is a Koszul contractible properad as the example of $\mathcal{NC}$-$\mathcal{F}rob$ shows. Lemma 46 shows that it is not contractible. Moreover we shall see below that it is not Koszul as a properad either.

Proposition 49. *Let $\mathcal{P} = \mathcal{F}(V)/(R)$ be a Koszul properad defined by a finite dimensional $\mathbb{S}$-bimodule V and by genus* 0 *relations, $R \subset \mathcal{F}^0(V)$. If the Koszul dual properad of $\mathcal{P}$ is equal, as an $\mathbb{S}$-bimodule, to the Koszul dual dioperad of the associated dioperad $\mathcal{D} := \mathcal{F}^0(V)/(R)$ then the properad $\mathcal{P}$ is contractible.*

Proof. In this case, the Koszul dual coproperad $\mathcal{P}^{\text{¡}} = \mathcal{P}^{!\vee}$ is equal to the Koszul dual dioperad $\mathcal{D}_{\text{¡}} = \mathcal{D}^{!\vee}$. Hence the image of the partial coproduct $\Delta_{(1,1)} : \mathcal{P}^{\text{¡}} \to \mathcal{P}^{\text{¡}} \boxtimes \mathcal{P}^{\text{¡}}$ is actually in $\mathcal{P}^{\text{¡}} \square \mathcal{P}^{\text{¡}}$ which is the part of genus 0 of $\mathcal{P}^{\text{¡}} \boxtimes \mathcal{P}^{\text{¡}}$. □

The Koszul dual properad is equal to the Koszul dual dioperad if and only if the part of genus > 0 of $\mathcal{P}^!$ vanished, that is $\mathcal{F}^g(V^\vee)/(R^\perp) = 0$ for $g > 0$. Proposition 49 allows us to give examples of Koszul contractible properads. One way to prove that a properad is Koszul is by means of *distributive laws* (see Proposition 8.4 of [Val07a]). Let $\mathcal{P}$ be a quadratic properad of the form $\mathcal{P} = \mathcal{F}(V, W)/(R \oplus D \oplus S)$, where $R \subset \mathcal{F}^{(2)}(V)$, $S \subset \mathcal{F}^{(2)}(W)$ and where

$$D \subset (I \oplus \underbrace{W}_{1}) \boxtimes_c (I \oplus \underbrace{V}_{1}) \bigoplus (I \oplus \underbrace{V}_{1}) \boxtimes_c (I \oplus \underbrace{W}_{1}).$$

The two pairs of $\mathbb{S}$-bimodules (V, R) and (W, S) generate two properads denoted $\mathcal{A} := \mathcal{F}(V)/(R)$ and $\mathcal{B} := \mathcal{F}(W)/(S)$.

Definition (Distributive law)**.** Let λ be a morphism of $\mathbb{S}$-bimodules

$$\lambda : (I \oplus \underbrace{W}_{1}) \boxtimes_c (I \oplus \underbrace{V}_{1}) \to (I \oplus \underbrace{V}_{1}) \boxtimes_c (I \oplus \underbrace{W}_{1}).$$

such that the $\mathbb{S}$-bimodule D is defined by the image of

$$(id, -\lambda) : (I \oplus \underbrace{W}_{1}) \boxtimes_c (I \oplus \underbrace{V}_{1}) \to (I \oplus \underbrace{W}_{1}) \boxtimes_c (I \oplus \underbrace{V}_{1}) \bigoplus (I \oplus \underbrace{V}_{1}) \boxtimes_c (I \oplus \underbrace{W}_{1}).$$

We call λ a *distributive law* and denote D by D_λ if the two following morphisms are injective

$$\begin{cases} \underbrace{\mathcal{A}}_{1} \boxtimes_c \underbrace{\mathcal{B}}_{2} \to \mathcal{P} \\ \underbrace{\mathcal{A}}_{2} \boxtimes_c \underbrace{\mathcal{B}}_{1} \to \mathcal{P}. \end{cases}$$

The last condition must be seen as a coherence axiom, which ensures that the natural morphism $\mathcal{A} \boxtimes \mathcal{B} \to \mathcal{P}$ is injective. In this case, Proposition 8.4 of [Val07b] states that $\mathcal{P}$ is Koszul if $\mathcal{A}$ and $\mathcal{B}$ are Koszul. A properad is called *binary* if it is generated by binary products and coproducts.

Proposition 50. *Let $\mathcal{D} = \mathcal{F}^0(V)/(R)$ be a binary Koszul dioperad defined by a distributive law such that V is finite dimensional. Then the associated properad $\mathcal{P} := \mathcal{F}(V)/(R)$ is Koszul and contractible.*

Proof. If a binary dioperad $\mathcal{D}$ defined by a distributive law verifies the hypotheses of Proposition 5.9 of [Gan03], then the associated properad $\mathcal{P}$ is also defined by distributive law and verifies the hypotheses od Proposition 8.4 of [Val07a]. In this case, the Koszul dual coproperad, given by Proposition 8.2 of [Val07a] has a genus 0 coproduct. $\square$

Corollary 51. *The properads $\mathcal{B}i\mathcal{L}ie$ of Lie bialgebras and $\varepsilon\mathcal{B}i$ of infinitesimal Hopf algebras are Koszul contractible.*

In this case, the Koszul dual (co)dioperad provides the good space of "homotopies" for the resolution of the properad. Therefore, it gives the proper notion of homotopy $\mathcal{P}$-gebra (see 6.1). An example of this fact for $\mathcal{B}i\mathcal{L}ie$ can be found in Section 9.2in [Gan05, Mer06].

Remark. Dually, in this case, the products of operations based on strictly positive genus graphs of the Koszul dual properad always vanish. If g denotes the genus of the underlying graph, it means that any such product is equivalent to products based on graphs with g simple loops , using the relations of the products and the relations of coproducts. Therefore, it is zero because of the relation  $= 0$ in the Koszul dual properad. This statement is a non-trivial result about the coherence of the relations of a properad.

To any binary properad $\mathcal{P}$, we associate a properad $\mathcal{P}_\diamond$ which codes $\mathcal{P}$-gebras satisfying the extra loop relation  $= 0$. Since the properad $\mathcal{B}i\mathcal{L}ie$ is Koszul, its Koszul dual properad $\mathcal{F}rob_\diamond$ is also Koszul by Koszul duality theory. This means that $\mathcal{F}rob_\diamond$ has a quadratic model. Since the properad $\mathcal{B}i\mathcal{L}ie$ has non trivial higher genus compositions, this model is not contractible, that is the boundary map creates higher genus graphs. The example $\mathcal{F}rob_\diamond$ provides an example of a Koszul non-contractible properad. (We do not know how to prove this result without the help of Koszul duality for properads).

Let $\mathcal{C}$ denote the Koszul dual coproperad of $\mathcal{NC}$-$\mathcal{F}rob$, that is $\mathcal{C} = s^{-1}\varepsilon\mathcal{B}i_\diamond{}^\vee$. Recall that a properad $\mathcal{P}$ is Koszul if and only if the cobar construction of the Koszul dual coproperad $\Omega(\mathcal{P}^{\text{i}}) = (\mathcal{F}(\mathcal{P}^{\text{i}}), \partial)$ is a resolution of $\mathcal{P}$. This statement is not true for $\mathcal{NC}$-$\mathcal{F}rob$. The cycle ξ given above induces a non-trivial element in homology.

Lemma 52. *The cycle ξ is not a boundary under ∂.*

Proof. We use the same notations as in Lemma 46 but applied to ∂ instead of ∂^0. The space $\mathcal{C}_1$ is generated by the elements R_{lm}, R_{rm}, R_a, R_c and some R_i for $i = 1, \ldots, 4$. For the same reason, ζ_1 must be an an element of $\mathcal{F}(\mathcal{C}_1)^{(2)}$. Since the image under ∂ of any element of $\mathcal{C}_1$ is a graph with two adjacent vertices indexed by  or , the element $\mu \circ R_{lm} \circ \Delta$ cannot belong to $\partial(\zeta_1)$. Hence $\mu \circ R_{lm} \circ \Delta$ must be an element of $\partial(\zeta_2)$. Since ∂ is quadratic, there exists an element S in $\mathcal{C}_2$ such that $\partial(S) = \mu \circ R_{lm} + \cdots$ or $\partial(S) = R_{lm} \circ \Delta + \cdots$. Such an S has to be an element of either $\varepsilon\mathcal{B}i_\diamond^\vee(1,2)^{(3)}$ or $\varepsilon\mathcal{B}i_\diamond^\vee(2,1)^{(3)}$. Consider the first case, the second one being symmetrical. The only element in $\varepsilon\mathcal{B}i_{\diamond}^{\vee}(1,2)^{(3)}$ whose partial coproduct includes $\mu \circ R_{lm}$ is the dual of the composite of



in $\varepsilon\mathcal{B}i_{\diamond}(1,2)^{(3)}$. The associativity relation and the loop relation in $\varepsilon\mathcal{B}i_{\diamond}$ show that this composite is equal to zero, which concludes the proof. $\square$

Theorem 53. *The properad $\mathcal{NC}$-$\mathcal{F}rob$ of non-commutative Frobenius bialgebras and the properad $\varepsilon\mathcal{B}i_{\diamond}$ of involutive infinitesimal bialgebras are not Koszul.*

We hope that this helps to clarify the general picture of models for prop(erad)s.

6. Homotopy $\mathcal{P}$-gebra

In this section, we define the notion of $\mathcal{P}$-gebra up to homotopy or *homotopy $\mathcal{P}$-gebra*. We make explicit structures of homotopy $\mathcal{P}$-gebras in terms of Maurer-Cartan elements. We also define and make explicit morphisms of homotopy $\mathcal{P}$-algebras, when $\mathcal{P}$ is an operad, in terms of Maurer-Cartan elements in an L_∞-algebra. This last part uses the notion of homotopy Koszul (colored) operads defined in the previous section.

6.1. $\mathcal{P}$-gebra, $\mathcal{P}_{(n)}$-gebra and homotopy $\mathcal{P}$-gebra.

Let $\mathcal{P}$ be a dg prop(erad) and $\Omega(\mathcal{C})$ be a model of $\mathcal{P}$.

Definition (Homotopy $\mathcal{P}$-gebra)**.** A structure of *homotopy $\mathcal{P}$-gebra* on a dg module X is a morphism of dg prop(erad)s : $\Omega(\mathcal{C}) \to \mathrm{End}_X$.

Any $\mathcal{P}$-gebra is a homotopy $\mathcal{P}$-gebra of particular type. In this case, the morphism of dg-properads factors through $\mathcal{P}$, that is $\Omega(\mathcal{C}) \xrightarrow{\sim} \mathcal{P} \to \mathrm{End}_X$. For the Koszul operads $\mathcal{A}ss$, $\mathcal{C}om$, $\mathcal{L}ie$, this notion coincides with homotopy associative, commutative, Lie algebras. For the properads $\mathcal{B}i\mathcal{L}ie$ and $\mathcal{A}ss\mathcal{B}i$, we get the notions of homotopy Lie bialgebras and homotopy bialgebras. Since $\mathcal{B}i\mathcal{L}ie$ is contractible, the explicit definition given in [Gan05, Mer06] coincides with this one.

Theorem 62 shows that a structure of homotopy $\mathcal{P}$-gebra on X is equivalent to a morphism of $\mathbb{S}$-bimodules in $s^{-1}\mathrm{Hom}_0^{\mathbb{S}}(\mathcal{C}, \mathrm{End}_X)$ which is a Maurer-Cartan element in the L_∞-convolution algebra $\mathrm{Hom}^{\mathbb{S}}(\mathcal{C}, \mathrm{End}_X)$.

Theorem 54. *A $\mathcal{P}$-gebra structure on X is equivalent to a Maurer-Cartan element in $\mathrm{Hom}^{\mathbb{S}}(\mathcal{C}, \mathrm{End}_X)$.*

This notion is well defined and independent of the choice of a model. By Theorem 70, if $\Omega(\mathcal{C}_1)$ and $\Omega(\mathcal{C}_2)$ are two models of $\mathcal{P}$, then the convolution L_∞-algebras are quasi-isomorphic, which induces a bijection between the set of Maurer-Cartan elements.

We can discuss the form of the solutions of the Maurer-Cartan equation. It gives the following definition.

Definition ($\mathcal{P}_{(n)}$-gebra)**.** A dg module X endowed with a Maurer-Cartan element γ in $\mathrm{Hom}^{\mathbb{S}}(\mathcal{C}, \mathrm{End}_X)$ such that $\gamma(c) = 0$ for every $c \in \mathcal{C}_{k>n}$ is called a *$\mathcal{P}_{(n)}$-gebra*.

This notion is the direct generalization of the notion of $A_{(n)}$-algebra of Stasheff [Sta63] or $L_{(n)}$-algebras. A $\mathcal{P}_{(n)}$-gebra is a homotopy $\mathcal{P}$-gebra with strict relations from degree n.

6.2. Morphisms of homotopy $\mathcal{P}$-algebras as Maurer-Cartan elements.

Another application of the notion of homotopy Koszul can be found in the study of morphisms between homotopy $\mathcal{P}$-algebras. A *colored properad* is an operad such that the inputs and outputs are labelled by an extra labelling and such that the composition is coherent with respect to this extra labelling. That is if the 'colors' (labelling) do not match, the composition of operations vanishes. It is proven in [VdL03] how to extend Koszul duality of operads to colored operads. It is straightforward to generalize Theorem 40 to this case.

Let $\mathcal{P} = \mathcal{F}(V)/(R)$ be a Koszul operad. One can define the 2-colored operad $\mathcal{P}_{\bullet\to\bullet}$ by $\mathcal{P} = \mathcal{F}(V_1 \oplus V_2 \oplus f)/(R_1 \oplus R_2 \oplus R_{\bullet\to\bullet})$, where V_1 and R_1 (resp. V_2 and R_2) are copies of V and R with inputs and outputs labelled by the color 1 (resp. 2), f is a generator of arity $(1,1)$ which goes from 1 to 2 and $R_{\bullet\to\bullet}$ is generated by $v \circ f^{\otimes n} - f \circ v$ for any element $v \in V(n)$ (see [Mar04a] for more details). The purpose of this definition lies in the following result. A structure of $\mathcal{P}_{\bullet\to\bullet}$-algebra is the data of two $\mathcal{P}$-algebras with a morphism of $\mathcal{P}$-algebras between them.

Lemma 55. *When $\mathcal{P}$ is Koszul generated by a finite dimensional $\mathbb{S}$-module V such that $V(1) = 0$, the 2-colored operad $\mathcal{P}_{\bullet\to\bullet}$ is homotopy Koszul.*

Proof. (i) The operad $(\mathcal{P}_{\bullet\to\bullet})_2$ is equal to $\mathcal{F}(V_1 \oplus V_2 \oplus f)/(R_1 \oplus R_2 \oplus R_\bullet)$, where $R_\bullet = f \circ V_1$. Hence, it is equal to $(\mathcal{P}_{\bullet\to\bullet})_2 \cong \mathcal{P}_1 \oplus \mathcal{P}_2 \oplus \mathcal{P} \circ (I \oplus \underbrace{f}_{\geqslant 1})$. Its Koszul dual is equal to $(\mathcal{P}_{\bullet\to\bullet})^{\text{¡}}_2 = \mathcal{P}^{\text{¡}}_1 \oplus \mathcal{P}^{\text{¡}}_2 \oplus s(f \circ \mathcal{P}^{\text{¡}})$. Therefore, $(\mathcal{F}(s^{-1}\bar{\mathcal{P}}^{\text{¡}}_1 \oplus s^{-1}\bar{\mathcal{P}}^{\text{¡}}_2 \oplus f \circ \bar{\mathcal{P}}^{\text{¡}})), \delta_2)$ is a quadratic model of $(\mathcal{P}_{\bullet\to\bullet})_2$, because δ_2 is equal to 3 copies of the Koszul resolution of $\mathcal{P}$.

(ii) Since $\mathcal{P}_{\bullet\to\bullet} \cong \mathcal{P}_1 \oplus \mathcal{P}_2 \oplus \mathcal{P} \circ (I \oplus \underbrace{f}_{\geqslant 1})$, it is equal to $(\mathcal{P}_{\bullet\to\bullet})_2$.

(iii) Since V is finite dimensional and $V(1) = 0$, the filtration with the number of leaves gives a suitable filtration. ☐

In this case, the minimal model of $\mathcal{P}_{\bullet\to\bullet}$ is given by $(\mathcal{F}(s^{-1}\bar{\mathcal{P}}^{\text{¡}}_1 \oplus s^{-1}\bar{\mathcal{P}}^{\text{¡}}_2 \oplus f \circ \bar{\mathcal{P}}^{\text{¡}})), \delta)$ by Theorem 40.

Proposition 56. *An algebra over the model of $\mathcal{P}_{\bullet\to\bullet}$ is the data of two homotopy $\mathcal{P}$-algebras with a homotopy (or weak) morphism between them.*

Proof. A morphism of 2-colored operads $(\mathcal{F}(s^{-1}\bar{\mathcal{P}}^{\text{¡}}_{\bullet\to\bullet 2}), \delta) \to \mathrm{End}_{X,Y}$ defines a homotopy $\mathcal{P}$-algebra structure on X and Y. The component on $\{\mathrm{Hom}(X^{\otimes n}, Y)\}_{n\geqslant 1}$ is equivalent to a morphism of dg $\mathcal{P}^{\text{¡}}$-coalgebras $\mathcal{P}^{\text{¡}}(X) \to \mathcal{P}^{\text{¡}}(Y)$, that is between the bar constructions of X and Y. ☐

Theorem 57. *Morphisms of homotopy $\mathcal{P}$-algebras between X and Y are in one-to-one correspondence with Maurer-Cartan elements in the L_∞-algebra* $(\mathrm{Hom}^{\mathbb{S}}(\mathcal{P}^{\text{¡}}_{\bullet\to\bullet 2}, \mathrm{End}_{X,Y}), \delta)$

Notice that this result was already proved by hands in [Dol07] in the case of homotopy Lie algebras.

Finally, a structure of homotopy $\mathcal{P}$-algebra on X is a Maurer-Cartan element in the strict Lie algebra $\mathrm{Hom}^{\mathbb{S}}(\mathcal{P}^{\text{¡}}, \mathrm{End}_X)$, whereas a morphism of homotopy $\mathcal{P}$-algebras between X and Y is a (generalized) Maurer-Cartan element in the homotopy Lie algebra $\mathrm{Hom}^{\mathbb{S}}(\mathcal{P}^{\text{¡}}, \mathrm{End}_{X,Y})$. The conceptual explanation of this phenomenon is the following. In the first case, we have a quadratic model of the Koszul operad $\mathcal{P}$ and the second case, we use a non-quadratic model of the homotopy Koszul 2-colored operad $\mathcal{P}_{\bullet\to\bullet}$.

7. L_∞-ALGEBRAS, DG MANIFOLDS, DG AFFINE SCHEMES AND MORPHISMS OF PROP(ERAD)S

7.1. **L_∞-algebras, dg manifolds and dg affine schemes.** Structure of a L_∞-algebra on a $\mathbb{Z}$-graded vector space $\mathfrak{g}$ is, by definition, a degree -1 coderivation, $Q : \odot^{\geq 1} s\mathfrak{g} \to \odot^{\geq 1} s\mathfrak{g}$, of the free cocommutative coalgebra without counit,

$$\odot^{\geq 1} s\mathfrak{g} := \bigoplus_{n\geq 1} \odot^n(s\mathfrak{g}) \ \subset\ \odot^\bullet s\mathfrak{g} := \bigoplus_{n\geq 0} \odot^n(s\mathfrak{g}),$$

which satisfies the condition $Q^2 = 0$. It is often very helpful to use geometric intuition and language when working with L_∞-algebras. Let us view the vector space $s\mathfrak{g}$ as a formal graded manifold (so that a choice of a basis in $\mathfrak{g}$ provides us with natural smooth coordinates on $s\mathfrak{g}$). If $\mathfrak{g}$ is finite-dimensional, then the structure ring, $\mathcal{O}_{s\mathfrak{g}}$, of formal smooth functions on the formal manifold $s\mathfrak{g}$ is equal to the completed graded commutative algebra $\widehat{\odot^\bullet}(s\mathfrak{g})^* := \prod_{n\geq 0} \odot^n(s\mathfrak{g})^*$ which is precisely the dual of the coalgebra $\odot^\bullet s\mathfrak{g}$. This dualization sends the augmentation, $\odot^{\geq 1} s\mathfrak{g}$, of the latter into the ideal, $I := \prod_{n\geq 1} \odot^n(s\mathfrak{g})^*$, of the distinguished point $0 \in s\mathfrak{g}$, while the coderivation Q into as a degree -1 derivation of $\mathcal{O}_{s\mathfrak{g}}$, i.e. into a formal vector field (denoted by the same letter Q) on the manifold $s\mathfrak{g}$ which vanishes at the distinguished point (as $QI \subset I$) and satisfies the condition $[Q,Q] = 2Q^2 = 0$. Such vector fields are often called *homological*.

In this geometric picture of L_∞-algebra structures on $\mathfrak{g}$, the subclass of dg Lie algebra structures gets represented by at most quadratic homological vector fields Q, that is $Q((s\mathfrak{g})^*) \subset (s\mathfrak{g})^* \oplus \odot^2(s\mathfrak{g})^*$ Such a vector field has a well-defined value at an arbitrary point $s\gamma \in s\mathfrak{g}$, not only at the distinguished point $0 \in s\mathfrak{g}$, i.e. it defines a smooth homological vector field on $s\mathfrak{g}$ viewed as an ordinary (non-formal) graded manifold. Given a particular dg Lie algebra $(\mathfrak{g}, d, [\ ,\ ])$, the associated homological vector field Q on $s\mathfrak{g}$ has the value at a point $s\gamma \in s\mathfrak{g}$ given explicitly by

$$Q(\gamma) := d\gamma + \frac{1}{2}[\gamma,\gamma], \tag{1}$$

where we used a canonical identification of the tangent space, $\mathcal{T}_\gamma$, at $s\gamma \in s\mathfrak{g}$ with $\mathfrak{g}$. One checks,

$$\begin{aligned} Q^2(\gamma) &= Q\left(d\gamma + \frac{1}{2}[\gamma,\gamma]\right) \\ &= -d(Q(\gamma)) + [Q(\gamma),\gamma] \\ &= -d\left(d\gamma + \frac{1}{2}[\gamma,\gamma]\right) + \left[d\gamma + \frac{1}{2}[\gamma,\gamma],\gamma\right] \\ &= 0. \end{aligned}$$

Notice that the zero locus of Q is the set of Maurer-Cartan elements in $\mathfrak{g}$.

A serious deficiency of the above geometric interpretation of L_∞-algebras is the necessity to work with the dual objects, $(\mathcal{O}_{s\mathfrak{g}}, Q)$, which make sense only for finite dimensional $\mathfrak{g}$. So we follow a suggestion of Kontsevich [Kon03] and understand from now on a *dg (smooth formal) manifold* as a pair, $(\odot^{\geq 1}X, Q)$, consisting of a cofree cocommutative algebra on a $\mathbb{Z}$-graded vector space X together with a degree -1 codifferential Q. Note that the dual of $\odot^{\geq 1}X$ is a well defined graded commutative algebra (without assumption on finite-dimensionality of X) and that dual of Q is a well-defined derivation of the latter. We identify from now on Q with its dual and call it a *homological* vector field on the dg manifold[3] X. This abuse of terminology is very helpful as it permits us to employ geometric intuition and use simple formulae of type (1) to define (in a mathematically rigorous way!) codifferentials Q on $\odot^{\geq 1}X$. Such codifferentials, $Q : \odot^{\geq 1}X \to \odot^{\geq 1}X$, are completely determined by the associated compositions,

$$Q_{proj} : \odot^{\geq 1}X \xrightarrow{Q} \odot^{\geq 1}X \xrightarrow{proj} X.$$

The restriction of Q_{proj} to $\odot^n X \subset \odot^{\geq 1}X$ is denoted by $Q^{(n)}$, $n \geq 1$.

Since we work with dual notions (coalgebras, coderivations), we will need the notion of *coideal*, which is the categorical dual to the notion of ideal. Hence, a coideal I of a coalgebra C is defined to be a quotient of C such that the kernel of the associated projection $C \twoheadrightarrow I$ is a subcoalgebra of C. For a complete study of this notion, we refer the reader to Appendix B "Categorical Algebra" of [Val06b]. This notion should not be confused with the notion of coideal used in Hopf algebra theory. Since a Hopf is an algebra and a coalgebra at the same time, a coideal in that sense is a submodule such the induced quotient carries again a bialgebra structure.

If I is a coideal of the coalgebra $\odot^{\geq 1}X$, we denote the associated sub-coalgebra of $\odot^{\geq 1}X$ by $(\mathcal{O}_I := I \setminus \odot^{\geq 1}X, Q)$. The latter is defined by the push-out diagram in the category of coalgebras,

$$\begin{tikzcd} \mathcal{O}_I \arrow[r] \arrow[d] & \odot^{\geq 1}X \arrow[d] \\ 0 \arrow[r] & I. \end{tikzcd}$$

---

[3] A warning about shift of grading: according to our definitions, a homological vector field on a graded vector space X is the same as a L_∞-structure on $s^{-1}X$.

If the coideal I is preserved by Q (i.e. admits a codifferential such that the right vertical arrow is a morphism of dg coalgebras), then the data $(\mathcal{O}_I, Q)$ is naturally a differential graded coalgebra which we often call a *dg affine scheme* (cf. [Bar06]). The coideal may not, in general, be homogeneous so the "weight" gradation, $\bigoplus_n \odot^n X$, may not survive in $\mathcal{O}_I$. A generic dg affine scheme by no means corresponds to a L_∞-algebra but, as we shall see below, some interesting examples (with non-trivial and non-homogeneous coideals) do.

A *morphism* of dg affine schemes is, by definition, a morphism of the associated dg coalgebras, $(I_1 \setminus \odot^{\geq 1} X_1, Q_1) \to (I_2 \setminus \odot^{\geq 1} X_2, Q_2)$.

7.2. **Another geometric model for a L_∞-structure.** One can interpret a L_∞-structure on a graded vector space $\mathfrak{g}$ as a linear total degree 1 polyvector field on the dual vector space $\mathfrak{g}^*$ viewed as a graded affine manifold. Note that there is no need to employ the degree shifting functors s and s^{-1} in this approach.

Indeed, let $(\wedge^\bullet \mathcal{T}_{\mathfrak{g}^*}, [\ ,\ ]_S)$ be the Schouten Lie algebra of polynomial polyvector fields on the affine manifold $\mathfrak{g}^*$. A generic total degree 1 polynomial polyvector field, $\nu = \{\nu_n \in \wedge^n \mathcal{T}_{\mathfrak{g}^*}\}_{n\geq 0}$, can be identified with a collection of its Taylor components with respect to affine coordinates on $\mathfrak{g}^*$, i.e. with a collection of linear maps,

$$\nu_{m,n} : \odot^m \mathfrak{g}^* \longrightarrow \wedge^n \mathfrak{g}^*, m \geq 0, n \geq 0,$$

of degree $n-2$. If ν is a linear polyvector field and lies in the Lie subalgebra $\wedge^{\geq 1}\mathcal{T}_{\mathfrak{g}^*}$, then only the Taylor components $\{\nu_{1,n}\}_{n\geq 1}$ can be non-zero. Their duals, $\nu_n := (\nu_{1,n})^*$, is a collection of linear maps, $\nu_n : \wedge^n \mathfrak{g} \to \mathfrak{g}$, $n \geq 1$, of degree $2-n$. It is easy to check the following

Proposition 58. *The data $\{\nu_n\}_{n\geq 1}$ defines a structure of L_∞-algebra on $\mathfrak{g}$ if and only if the linear polyvector field ν on $\mathfrak{g}^*$ satisfies the equation $[\nu, \nu]_S = 0$.*

Corollary 59. *There is a one-to-one correspondence between structures of L_∞-algebra on a finite-dimensional vector space $\mathfrak{g}$ and linear degree one polyvector fields, $\nu \in \wedge^{\geq 1}\mathcal{T}_{\mathfrak{g}^*}$, satisfying the equation $[\nu, \nu]_S = 0$.*

Kontsevich's formality morphism [Kon03], $\mathcal{F}$, associates to an arbitrary Maurer-Cartan element in the Schouten Lie algebra $\nu \in \wedge^\bullet \mathcal{T}_{\mathfrak{g}^*}$ a Maurer-Cartan element, $\mathcal{F}(\nu)$, in the Hochschild dg Lie algebra, $\oplus_{n\geq 0}\mathrm{Hom}_{poly}(\mathcal{O}_{\mathfrak{g}^*}^{\otimes n}, \mathcal{O}_{\mathfrak{g}^*})[[\hbar]]$, of polydifferential operators on the graded commutative algebra, $\mathcal{O}_{\mathfrak{g}^*} := \odot^\bullet \mathfrak{g}$, of smooth functions on the affine manifold $\mathfrak{g}^*$. If ν is a linear polyvector field on $\mathfrak{g}^*$ satisfying the equation $[\nu, \nu]_S = 0$, then one can set to zero all contributions to the formality morphism $\mathcal{F}$ coming from graphs with closed directed paths (wheels) [Sho03] and the resulting element $\mathcal{F}_{no-wheels}(\nu) \in \oplus_{n\geq 0}\mathrm{Hom}_{poly}(\mathcal{O}_{\mathfrak{g}^*}^{\otimes n}, \mathcal{O}_{\mathfrak{g}^*})[[\hbar]]$ is still Maurer-Cartan. It is easy to check that $\mathcal{F}_{no-wheels}(\nu)$ has no summand with weight $n = 0$ and hence defines an A_∞-structure on $\odot^\bullet \mathfrak{g}$ which also makes sense for $\hbar = 1$. Moreover, as $\mathcal{F}_{no-wheels}(\nu)$ involves no wheels (and hence no associated traces of linear maps), this A_∞-structure makes sense for arbitrary (not necessarily finite-dimensional) L_∞-algebra.

Definition. Let $\{\nu_n : \wedge^n \mathfrak{g} \to \mathfrak{g}\}_{n\geq 1}$ be a L_∞-structure on a graded vector space $\mathfrak{g}$. The A_∞-structure, $\mathcal{F}_{no-wheels}(\nu)$, on $\odot^\bullet \mathfrak{g}$ obtained via Kontsevich's "no-wheels" quantization of the associated linear polyvector field ν is called the *universal enveloping algebra* of the L_∞-algebra.

In a recent interesting paper [Bar07] Baranovsky also defined a universal enveloping for a L_∞-algebra $\mathfrak{g}$ as a certain A_∞-structure on the space $\odot^\bullet \mathfrak{g}$. In his approach the A_∞-structure is constructed with the help of the homological perturbation and the natural homotopy transfer of the canonical dg associative algebra structure on the cobar construction on the dg coalgebra $\odot^\bullet s\mathfrak{g}$.

7.3. **Maurer-Cartan elements in a filtered L_∞-algebra.** A L_∞-algebra $(\mathfrak{g}, Q = \{Q^{(n)}\}_{n\geq 1})$ is called *filtered* if $\mathfrak{g}$ admits a non-negative decreasing Hausdorff filtration,

$$\mathfrak{g}_0 = \mathfrak{g} \supseteq \mathfrak{g}_1 \supseteq \ldots \supseteq \mathfrak{g}_i \supseteq \ldots,$$

such that the linear map $Q^{(n)} : \odot^n(s\mathfrak{g}) \to s\mathfrak{g}$ takes values in $s\mathfrak{g}_n$ for all $n \geq n_0$ and some $n_0 \in \mathbb{N}$. In this case Q extends naturally to a coderivation of the cocommutative coalgebra, $\odot^{\geq 1} s\hat{\mathfrak{g}}$, with $\hat{\mathfrak{g}}$

being the completion of $\mathfrak g$ with respect to the topology induced by the filtration, and the equation,

$$Q(\sum_{n\geq 1}\frac{1}{n!}\gamma^{\odot n})=0,$$

for a degree zero element $\gamma\in s\hat{\mathfrak g}$ (i.e. for a degree -1 element in $\hat{\mathfrak g}$) makes sense. Its solutions are called *(generalized) Maurer-Cartan elements* (or, shortly, *MC elements*) in $(\mathfrak g, Q)$. Geometrically, an MC element is a degree -1 element in $\hat{\mathfrak g}$ at which the homological vector field Q vanishes. From now on we do not distinguish between $\mathfrak g$ and its completion $\hat{\mathfrak g}$.

To every MC element γ in a filtered L_∞-algebra $(\mathfrak g, Q)$ there corresponds, by Theorem 2.6.1 in [Mer00], a twisted L_∞-algebra, $(\mathfrak g, Q_\gamma)$, with

$$Q_\gamma(\alpha):=Q(\sum_{n\geq 0}\frac{1}{n!}\gamma^{\odot n}\odot\alpha)$$

for an arbitrary $\alpha\in\odot^{\geq 1}s\mathfrak g$. The geometric meaning of this twisted L_∞-structure is simple [Mer00]: if a homological vector field Q vanishes at a degree 0 point $\gamma\in s\mathfrak g$, then applying to Q a formal diffeomorphism, ϕ_γ, which is a translation sending γ into the origin 0 (and which is nothing but the unit shift, $e^{\mathrm{ad}\gamma}$, along the formal integral lines of the constant vector field $-\gamma$) will give us a new formal vector field, $Q_\gamma:=d\phi_\gamma(Q)$, which is *homological* and *vanishes* at the distinguished point; thus Q_γ defines a L_∞ structure on the underlying space $\mathfrak g$. In fact, we can apply this "translation diffeomorphism" trick to arbitrary (i.e. not necessarily MC) elements γ of degree 0 in $s\mathfrak g$ and get *homological* vector fields, $Q_\gamma:=d\phi_\gamma(Q)$, which do not vanish at 0 and hence define generalized L_∞ structures on $\mathfrak g$ with "zero term" $Q_\gamma^{(0)}\neq 0$.

7.4. **Extended morphisms of dg props as a dg affine scheme.** Let $(\mathcal P,\partial_{\mathcal P})$ and $(\mathcal E,\partial_{\mathcal E})$ be dg prop(erad)s with differentials $\partial_{\mathcal P}$ and $\partial_{\mathcal E}$ of degree -1. Let $\mathrm{Hom}^{\mathbb S}_\bullet(\mathcal P,\mathcal E)$ denote the graded vector space of all possible morphisms $\mathcal P\to\mathcal E$ in category of $\mathbb Z$-graded $\mathbb S$-bimodules, and let $\mathrm{Mor}(\mathcal P,\mathcal E)$ denote the set of all possible morphisms $\mathcal P\to\mathcal E$ in category of prop(erad)s, (note that we do *not* assume that elements of $\mathrm{Hom}^{\mathbb S}_\bullet(\mathcal P,\mathcal E)$ or $\mathrm{Mor}(\mathcal P,\mathcal E)$ respect differentials). It is clear that

$$\mathrm{Mor}(\mathcal P,\mathcal E)=\left\{\gamma\in\mathrm{Hom}_{\mathbb Z}(\mathcal P,\mathcal E)\,|\,\gamma\circ\mu_{\mathcal P}\left(\mathcal P\boxtimes_{(1,1)}\mathcal P\right)=\mu_{\mathcal E}\left(\gamma(\mathcal P)\boxtimes_{(1,1)}\gamma(\mathcal P)\right)\ \text{ and }|\gamma|=0\right\}.$$

We need a $\mathbb Z$-graded extension of this set,

$$\mathrm{Mor}_{\mathbb Z}(\mathcal P,\mathcal E):=\left\{\gamma\in\mathrm{Hom}_{\mathbb Z}(\mathcal P,\mathcal E)\,|\,\gamma\circ\mu_{\mathcal P}\left(\mathcal P\boxtimes_{(1,1)}\mathcal P\right)=\mu_{\mathcal E}\left(\gamma(\mathcal P)\boxtimes_{(1,1)}\gamma(\mathcal P)\right)\right\},\tag{2}$$

which we define by the same algebraic equations but dropping the assumption on the degree and homogeneity of γ.

Lemma 60. *The vector space* $\mathrm{Hom}^{\mathbb S}_\bullet(\mathcal P,\mathcal E)$ *is naturally a dg manifold.*

Proof. We define a degree -1 coderivation of the free cocommutative coalgebra, $\odot^{\geq 1}\mathrm{Hom}^{\mathbb S}_\bullet(\mathcal P,\mathcal E)$ by setting (in the dual picture, cf. § 7.1)

$$Q(\gamma):=\partial_{\mathcal E}\circ\gamma-(-1)^\gamma\gamma\circ\partial_{\mathcal P}\tag{3}$$

for an arbitrary $\gamma\in\mathrm{Hom}^{\mathbb S}_\bullet(\mathcal P,\mathcal E)$. As

$$\begin{aligned}
Q^2(\gamma) &= Q(\partial_{\mathcal E}\circ\gamma-(-1)^\gamma\gamma\circ\partial_{\mathcal P})\\
&= -\partial_{\mathcal E}\circ Q(\gamma)-(-1)^\gamma Q(\gamma)\circ\partial_{\mathcal P})\\
&= -(-1)^\gamma\partial_{\mathcal E}\circ\gamma\circ\partial_{\mathcal P}+(-1)^\gamma\partial_{\mathcal E}\circ\gamma\circ\partial_{\mathcal P}\\
&= 0,
\end{aligned}$$

Q is a linear homological field on $\mathrm{Hom}^{\mathbb S}_\bullet(\mathcal P,\mathcal E)$. (By the way, the zero locus of Q is a linear subspace of $\mathrm{Hom}^{\mathbb S}_\bullet(\mathcal P,\mathcal E)$ describing morphisms of *complexes*.) □

Proposition 61. *The set* $\mathrm{Mor}_{\mathbb Z}(\mathcal P,\mathcal E)$ *is naturally a dg affine scheme.*

Proof. Let I be the coideal in $\odot^{\geq 1}\mathrm{Hom}^{\mathbb{S}}_{\bullet}(\mathcal{P},\mathcal{E})$ cogenerated by the algebraic relations,

$$\gamma\circ\mu_{\mathcal{P}}\left(\mathcal{P}\boxtimes_{(1,1)}\mathcal{P}\right)-\mu_{\mathcal{E}}\left(\gamma(\mathcal{P})\boxtimes_{(1,1)}\gamma(\mathcal{P})\right), \tag{4}$$

on the "variable" $\gamma\in\mathrm{Hom}^{\mathbb{S}}_{\bullet}(\mathcal{P},\mathcal{E})$. The sub-coalgebra,

$$\mathcal{O}_{\mathrm{Mor}_{\mathbb{Z}}(\mathcal{P},\mathcal{E})}:=I\setminus\odot^{\geq 1}\mathrm{Hom}^{\mathbb{S}}_{\bullet}(\mathcal{P},\mathcal{E}),$$

of $\odot^{\geq 1}\mathrm{Hom}^{\mathbb{S}}_{\bullet}(\mathcal{P},\mathcal{E})$ makes the set $\mathrm{Mor}_{\mathbb{Z}}(\mathcal{P},\mathcal{E})$ into a $\mathbb{Z}$-graded affine scheme. Next we show that the homological vector field Q defined in Lemma 60 is tangent to $\mathrm{Mor}_{\mathbb{Z}}(\mathcal{P},\mathcal{E})$. Indeed, identifying Q and I with their duals (as in subsection 7.1 and the proof of Lemma 60), we have

$$\begin{aligned}Q\left(\gamma\circ\mu_{\mathcal{P}}\left(\mathcal{P}\boxtimes_{(1,1)}\mathcal{P}\right)-\mu_{\mathcal{E}}\left(\gamma(\mathcal{P})\boxtimes_{(1,1)}\gamma(\mathcal{P})\right)\right) \;=\; & Q(\gamma)\circ\mu_{\mathcal{P}}\left(\mathcal{P}\boxtimes_{(1,1)}\mathcal{P}\right)-\mu_{\mathcal{E}}\left(Q(\gamma)(\mathcal{P})\boxtimes_{(1,1)}\gamma(\mathcal{P})\right)\\ & -(-1)^{|\gamma|}\mu_{\mathcal{E}}\left(\gamma(\mathcal{P})\boxtimes_{(1,1)}Q(\gamma)(\mathcal{P})\right).\end{aligned}$$

Consistency of $\partial_{\mathcal{P}}$ and $\partial_{\mathcal{E}}$ with $\mu_{\mathcal{P}}$ and, respectively, $\mu_{\mathcal{E}}$ implies,

$$\begin{aligned}Q(\gamma)\circ\mu_{\mathcal{P}}\left(\mathcal{P}\boxtimes_{(1,1)}\mathcal{P}\right) \quad &= \quad \partial_{\mathcal{E}}\circ\gamma\circ\mu_{\mathcal{P}}\left(\mathcal{P}\boxtimes_{(1,1)}\mathcal{P}\right)-(-1)^{\gamma}\gamma\circ\partial_{\mathcal{P}}\circ\mu_{\mathcal{P}}\left(\mathcal{P}\boxtimes_{(1,1)}\mathcal{P}\right)\\ &= \quad \partial_{\mathcal{E}}\circ\gamma\circ\mu_{\mathcal{P}}\left(\mathcal{P}\boxtimes_{(1,1)}\mathcal{P}\right)-(-1)^{\gamma}\gamma\circ\mu_{\mathcal{P}}\left(\partial_{\mathcal{P}}(\mathcal{P})\boxtimes_{(1,1)}\mathcal{P}\right)\\ &\quad\quad -(-1)^{\gamma}\gamma\circ\mu_{\mathcal{P}}\left(\mathcal{P}\boxtimes_{(1,1)}\partial_{\mathcal{P}}(\mathcal{P})\right)\\ &=_{\text{mod } I} \quad \partial_{\mathcal{E}}\circ\mu_{\mathcal{E}}\left(\gamma(\mathcal{P})\boxtimes_{(1,1)}\gamma(\mathcal{P})\right)-(-1)^{\gamma}\mu_{\mathcal{E}}\left(\gamma\circ\partial_{\mathcal{P}}(\mathcal{P})\boxtimes_{(1,1)}\gamma(\mathcal{P})\right)\\ &\quad\quad -\mu_{\mathcal{E}}\left(\gamma(\mathcal{P})\boxtimes_{(1,1)}\gamma\circ\partial_{\mathcal{P}}(\mathcal{P})\right)\\ &=_{\text{mod } I} \quad \mu_{\mathcal{E}}\left(Q(\gamma)(\mathcal{P})\boxtimes_{(1,1)}\gamma(\mathcal{P})\right)+(-1)^{|\gamma|}\mu_{\mathcal{E}}\left(\gamma(\mathcal{P})\boxtimes_{(1,1)}Q(\gamma)(\mathcal{P})\right).\end{aligned}$$

Thus $Q(I)\subset I$, and hence Q gives rise to a degree -1 codifferential on the coalgebra $\mathcal{O}_{\mathrm{Mor}_{\mathbb{Z}}(\mathcal{P},\mathcal{E})}$ proving the claim. ☐

In the following theorem, we study the properties of the convolution L_∞-algebra defined in Theorem 28.

Theorem 62. *Let $(\mathcal{P}=\mathcal{F}(s^{-1}\mathcal{C}),\partial_{\mathcal{P}})$ be a quasi-free prop(erad) generated by an $\mathbb{S}$-bimodule $s^{-1}\mathcal{C}$ (so that $\mathcal{C}$ is a homotopy coprop(erad)), and let $(\mathcal{E},\partial_{\mathcal{E}})$ be an arbitrary dg prop(erad). Then*

- (i) *The graded vector space,* $\mathrm{Hom}^{\mathbb{S}}_{\bullet}(\mathcal{C},\mathcal{E})$*, is canonically a L_∞-algebra;*
- (ii) *The canonical L_∞-structure in (i) is filtered and its MC elements are morphisms, $(\mathcal{P},\partial_{\mathcal{P}})\to(\mathcal{E},\partial_{\mathcal{E}})$, of* dg *prop(erad)s;*
- (iii) *if $\partial_{\mathcal{P}}(s^{-1}\mathcal{C})\subset\mathcal{F}(s^{-1}\mathcal{C})^{(\leq 2)}$, where $\mathcal{F}(s^{-1}\mathcal{C})^{(\leq 2)}$ is the subspace of $\mathcal{F}(s^{-1}\mathcal{C})$ spanned by decorated graphs with at most two vertices, then* $\mathrm{Hom}^{\mathbb{S}}_{\bullet}(\mathcal{C},\mathcal{E})$ *is canonically a dg Lie algebra.*

Proof. (i) If $\mathcal{P}$ is free as a prop(erad), then extended morphisms from $\mathcal{P}$ to $\mathcal{Q}$ are uniquely determined by their values on the generators $s^{-1}\mathcal{C}$ so that $\mathcal{O}_{\mathrm{Mor}_{\mathbb{Z}}(\mathcal{P},\mathcal{E})}=\odot^{\geq 1}\mathrm{Hom}^{\mathbb{S}}_{\bullet}(s^{-1}\mathcal{C},\mathcal{E})$ and the claim follows from the definition of L_∞-structure in § 7.1.

(ii) The canonical L_∞ structure on $\mathrm{Hom}^{\mathbb{S}}_{\bullet}(\mathcal{C},\mathcal{E})$ is given by the restriction of the homological vector field (3) on $s\mathrm{Hom}^{\mathbb{S}}_{\bullet}(\mathcal{P},\mathcal{E})$ to the subspace $\mathrm{Hom}^{\mathbb{S}}_{\bullet}(\mathcal{C},\mathcal{E})$. This field is a formal power series in coordinates on $\mathrm{Hom}^{\mathbb{S}}_{\bullet}(\mathcal{C},\mathcal{E})$ and its part, $Q^{(n)}$, corresponding to monomials of (polynomial) degree n is given precisely by

$$Q^{(1)}(\gamma):=\partial_{\mathcal{E}}\circ\gamma-(-1)^{\gamma}\gamma\circ\partial^{(1)}_{\mathcal{P}}\quad\text{and}\quad Q^{(n)}(\gamma):=-(-1)^{\gamma}\gamma\circ\partial^{(n)}_{\mathcal{P}}\ \text{for } n>1, \tag{5}$$

where $\partial^{(n)}_{\mathcal{P}}$ is the composition[4]

$$\partial^{(n)}_{\mathcal{P}}:s^{-1}\mathcal{C}\xrightarrow{\partial_{\mathcal{P}}}\mathcal{F}(s^{-1}\mathcal{C})\xrightarrow{proj}\mathcal{F}(s^{-1}\mathcal{C})^{(n)}.$$

Note that the first summand on the r.h.s. of (3) contributes only to $Q^{(1)}$.

Define an exhaustive increasing filtration on the $\mathbb{S}$-bimodule $\mathcal{C}$ by

$$\mathcal{C}_0=0,\quad \mathcal{C}_i:=s\bigcap_{n\geq i}\mathrm{Ker}\,\partial^{(n)}_{\mathcal{P}}\ \text{for } i\geq 1,$$

---

[4] Note that for any differential $\partial_{\mathcal{P}}$ in a free properad $\mathcal{F}(s^{-1}\mathcal{C})$ the induced map $\partial^{(1)}_{\mathcal{P}}:s^{-1}\mathcal{C}\to s^{-1}\mathcal{C}$ is also a differential.

and the associated decreasing filtration on $\mathrm{Hom}_\bullet^{\mathbb{S}}(\mathcal{C},\mathcal{E})$ by

$$\mathrm{Hom}_\bullet^{\mathbb{S}}(\mathcal{C},\mathcal{E})_i := \{\gamma \in \mathrm{Hom}_\bullet^{\mathbb{S}}(\mathcal{C},\mathcal{E}) \mid \gamma(v) = 0\ \forall v \in \mathcal{C}_i\},\ i \geq 0.$$

Then, for all $n \geq 2$ and any $f_1, \ldots, f_n \in \mathrm{Hom}_\bullet^{\mathbb{S}}(\mathcal{C},\mathcal{E})$, equality (5) implies that the value of the map $Q^{(n)}(f_1, \ldots, f_n) \in s^{-1}\mathrm{Hom}_\bullet^{\mathbb{S}}(\mathcal{C},\mathcal{E})$ on arbitrary element of $\mathcal{C}_n \subset \mathrm{Ker}\,\partial_{\mathcal{P}}^{(n)}$ is equal to zero, i.e.

$$Q^{(n)}(f_1, \ldots, f_n) \in \mathrm{Hom}_\bullet^{\mathbb{S}}(\mathcal{C},\mathcal{E})_n.$$

Which in turn implies the claim that the canonical L_∞ structure on $\mathrm{Hom}_\bullet^{\mathbb{S}}(\mathcal{C},\mathcal{E})$ is filtered with respect to the constructed filtration. The claim about MC elements follows immediately from the definition (3) of the homological vector field.

(iii) As $\partial_{\mathcal{P}}^{(n)} = 0$ for $n > 2$ we conclude using formula (5) that $Q^{(n)} = 0$ for all $n > 2$. □

A special case of the above Theorem when both $\mathcal{P}$ and $\mathcal{E}$ are operads was proven earlier by van der Laan [VdL02] using different ideas.

The main point of our proof of Theorem 62 is an observation that, for a free prop(erad) $\mathcal{P} = \mathcal{F}(s^{-1}\mathcal{C})$, the set, $\mathrm{Mor}_{\mathbb{Z}}(\mathcal{P},\mathcal{E})$, of extended morphisms from $\mathcal{P}$ to an arbitrary prop(erad) $\mathcal{E}$, i.e. the set of solutions of equation (2), can be canonically identified with the graded *vector* space $s\mathrm{Hom}_\bullet^{\mathbb{S}}(\mathcal{C},\mathcal{E})$. This simple fact makes the dg affine scheme $\mathrm{Mor}_{\mathbb{Z}}(\mathcal{P},\mathcal{E})$ into a dg *smooth* manifold and hence provides us with a canonical L_∞-structure on $\mathrm{Hom}_\bullet^{\mathbb{S}}(\mathcal{C},\mathcal{E})$. A similar phenomenon occurs for the set of extended morphisms, $\mathrm{CoMor}_{\mathbb{Z}}(\mathcal{E}^c,\mathcal{P}^c)$, from an arbitrary coprop(erad) $\mathcal{E}^c$ to a cofree coprop(erad) $\mathcal{P}^c := \mathcal{F}^c(s\mathcal{E})$, and hence the arguments very similar to the ones used in the proof of Theorem 62 (and which we leave to the reader as an exercise) establish the following,

Theorem 63. *Let $(\mathcal{P}^c = \mathcal{F}^c(s\mathcal{C}), d_{\mathcal{P}})$ be a quasi-free coprop(erad), that is $\mathcal{C}$ is a homotopy prop(erad), and let $(\mathcal{E}^c, d_{\mathcal{E}})$ be an arbitrary dg coprop(erad). Then*

- (i) *The graded vector space, $\mathrm{Hom}_\bullet^{\mathbb{S}}(\mathcal{E}^c,\mathcal{C})$, is canonically a L_∞-algebra;*
- (ii) *The canonical L_∞-structure in (i) is filtered and its MC elements are morphisms, $(\mathcal{E}^c, d_{\mathcal{E}}) \to (\mathcal{P}^c, d_{\mathcal{P}})$, of* dg *coprop(erad)s;*
- (iii) *if $d_{\mathcal{P}}$ is quadratic, that is $\mathcal{C}$ is a usual prop(erad), then* $\mathrm{Hom}_\bullet^{\mathbb{S}}(\mathcal{E}^c,\mathcal{C})$ *is canonically a dg Lie algebra.*

For finite-dimensional $\mathcal{Q}$ and $\mathcal{C}$ Theorems 62 and 63 are, of course, equivalent to each other.

A morphism of L_∞-algebras, $(\mathfrak{g}_1, \mathcal{Q}_1) \to (\mathfrak{g}_2, \mathcal{Q}_2)$, is, by definition [Kon99], a morphism, $\lambda : (\odot^\bullet(s\mathfrak{g}_1), \mathcal{Q}_1) \to (\odot^\bullet(s\mathfrak{g}_2), \mathcal{Q}_2)$, of the associated dg coalgebras. It is called a *quasi-isomorphism* if the composition,

$$s\mathfrak{g}_1 \xrightarrow{i} \odot^\bullet(s\mathfrak{g}_1) \xrightarrow{\lambda} \odot^\bullet(s\mathfrak{g}_2) \xrightarrow{p} s\mathfrak{g}_2$$

induces an isomorphism, $H(s\mathfrak{g}_1, \mathcal{Q}_1^{(1)}) \to H(s\mathfrak{g}_2, \mathcal{Q}_2^{(1)})$, of the associated homology groups with respect to the *linear* (in cogenerators) parts of the codifferentials. Here i is a natural inclusion and p a natural projection.

By analogy, a map $\phi : (\mathcal{F}(s^{-1}\mathcal{C}_1), \partial_1) \to (\mathcal{F}(s^{-1}\mathcal{C}_2), \partial_2)$ of quasi-free properads is called a *tangent* quasi-isomorphism if the composition

$$s^{-1}\mathcal{C}_1 \xrightarrow{i} \mathcal{F}(s^{-1}\mathcal{C}_1) \xrightarrow{\phi} \mathcal{F}(s^{-1}\mathcal{C}_2) \xrightarrow{p} s^{-1}\mathcal{C}_2$$

induces an isomorphism of cohomology groups, $H(s^{-1}\mathcal{C}_2, \partial_2^{(1)}) \to H(s^{-1}\mathcal{C}_2, \partial_2^{(1)})$.

If we assume that properads $\mathcal{F}(s^{-1}\mathcal{C}_1)$ and $(\mathcal{F}(s^{-1}\mathcal{C}_2)$ are completed by the number of vertices (see §5.4) and that their differentials are bounded,

$$\partial_i(s^{-1}\mathcal{C}_i) \subset \mathcal{F}(s^{-1}\mathcal{C}_i)^{(\leq n_i)} \text{ for some } n_i \in \mathbb{N},\ i = 1, 2,$$

then it is not hard to show (using filtrations by the number of vertices as in §5.4 and the classical Comparison Theorem of spectral sequences) that any continuous tangent quasi-isomorphism $\phi : (\mathcal{F}(s^{-1}\mathcal{C}_1), \partial_1) \to (\mathcal{F}(s^{-1}\mathcal{C}_2), \partial_2)$ is actually a quasi-isomorphism in the ordinary sense.

Theorem 64. *(i) Let* $(\mathcal{P}_1 := \mathcal{F}(s^{-1}\mathcal{C}_1), \partial_1)$ *and* $(\mathcal{P}_2 := \mathcal{F}(s^{-1}\mathcal{C}_2), \partial_2)$ *be quasi-free prop(erad)s,* $(\mathcal{E}, \partial_\mathcal{E})$ *a dg prop(erad), and* $(\mathrm{Hom}^{\mathbb{S}}_{\bullet}(\mathcal{C}_1, \mathcal{E}), Q_1)$ *and* $(\mathrm{Hom}^{\mathbb{S}}_{\bullet}(\mathcal{C}_1, \mathcal{E}), Q_2)$ *the associated* L_∞*-algebras. Then any morphism,*

$$\phi : (\mathcal{P}_1, \partial_1) \longrightarrow (\mathcal{P}_2, \partial_2),$$

of dg prop(erad)s induces canonically an associated morphism,

$$\phi_{ind} : (\mathrm{Hom}^{\mathbb{S}}_{\bullet}(\mathcal{C}_2, \mathcal{E}), Q_2) \longrightarrow (\mathrm{Hom}^{\mathbb{S}}_{\bullet}(\mathcal{C}_1, \mathcal{E}), Q_1)$$

of L_∞*-algebras. Moreover, if* ϕ *is a tangent quasi-isomorphism of dg prop(erad)s, then* ϕ_{ind} *is a quasi-isomorphism of* L_∞*-algebras.*

(ii) Let $(\mathcal{P} := \mathcal{F}(s^{-1}\mathcal{C}), \partial)$ *be a quasi-free prop(erad),* $(\mathcal{E}_1, \partial_{\mathcal{E}_1})$ *and* $(\mathcal{E}_2, \partial_{\mathcal{E}_2})$ *arbitrary dg prop(erad)s, and* $(\mathrm{Hom}^{\mathbb{S}}_{\bullet}(\mathcal{C}, \mathcal{E}_1), Q_1)$ *and* $(\mathrm{Hom}^{\mathbb{S}}_{\bullet}(\mathcal{C}, \mathcal{E}_2), Q_2)$ *the associated* L_∞*-algebras. Then any morphism,*

$$\psi : (\mathcal{E}_1, \partial_{\mathcal{E}_1}) \longrightarrow (\mathcal{E}_2, \partial_{\mathcal{E}_2}),$$

of dg prop(erad)s induces canonically an associated morphism,

$$\psi_{ind} : (\mathrm{Hom}^{\mathbb{S}}_{\bullet}(\mathcal{C}, \mathcal{E}_1), Q_1) \longrightarrow (\mathrm{Hom}^{\mathbb{S}}_{\bullet}(\mathcal{C}, \mathcal{E}_2), Q_2)$$

of L_∞*-algebras. Moreover, if* ψ *is a quasi-isomorphism of dg prop(erad)s, then* ψ_{ind} *is a quasi-isomorphism of* L_∞*-algebras.*

Proof. (i) The map ϕ induces a degree 0 linear map,

$$\begin{array}{ccc} \mathrm{Hom}^{\mathbb{S}}_{\bullet}(\mathcal{P}_2, \mathcal{E}) & \longrightarrow & \mathrm{Hom}^{\mathbb{S}}_{\bullet}(\mathcal{P}_1, \mathcal{E}) \\ \gamma & \longrightarrow & \gamma \circ \phi \end{array}$$

Using definition (3) of the codifferentials Q_1 and Q_2, and the fact that ϕ respects differentials ∂_1 and ∂_2, we obtain, for any $\gamma \in \mathrm{Hom}^{\mathbb{S}}_{\bullet}(\mathcal{P}_2, \mathcal{E})$,

$$\begin{aligned} Q_1(\gamma \circ \phi) &= \partial_\mathcal{E} \circ \gamma \circ \phi - (-1)^\gamma \gamma \circ \phi \circ \partial_1 \\ &= \partial_\mathcal{E} \circ \gamma \circ \phi - (-1)^\gamma \gamma \circ \partial_2 \circ \phi \\ &= Q_2(\gamma) \circ \phi, \end{aligned}$$

and hence conclude that ϕ induces a morphism of dg coalgebras,

$$\phi_{ind} : (\odot^{\geq 1}\mathrm{Hom}^{\mathbb{S}}_{\bullet}(\mathcal{P}_2, \mathcal{E}), Q_2) \longrightarrow (\odot^{\geq 1}\mathrm{Hom}^{\mathbb{S}}_{\bullet}(\mathcal{P}_1, \mathcal{E}), Q_1).$$

As

$$\phi \circ \mu_{\mathcal{P}_1}\left(\mathcal{P}_1 \boxtimes_{(1,1)} \mathcal{P}_1\right) = \mu_{\mathcal{P}_2}\left(\phi(\mathcal{P}_1) \boxtimes_{(1,1)} \phi(\mathcal{P}_1)\right) \subset \mu_{\mathcal{P}_2}\left(\mathcal{P}_2 \boxtimes_{(1,1)} \mathcal{P}_2\right)$$

we have

$$\begin{aligned} &\gamma \circ \phi \circ \mu_{\mathcal{P}_1}\left(\mathcal{P}_1 \boxtimes_{(1,1)} \mathcal{P}_1\right) - \mu_\mathcal{E}\left(\gamma \circ \phi(\mathcal{P}_1) \boxtimes_{(1,1)} \gamma \circ \phi(\mathcal{P}_1)\right) \subset \\ &\qquad \subset \gamma \circ \mu_{\mathcal{P}_2}\left(\mathcal{P}_2 \boxtimes_{(1,1)} \mathcal{P}_2\right) - \mu_\mathcal{E}\left(\gamma(\mathcal{P}_2) \boxtimes_{(1,1)} \gamma(\mathcal{P}_1)\right). \end{aligned}$$

Thus the map ϕ_{ind} sends cogenerators (4) of the coideal I_2 in $\odot^{\geq 1}\mathrm{Hom}^{\mathbb{S}}_{\bullet}(\mathcal{P}_2, \mathcal{E})$ into cogenerators of the coideal I_1 in $\odot^{\geq 1}\mathrm{Hom}^{\mathbb{S}}_{\bullet}(\mathcal{P}_1, \mathcal{E})$, and hence gives rise to a morphism of dg coalgebras,

$$\phi_{ind} : \left(\mathcal{O}_{\mathrm{Mor}_\mathbb{Z}(\mathcal{P}_2, \mathcal{E})}, Q_2\right) \longrightarrow \left(\mathcal{O}_{\mathrm{Mor}_\mathbb{Z}(\mathcal{P}_1, \mathcal{E})}, Q_1\right),$$

i.e. to a morphism of dg affine schemes, $\phi_{ind} : (\mathrm{Mor}_\mathbb{Z}(\mathcal{P}_2, \mathcal{E}), Q_2) \to (\mathrm{Mor}_\mathbb{Z}(\mathcal{P}_1, \mathcal{E}), Q_1)$.

If the dg prop(erad)s $\mathcal{P}_1$ and $\mathcal{P}_2$ are quasi-free, then the above morphism of dg affine schemes is the same as a morphism of smooth dg manifolds, i.e. a morphism,

$$\phi_{ind} : (\mathrm{Hom}^{\mathbb{S}}_{\bullet}(\mathcal{C}_2, \mathcal{E}), Q_2) \longrightarrow (\mathrm{Hom}^{\mathbb{S}}_{\bullet}(\mathcal{C}_1, \mathcal{E}), Q_1),$$

of L_∞-algebras. The last statement of Theorem 64 follows immediately from the formulae (5) for $n = 1$ and the Künneth formula completing the proof of Claim (i).

Claim (ii) is much easier than Claim (i): it follows directly from the formulae (5) for $n = 1$. □

7.5. **Proofs via local coordinate computations.** Calculations in local coordinates is a powerful and useful tool in differential geometry. In this section we show a new proofs of Lemma 60, Proposition 61 and Theorem 62 by explicitly describing all the notions and constructions of §7.4 in local coordinates and justifying thereby the geometric language we used in that section. For simplicity, we show the proofs only for the case when $(\mathcal{P}, \partial_P)$ and $(\mathcal{E}, \partial_{\mathcal{E}})$ are dg associative algebras, that is, dg properads concentrated in biarity $(1,1)$ (a generalization to arbitrary dg (prop)erads is straightforward); moreover, to simplify Koszul signs in the formulae below we also assume that both $\mathcal{P}$ and $\mathcal{E}$ are free modules over a graded commutative ring, $R = \oplus_{i\in\mathbb{Z}} R^i$, with degree 0 generators $\{e_a\}_{a\in I}$ and, respectively, $\{e_\alpha\}_{\alpha\in J}$. Then multiplications and differentials in $\mathcal{P}$ and $\mathcal{E}$ have the following coordinate representations,

$$e_a \cdot e_b = \sum_{c\in I} \mu_{ab}^c e_c, \quad e_\alpha \cdot a_\beta = \sum_{\gamma\in J} \mu_{\alpha\beta}^\gamma e_\gamma,$$

$$\partial_{\mathcal{P}} e_a = \sum_{b\in I} D_a^b e_b, \quad \partial_{\mathcal{E}} e_\alpha = \sum_{\beta\in J} D_\alpha^\beta e_\beta,$$

for some coefficients $\mu_{ab}^c, \mu_{\alpha\beta}^\gamma \in R^0$ and $D_a^b, D_\alpha^\beta \in R^{-1}$. Equations $\partial_{\mathcal{P}}^2 = \partial_{\mathcal{E}}^2 = 0$ as well as equations for compatibility of differentials with products are given in coordinates as follows,

$$\sum_{b\in I} D_a^b D_b^c = 0, \qquad \sum_{\beta\in J} D_\alpha^\beta D_\beta^\gamma = 0, \tag{6}$$

$$D_a^m \mu_{mb}^c + D_b^m \mu_{am}^c = \mu_{ab}^m D_m^c, \qquad D_\alpha^\nu \mu_{\nu\beta}^\gamma + D_\beta^\nu \mu_{\alpha\nu}^\gamma = \mu_{\alpha\beta}^\nu D_\nu^\gamma. \tag{7}$$

A generic homogeneous map of graded vector space, $\gamma : \mathcal{P} \to \mathcal{E}$, of degree $i \in \mathbb{Z}$ is unique;y determined by its values on the generators,

$$\gamma(e_a) = \sum_{\alpha\in J} \gamma_{a(i)}^\alpha e_\alpha,$$

for some coefficients $\gamma_{a(i)}^\alpha \in R^i$. We shall understand these coefficients as coordinates on the flat manifold $\mathrm{Hom}_\bullet^{\mathbb{S}}(\mathcal{P}, \mathcal{E})$.

Consider now a completed free graded commutative algebra, $R[[\gamma_{a(i)}^\alpha]]$, generated by formal variables $\gamma_{a(i)}^\alpha$ to which we assign degree i. This algebra is precisely the algebra of smooth functions, $\mathcal{O}_{\mathrm{Hom}_\bullet^{\mathbb{S}}(\mathcal{P},\mathcal{E})}$, on the manifold $\mathrm{Hom}_\bullet^{\mathbb{S}}(\mathcal{P}, \mathcal{E})$. Let us consider a degree -1 vector field (that is, a derivation of $\mathcal{O}_{\mathrm{Hom}_\bullet^{\mathbb{S}}(\mathcal{P},\mathcal{E})}$),

$$Q = \left(\sum_{\alpha,\beta,a,i} D_\beta^\alpha \gamma_{a(i)}^\beta - \sum_{a,b,\alpha,i} (-1)^i D_a^b \gamma_{b(i)}^\alpha \right) \frac{\partial}{\partial \gamma_{a(i)}^\alpha},$$

on $\mathrm{Hom}_\bullet^{\mathbb{S}}(\mathcal{P}, \mathcal{E})$. In view of (6), we have

$$\begin{aligned} [Q,Q] &= 2\left(\sum_{\alpha,\beta,\gamma,a,i} -D_\beta^\alpha D_\gamma^\beta \gamma_{a(i)}^\gamma - \sum_{\alpha,\beta,a,b,i} (-1)^i D_a^b D_\beta^\alpha \gamma_{b(i)}^\beta \right) \frac{\partial}{\partial \gamma_{a(i)}^\alpha} \\ &\quad +2\left(- \sum_{\alpha,a,b,c,i} (-1)^i D_\beta^\alpha D_a^b \gamma_{b(i)}^\beta - \sum_{\alpha,\beta,\gamma,a,i} D_a^b D_b^c \gamma_{c(i)}^\alpha \right) \frac{\partial}{\partial \gamma_{a(i)}^\alpha} \\ &= 0, \end{aligned}$$

proving thereby Lemma 60 which claims that $(\mathrm{Hom}_\bullet^{\mathbb{S}}(\mathcal{P}, \mathcal{E}), Q)$ is a dg manifold.

The space of extended morphisms of associative R-algebras $\mathrm{Mor}_{\mathbb{Z}}(\mathcal{P}, \mathcal{E})$ is, by its definition, a (singular, in general) subspace of the manifold $\mathrm{Hom}_\bullet^{\mathbb{S}}(\mathcal{P}, \mathcal{E})$ given explicitly by the following equations,

$$\mathrm{Mor}_{\mathbb{Z}}(\mathcal{P}, \mathcal{E}) := \left\{ \gamma_{a(i)}^\alpha \in \mathrm{Hom}_\bullet^{\mathbb{S}}(\mathcal{P}, \mathcal{E}) : \sum_{c\in I} \mu_{ab}^c \gamma_{c(i)}^\alpha - \sum_{\substack{\beta,\gamma\in J \\ j+k=i}} \mu_{\beta\gamma}^\alpha \gamma_{a(j)}^\beta \gamma_{b(k)}^\gamma = 0 \right\}.$$

Let I be an ideal in $\mathcal{O}_{\mathrm{Hom}^{\mathbb{S}}_{\bullet}(\mathcal{P},\mathcal{E})} = \mathbb{K}[[\gamma^{\alpha}_{a(i)}]]$ generated by the functions $\{\sum_{c\in I}\mu^{c}_{ab}\gamma^{\alpha}_{c(i)} - \sum_{\substack{\beta,\gamma\in J\\ j+k=i}}\mu^{\alpha}_{\beta\gamma}\gamma^{\beta}_{a(j)}\gamma^{\gamma}_{b(k)}\}$. Then the structure sheaf, $\mathcal{O}_{\mathrm{Mor}_{\mathbb{Z}}(\mathcal{P},\mathcal{E})}$, of the scheme $\mathrm{Mor}_{\mathbb{Z}}(\mathcal{P},\mathcal{E})$ is given, by definition, by the quotient algebra $\mathcal{O}_{\mathrm{Hom}^{\mathbb{S}}_{\bullet}(\mathcal{P},\mathcal{E})}/I$ (which, in general, is not freely generated, i.e. is not smooth). We claim that the vector field Q on $\mathrm{Hom}^{\mathbb{S}}_{\bullet}(\mathcal{P},\mathcal{E})$ ia tangent to the subspace $\mathrm{Mor}_{\mathbb{Z}}(\mathcal{P},\mathcal{E})$. Indeed, in view of (7), we have

$$
\begin{aligned}
Q\left(\sum_{c\in I}\mu^{c}_{ab}\gamma^{\alpha}_{c(i)} - \sum_{\substack{\beta,\gamma\in J\\ j+k=i}}\mu^{\alpha}_{\beta\gamma}\gamma^{\beta}_{a(j)}\gamma^{\gamma}_{b(k)}\right) \quad = \quad & \sum_{\sigma\in J} D^{\alpha}_{\sigma}\left(\sum_{c\in I}\mu^{c}_{ab}\gamma^{\sigma}_{c(i)} - \sum_{\substack{\beta\gamma\in J\\ j+k=i}}\mu^{\sigma}_{\beta\gamma}\gamma^{\beta}_{a(j)}\gamma^{\gamma}_{b(k)}\right)\\
& -(-1)^{i}\sum_{e\in E} D^{m}_{a}\left(\sum_{c\in I}\mu^{c}_{mb}\gamma^{\sigma}_{c(i)} - \sum_{\substack{\beta\gamma\in J\\ j+k=i}}\mu^{\sigma}_{\beta\gamma}\gamma^{\beta}_{m(j)}\gamma^{\gamma}_{b(k)}\right)\\
& -(-1)^{i}\sum_{e\in E} D^{m}_{b}\left(\sum_{c\in I}\mu^{c}_{am}\gamma^{\sigma}_{c(i)} - \sum_{\substack{\beta\gamma\in J\\ j+k=i}}\mu^{\sigma}_{\beta\gamma}\gamma^{\beta}_{a(j)}\gamma^{\gamma}_{m(k)}\right)
\end{aligned}
$$

Thus $Q(I)\subset I$ so that Q makes $\mathcal{O}_{\mathrm{Mor}_{\mathbb{Z}}(\mathcal{P},\mathcal{E})}$ into a *differential* graded algebra proving thereby Proposition 61.

To prove Theorem 62 we have to assume from now on that $\mathcal{P}$ is a free algebra, $\otimes^{\bullet}V$, generated by some free R-module V. Let $\{e_A\}_{A\in K}$ stand for a set of generators of V so that the basis $\{e_a\}$ we used above can be identified with the following set,

$$
\{e_a\}_{a\in I} = \{e_A,\ e_{A_1A_2} := e_{A_1}\otimes e_{A_2},\ e_{A_1A_2A_3} = e_{A_1}\otimes e_{A_2}\otimes e_{A_3},\ \ldots\}_{A_{\bullet}\in K}.
$$

The differential $\partial_{\mathcal{P}}$ is now completely determined by its values on the generators $\{e_A\}$,

$$
\partial_{\mathcal{P}} e_A = \sum_{k\geq 1}\ \sum_{A_1,\ldots,A_k\in K} D^{A_1\ldots A_k}_{A}\, e_{A_1\ldots A_k},
$$

for some coefficients $D^{A_1\ldots A_k}_{A}\in R^{-1}$. On the other hand, the R-algebra of smooth formal functions on the manifold $\mathrm{Hom}^{\mathbb{S}}_{\bullet}(\mathcal{P},\mathcal{E})$ gets the following explicit representation,

$$
\mathcal{O}_{\mathrm{Hom}^{\mathbb{S}}_{\bullet}(\mathcal{P},\mathcal{E})} = R[[\gamma^{\alpha}_{A(i)},\ \gamma^{\alpha}_{A_1A_2(i)},\ \gamma^{\alpha}_{A_1A_2A_3(i)},\ldots]].
$$

The key point is that the system of equations defining the subspace $\mathrm{Mor}_{\mathbb{Z}}(\mathcal{P},\mathcal{E})\subset\mathrm{Hom}^{\mathbb{S}}_{\bullet}(\mathcal{P},\mathcal{E})$ can now be easily solved,

$$
\begin{aligned}
\gamma^{\alpha}_{A_1A_2(i)} \quad &= \quad \sum_{\substack{\beta_1,\beta_2\in J\\ i_1+i_2=i}}\mu^{\alpha}_{\beta_1\beta_2}\gamma^{\beta_1}_{A_1(i_1)}\gamma^{\beta_2}_{A_2(i_2)},\\
\gamma^{\alpha}_{A_1A_2A_3(i)} \quad &= \quad \sum_{\substack{\beta_{\bullet},\gamma\in J\\ i_1+i_2+i_3=i}}\mu^{\alpha}_{\beta_1\gamma}\mu^{\gamma}_{\beta_2\beta_3}\gamma^{\beta_1}_{A_1(i_1)}\gamma^{\beta_2}_{A_2(i_2)}\gamma^{\beta_3}_{A_3(i_3)},,\\
&\ldots
\end{aligned}
$$

in terms of the independent variables $\gamma^{\alpha}_{A(i)}$. Thus $\mathrm{Mor}_{\mathbb{Z}}(\mathcal{P},\mathcal{E})$ is itself a smooth formal manifold with the structure sheaf $\mathcal{O}_{\mathrm{Mor}_{\mathbb{Z}}(\mathcal{P},\mathcal{E})}\simeq R[[\gamma^{\alpha}_{A(i)}]]$. The vector field Q on the manifold $\mathrm{Hom}^{\mathbb{S}}_{\bullet}(\mathcal{P},\mathcal{E})$ restricts to a smooth degree -1 homological vector field on the subspace $\mathrm{Mor}_{\mathbb{Z}}(\mathcal{P},\mathcal{E})\subset\mathrm{Hom}^{\mathbb{S}}_{\bullet}(\mathcal{P},\mathcal{E})$ which is given explicitly as follows,

$$
Q|_{\mathrm{Mor}_{\mathbb{Z}}(\mathcal{P},\mathcal{E})} = \left(\sum_{\alpha,\beta,a,i} D^{\alpha}_{\beta}\gamma^{\beta}_{A(i)} - \sum_{A,A_{\bullet},\alpha,i}(-1)^{i} D^{A_1\ldots A_k}_{A}\gamma^{\alpha}_{A_1\ldots A_k(i)}\right)\frac{\partial}{\partial\gamma^{\alpha}_{A(i)}}
$$

where, for $k\geq 2$,

$$
\gamma^{\alpha}_{A_1A_2\ldots A_k(i)} = \sum_{\substack{\beta_{\bullet},\gamma_{\bullet}\in J\\ i_1+\ldots+i_k=i}}\mu^{\alpha}_{\beta_1\gamma_1}\mu^{\gamma_1}_{\beta_2\gamma_2}\ldots\mu^{\gamma_{k-2}}_{\beta_{k-1}\beta_k}\gamma^{\beta_1}_{A_1(i_1)}\gamma^{\beta_2}_{A_2(i_2)}\ldots\gamma^{\beta_k}_{A_k(i_k)}.
$$

Thus $\mathrm{Mor}_{\mathbb{Z}}(\mathcal{P},\mathcal{E})=\mathrm{Hom}_{\mathbb{Z}}(V,\mathcal{E})$ is canonically a dg manifold, i.e. $s\mathrm{Hom}_{\mathbb{Z}}(V,\mathcal{E})$ is canonically a L_∞-algebra, and Theorem 62(i) is proved. Theorem 62(ii) follows from the above explicit expression for the homological vector field $Q|_{\mathrm{Mor}_{\mathbb{Z}}(\mathcal{P},\mathcal{E})}$ as its zero set is precisely the set of morphisms $\mathcal{P}\rightarrow\mathcal{E}$ which commute with the differentials. Finally, if the differential $\partial_{\mathcal{P}}$ is at most quadratic in generators, then $D_A^{A_1\ldots A_k}=0$ for $k\geq 3$ and hence $Q|_{\mathrm{Mor}_{\mathbb{Z}}(\mathcal{P},\mathcal{E})}$ is evidently at most quadratic homological vector field so that Theorem 62(iii) is also done.
In a similar purely geometric way one can prove anew Theorem 64. We leave the details as an exercise to the interested reader.

7.6. **Enlarged category of dg prop(erad)s.** For any dg prop(erad)s $(\mathcal{P}_1,\partial_1)$, $(\mathcal{P}_2,\partial_2)$ and $(\mathcal{P}_3,\partial_3)$, the natural composition map,

$$\begin{array}{ccc}\mathrm{Hom}_\bullet^{\mathbb{S}}(\mathcal{P}_2,\mathcal{P}_3)\otimes\mathrm{Hom}_\bullet^{\mathbb{S}}(\mathcal{P}_1,\mathcal{P}_2) & \longrightarrow & \mathrm{Hom}_\bullet^{\mathbb{S}}(\mathcal{P}_1,\mathcal{P}_3)\\ \gamma_2\otimes\gamma_1 & \longrightarrow & \gamma_2\circ\gamma_1\end{array}$$

respects the relations (4) and hence induces a map of coalgebras (cf. the proof of Theorem 64(i)),

$$\circ:\mathcal{O}_{Mor_{\mathbb{Z}}(\mathcal{P}_2,\mathcal{P}_3)}\otimes\mathcal{O}_{\mathrm{Mor}_{\mathbb{Z}}(\mathcal{P}_1,\mathcal{P}_2)}\longrightarrow\mathcal{O}_{Mor_{\mathbb{Z}}(\mathcal{P}_1,\mathcal{Q})}.$$

Proposition 65. *The map $\circ$ respects the codifferentials (3), i.e. induces a morphism of dg affine schemes,*

$$(\mathrm{Mor}_{\mathbb{Z}}(\mathcal{P}_2,\mathcal{P}_3),Q_{23})\times(\mathrm{Mor}_{\mathbb{Z}}(\mathcal{P}_1,\mathcal{P}_2),Q_{12})\longrightarrow(\mathrm{Mor}_{\mathbb{Z}}(\mathcal{P}_1,\mathcal{P}_3),Q_{13})\,.$$

Proof. We have, for any $\gamma_1\in\mathrm{Hom}_\bullet^{\mathbb{S}}(\mathcal{P}_1,\mathcal{P}_2)$ and $\gamma_2\in\mathrm{Hom}_\bullet^{\mathbb{S}}(\mathcal{P}_2,\mathcal{P}_3)$,

$$\begin{array}{rcl}Q_{13}(\gamma_2\circ\gamma_1) & \overset{\mathrm{by}(3)}{=} & \partial_3\circ\gamma_2\circ\gamma_1-(-1)^{\gamma_1+\gamma_2}\gamma_2\circ\gamma_1\circ\partial_1\\ & = & \partial_3\circ\gamma_2\circ\gamma_1-(-1)^{\gamma_2}\gamma_2\circ\partial_2\circ\gamma_1\\ & & +(-1)^{\gamma_2}\circ\partial_2\circ\gamma_1-(-1)^{\gamma_1+\gamma_2}\gamma_2\circ\gamma_1\circ\partial_1\\ & = & Q_{23}(\gamma_2)\circ\gamma_1+(-1)^{\gamma_2}\gamma_2\circ Q_{12}(\gamma_1).\end{array}$$

□

As the composition $\circ$ is obviously associative, we end up with the following canonical enlargement of the category of dg prop(erad)s.

Corollary 66. *The data,*

$$\begin{array}{rcl}\mathsf{Objects} & := & \textit{dg prop(erad)s},\\ \mathsf{Hom}(\mathcal{P}_1,\mathcal{P}_2) & := & \textit{the dg affine scheme }\ (\mathrm{Mor}_{\mathbb{Z}}(\mathcal{P}_1,\mathcal{P}_2),Q_{12})\end{array}$$

is a category. Moreover, the composition,

$$\circ:\mathsf{Hom}(\mathcal{P}_2,\mathcal{P}_3)\times\mathsf{Hom}(\mathcal{P}_1,\mathcal{P}_2)\longrightarrow\mathsf{Hom}(\mathcal{P}_1,\mathcal{P}_3)$$

is a morphism of dg affine schemes.

Note that if $\mathcal{P}_1$ is quasi-free then, by Theorem 62, $\mathsf{Hom}(\mathcal{P}_1,\mathcal{P}_2)$ is precisely the filtered L_∞-algebra whose Maurer-Cartan elements are ordinary morphisms of dg prop(erad)s from $\mathcal{P}_1$ to $\mathcal{P}_2$. Note also that if $\phi:\mathcal{P}_1\rightarrow\mathcal{P}_2$ is an ordinary morphism of quasi-free dg prop(erad)s, then the composition map,

$$\circ:\mathsf{Hom}(\mathcal{P}_2,\mathcal{P}_3)\times\phi\longrightarrow\mathsf{Hom}(\mathcal{P}_1,\mathcal{P}_3)$$

is precisely the L_∞-morphism of Theorem 64(i).

7.7. **Families of natural L_∞-structures on $\oplus\mathcal{P}$.** It was shown in Section 4.5 that for any homotopy properad $\mathcal{P}$ the associated direct sum $\oplus\mathcal{P} := \bigoplus_{m,n}\mathcal{P}(m,n)$ has a natural structure of L_∞-algebra which encodes all possible compositions in $\mathcal{P}$. In this section we show a new proof of this result which is independent of Section 4.5 and the earlier works [KM01, VdL02] which treated the special case of operads. The present approach is based on certain universal properties of the properad of Frobenius algebras (and its non-commutative versions) and Theorem 62; it provides a conceptual explanation of the phenomenon in terms of convolution properads.

Theorem 67. *Let $\mathcal{P} = \{P(m,n)\}$ be a homotopy prop(erad). Then*

(i) $\bigoplus_{m,n} P(m,n)$ *is canonically a L_∞-algebra;*
(ii) $\bigoplus_{m,n} P(m,n)^{\mathbb{S}_m}$ *is canonically a L_∞-algebra;*
(iii) $\bigoplus_{m,n} P(m,n)^{\mathbb{S}_n}$ *is canonically a L_∞-algebra;*
(iv) $\bigoplus_{m,n} P(m,n)^{\mathbb{S}_m\times\mathbb{S}_n}$ *is canonically a L_∞-algebra;*
(v) *there is a natural commutative diagram of L_∞-morphisms,*

$$\begin{tikzcd}
 & \bigoplus_{m,n} P(m,n)^{\mathbb{S}_m} \arrow[dr] & \\
\bigoplus_{m,n} P(m,n) \arrow[ur] \arrow[dr] & & \bigoplus_{m,n} P(m,n)^{\mathbb{S}_m\times\mathbb{S}_n} \\
 & \bigoplus_{m,n} P(m,n)^{\mathbb{S}_n} \arrow[ur] &
\end{tikzcd}$$

Finally, if $\mathcal{P}$ is a dg properad, then all the above data are dg Lie algebras and morphisms of dg Lie algebras.

Proof. Recall that the prop(erad) of Frobenius algebras can be defined it as a quotient,

$$\mathcal{F}rob := \mathcal{F}\langle V\rangle/(R)$$

of the free prop(erad), $\mathcal{F}(V)$, generated by the $\mathbb{S}$-bimodule $V = \{V(m,n)\}$,

$$V(m,n) := \begin{cases} \mathrm{Id}_2 \otimes \mathrm{Id}_1 \equiv \mathrm{span}\left\langle \quad = \quad \right\rangle & \text{if } m=2, n=1,\\ \mathrm{Id}_1 \otimes \mathrm{Id}_2 \equiv \mathrm{span}\left\langle \quad = \quad \right\rangle & \text{if } m=1, n=2,\\ 0 & \text{otherwise}\end{cases}$$





modulo the ideal generated by relations



Here Id_n stands for the trivial one dimensional representation of the group $\mathbb{S}_n$. It is clear that $\mathcal{F}rob(m,n) = \mathrm{Id}_m \otimes \mathrm{Id}_n$ and the compositions in $\mathcal{F}rob$ are determined by the canonical isomorphism $\mathbb{K}\otimes\mathbb{K}\to\mathbb{K}$ (thus $\mathcal{F}rob$ is a prop(erad)ic analogue of $\mathcal{C}om$ in the theory of operads). The dual space, $\mathcal{F}rob^*$, is naturally a coprop(erad) [5]. Homotopy prop(erad) structure on $\mathcal{P}$ is the same as a differential, $d_{\mathcal{P}}$, in the free coprop(erad) $\mathcal{F}^c(s\bar{\mathcal{P}})$. Theorem 63(i) applied to the coprop(erad)s $\mathcal{F}rob^*$ and $(\mathcal{F}^c(s\bar{\mathcal{P}}), d_{\mathcal{P}})$, asserts that the vector space

---

[5] In fact, $\mathcal{F}rob^*$ is a completed coproperad with respect to the topology induced by the number of vertices. The formulae for the composite coproduct in infinite. But since we 'dualize' it by considering the convolution homotopy properad $\mathrm{Hom}^{\mathbb{S}}(\mathcal{F}rob^*, -)$ it does not matter.

$$\mathrm{Hom}^{\mathbb{S}}_{\bullet}(\mathcal{F}rob^*, \mathcal{P}) = \bigoplus_{m,n} (\mathrm{Id}_m \otimes \mathrm{Id}_n) \otimes_{\mathbb{S}_m \times \mathbb{S}_n} \mathcal{P}(m,n) = \bigoplus_{m,n} \mathcal{P}(m,n)^{\mathbb{S}_m \times \mathbb{S}_n}$$

is canonically a L_∞-algebra. Hence the claim (iv).

Let us next define a non-commutative analogue of $\mathcal{F}rob$ as a quotient,

$$\mathcal{F}rob_+^+ := \mathcal{F}(V)/(R)$$

of the free prop(erad), $\mathcal{F}(V)$, generated by the $\mathbb{S}$-bimodule $V = \{V(m,n)\}$,

$V(m,n) :=$

$\mathbb{K}[\mathbb{S}_2] \otimes \mathrm{Id}_1 \equiv \mathrm{span}$  if $m=2, n=1$,

$\mathrm{Id}_1 \otimes \mathbb{K}[\mathbb{S}_2] \equiv \mathrm{span}$  if $m=1, n=2$,

0 otherwise

modulo the ideal generated by relations





It is clear that $\mathcal{F}rob_+^+(m,n) = \mathbb{K}[\mathbb{S}_m] \otimes \mathbb{K}[\mathbb{S}_n]$. Analogously one defines two other versions of $\mathcal{F}rob$,

$$\mathcal{F}rob_+ = \{\mathcal{F}rob_+(m,n) = \mathrm{Id}_m \otimes \mathbb{K}[\mathbb{S}_n]\} \quad \text{and} \quad \mathcal{F}rob^+ = \{\mathcal{F}rob_+(m,n) = \mathbb{K}[\mathbb{S}_m] \otimes \mathrm{Id}_n\},$$

with comultiplication (resp. multiplication) commutative but multiplication (resp. comultiplication) noncommutative. Then applying again Theorem 63(i) or Theorem 27 to $\mathcal{Q}$ being $(\mathcal{F}rob_+^+)^*$, $(\mathcal{F}rob_+)^*$ or $(\mathcal{F}rob^+)^*$ and $\mathcal{D}$ being $\mathcal{F}^c(s\bar{\mathcal{P}})$ we conclude that the vector spaces,

$$\mathrm{Hom}^{\mathbb{S}}_{\bullet}\big((\mathcal{F}rob_+^+)^*, \mathcal{P}\big) = \bigoplus_{m,n} \mathcal{P}(m,n),$$

$$\mathrm{Hom}^{\mathbb{S}}_{\bullet}\big((\mathcal{F}rob_+)^*, \mathcal{P}\big) = \bigoplus_{m,n} \mathcal{P}(m,n)^{\mathbb{S}_m},$$

$$\mathrm{Hom}^{\mathbb{S}}_{\bullet}\big((\mathcal{F}rob^+)^*, \mathcal{P}\big) = \bigoplus_{m,n} \mathcal{P}(m,n)^{\mathbb{S}_n},$$

admit canonically L_∞-structures proving thereby Claim (i)-(iii).

Finally, the natural commutative diagram of morphisms of properads,

$$\begin{tikzcd} & \mathcal{F}rob_+ \arrow[dr] & \\ \mathcal{F}rob \arrow[ur] \arrow[dr] & & \mathcal{F}rob_+^+ \\ & \mathcal{F}rob^+ \arrow[ur] & \end{tikzcd}$$

proves claim (v). □

The prop(erad) $\mathcal{F}rob_+^+$ was designed so that it is generated as an $\mathbb{S}$-module by arbitrary (m,n)-corollas, and the comultiplication $\Delta_{(1,1)}$ in its dual, $(\mathcal{F}rob_+^+)^*$ splits such a corolla into *all* possible two vertex (m,n)-graphs. Hence the L_∞-structure claimed in Theorem 67 is exactly the same as in Theorem 25.

The L_∞ structures on the direct sum $\oplus P$ and its subspaces of invariants constructed in the proof of Theorem 67 are the most natural ones to consider as they involve *all* possible compositions in $\mathcal{P}$. However, they are by no means unique in the case of prop(erad)s (as opposite to the case of operads). For example, the part of prop(erad) compositions which correspond to so called $\frac{1}{2}$-propic graphs or dioperadic graphs, that is graphs of genus 0, (see pictures (8) below) also combine into a L_∞-structure on $\oplus P$ (and its subspaces of invariants) as the following argument shows.

For a prop(erad) $\mathcal{P} = \{\mathcal{P}(m,n)\}$ we denote by $\mathcal{P}^\dagger = \{\mathcal{P}^\dagger(m,n)\}$ the associated "flow reversed" prop(erad) with $\mathcal{P}^\dagger(m,n) := \mathcal{P}(n,m)$. Let $\mathcal{A}ss$ be the operad of associative algebras and define the properad, $\mathcal{A}ss^\dagger \bullet \mathcal{A}ss = \{\mathcal{A}ss^\dagger \bullet \mathcal{A}ss(m,n)\}$ by setting

$$\mathcal{A}ss^\dagger \bullet \mathcal{A}ss(m,n) := \mathcal{A}ss^\dagger(m) \otimes \mathcal{A}ss(n) \simeq \mathbb{K}[\mathbb{S}_m] \otimes \mathbb{K}[\mathbb{S}_n]$$

and defining the compositions $\mu_{(1,1)}$ to be non-zero only on decorated graphs of the form

(8)  and 

on which it is equal to the operadic compositions in $\mathcal{A}ss$. The properad $\mathcal{A}ss^\dagger \bullet \mathcal{A}ss$ corresponds to the $\frac{1}{2}$-prop $\mathcal{U}^{properad}_{\frac{1}{2}\text{-prop}}(\mathcal{F}rob_+^+)$.

Let $\mathcal{C}om$ be the properad of commutative algebras, and define the properads $\mathcal{C}om^\dagger \bullet \mathcal{A}ss$, $\mathcal{A}ss^\dagger \bullet \mathcal{C}om$, and $\mathcal{C}om^\dagger \bullet \mathcal{C}om$ by analogy to $\mathcal{A}ss^\dagger \bullet \mathcal{A}ss$. Similarly, they correspond to the $\frac{1}{2}$-props $\mathcal{U}^{properad}_{\frac{1}{2}\text{-prop}}(\mathcal{F}rob^+)$, $\mathcal{U}^{properad}_{\frac{1}{2}\text{-prop}}(\mathcal{F}rob_+)$ and $\mathcal{U}^{properad}_{\frac{1}{2}\text{-prop}}(\mathcal{F}rob)$. Applying Theorem 63(i) to $\mathcal{Q}$ being coproperads $(\mathcal{A}ss^\dagger \bullet \mathcal{A}ss)^*$, $(\mathcal{A}ss^\dagger \bullet \mathcal{C}om)^*$, $(\mathcal{C}om^\dagger \bullet \mathcal{A}ss)^*$, or $(\mathcal{C}om^\dagger \bullet \mathcal{C}om)^*$, we conclude that the vector spaces,

$$\bigoplus_{m,n} P(m,n),\ \bigoplus_{m,n} P(m,n)^{\mathbb{S}_m},\ \bigoplus_{m,n} P(m,n)^{\mathbb{S}_n},\ \bigoplus_{m,n} P(m,n)^{\mathbb{S}_m \times \mathbb{S}_n},$$

admit canonically L_∞-structures encoding $\frac{1}{2}$-prop compositions of the form (8). The natural morphism of operads,

$$\mathcal{A}ss \longrightarrow \mathcal{C}om,$$

implies that these L_∞-structures are related to each other via the same commutative diagram of L_∞ morphisms as in Theorem 67(v). In the case of operads the constructed L_∞-structures are exactly the same as in Theorem 67 but for prop(erad)s they are different.

8. Deformation theory of morphisms of prop(erad)s

In this section, we define the deformation theory of morphisms of prop(erad)s. We follow the conceptual method proposed by Quillen in [Qui67, Qui70].

8.1. **Basic Definition.** Let $(\mathcal{P}, d_\mathcal{P}) \xrightarrow{\varphi} (\mathcal{Q}, d_\mathcal{Q})$ be a morphism of dg prop(erad)s. We would like to define a chain complex with which we could study the deformation theory of this map. Following Quillen [Qui70], the conceptual method is to take the total right derived functor of the functor Der of derivations from the category of prop(erad)s above $\mathcal{Q}$ (see also [Mar96a, VdL02]). That is, we consider a cofibrant replacement $(\mathcal{R}, \partial)$ of $\mathcal{P}$ is the category of dg prop(erad)s

$$\begin{tikzcd}
\mathcal{R} \arrow[r, "\varepsilon"] \arrow[dr, "\gamma"'] & \mathcal{P} \arrow[d, "\varphi"] \\
 & \mathcal{Q}.
\end{tikzcd}$$

Recall that $\mathcal{Q}$ has an infinitesimal $\mathcal{P}$-bimodule (respectively infinitesimal $\mathcal{R}$-bimodule) structure given by φ (respectively γ).

Lemma 68. *Let $(\mathcal{R}, \partial)$ be a resolution of $\mathcal{P}$ and let f be a homogenous derivation of degree n in $\mathrm{Der}_n(\mathcal{R}, \mathcal{Q})$, the derivative $D(f) = d_{\mathcal{Q}} \circ f - (-1)^{|f|} f \circ \partial$ is a derivation of degree $n-1$ of $\mathrm{Der}_{n-1}(\mathcal{R}, \mathcal{Q})$.*

Proof. The degree of $D(f)$ is $n-1$. It remains to show that it is a derivation. For every pair r_1 and r_2 of homogenous elements of $\mathcal{R}$, $D(f)\left(\mu_{\mathcal{R}}(r_1 \boxtimes_{(1,1)} r_2)\right)$ is equal to

$$\begin{aligned}
D(f)\left(\mu_{\mathcal{R}}(r_1 \boxtimes_{(1,1)} r_2)\right) &= (d_{\mathcal{Q}} \circ f - (-1)^n f \circ \partial)\left(\mu_{\mathcal{R}}(r_1 \boxtimes_{(1,1)} r_2)\right) \\
&= d_{\mathcal{Q}}\Big(\mu_{\mathcal{Q}}\big(f(r_1) \boxtimes_{(1,1)} \gamma(r_2) + (-1)^{n|r_1|}\gamma(r_1) \boxtimes_{(1,1)} f(r_2)\big)\Big) \\
&\quad -(-1)^n f\Big(\mu_{\mathcal{R}}\big(\partial(r_1) \boxtimes_{(1,1)} r_2 + (-1)^{|r_1|} r_1 \boxtimes_{(1,1)} \partial(r_2)\big)\Big) \\
&= \mu_{\mathcal{Q}}\Big((d_{\mathcal{Q}} \circ f)(r_1) \boxtimes_{(1,1)} \gamma(r_2) + (-1)^{n+|r_1|} f(r_1) \boxtimes_{(1,1)} (d_{\mathcal{Q}} \circ \gamma)(r_2) \\
&\quad +(-1)^{n|r_1|}(d_{\mathcal{Q}} \circ \gamma)(r_1) \boxtimes_{(1,1)} f(r_2) + (-1)^{|r_1|(n-1)}\gamma(r_1) \boxtimes_{(1,1)} (d_{\mathcal{Q}} \circ f)(r_2)\Big) \\
&\quad -(-1)^n \mu_{\mathcal{Q}}\Big((f \circ \partial)(r_1) \boxtimes_{(1,1)} \gamma(r_2) + (-1)^{n(|r_1|-1)}(\gamma \circ \partial)(r_1)) \boxtimes_{(1,1)} f(r_2) \\
&\quad +(-1)^{|r_1|} f(r_1) \boxtimes_{(1,1)} (\gamma \circ \partial)(r_2)) + (-1)^{(n-1)|r_1|}\gamma(r_1) \boxtimes_{(1,1)} (f \circ \partial)(r_2)\Big).
\end{aligned}$$

Since γ is morphism of dg prop(erad)s, it commutes with the differentials, that is $\gamma \circ \partial = d_{\mathcal{Q}} \circ \gamma$. This gives

$$\begin{aligned}
D(f)\left(\mu_{\mathcal{R}}(r_1 \boxtimes_{(1,1)} r_2)\right) &= \mu_{\mathcal{Q}}\Big((d_{\mathcal{Q}} \circ f)(r_1) \boxtimes_{(1,1)} \gamma(r_2) - (-1)^n (f \circ \partial)(r_1) \boxtimes_{(1,1)} \gamma(r_2) \\
&\quad +(-1)^{|r_1|(n-1)}\big(\gamma(r_1) \boxtimes_{(1,1)} (d_{\mathcal{Q}} \circ f)(r_2) - (-1)^n \gamma(r_1) \boxtimes_{(1,1)} (f \circ \partial)(r_2)\big)\Big) \\
&= \mu_{\mathcal{Q}}\Big(D(f)(r_1) \boxtimes_{(1,1)} \gamma(r_2) + (-1)^{|r_1|(n-1)}\gamma(r_1) \boxtimes_{(1,1)} D(f)(r_2)\Big).
\end{aligned}$$

□

In other words, the space of derivations $\mathrm{Der}(\mathcal{R}, \mathcal{Q})$ is a sub-dg-module of the space of morphisms $\mathrm{Hom}^{\mathbb{S}}(\mathcal{R}, \mathcal{Q})$. We define the deformation complex of the morphism φ by $C_\bullet(\varphi) := \big(\mathrm{Der}_\bullet(\mathcal{R}, \mathcal{Q}), D\big)$. By Theorem 98 and Theorem 99, there always exists a quasi-free cofibrant resolutions. For instance, we can consider the bar-cobar resolution by Theorem 19. This will produce an explicit but huge complex which is difficult to compute. Instead of that, we will work with the chain complex obtained from a minimal model of $\mathcal{P}$ when it exists. Its size is much smaller but its differential can be not so easy to make explicit. In this sequel, our main example be the deformation theory of representations of $\mathcal{P}$ of the form $\mathcal{Q} = \mathrm{End}_X$, that is $\mathcal{P}$-gebras.

8.2. **Deformation theory of representations of prop(erad)s.** Let $(\mathcal{P}, d_{\mathcal{P}})$ be a dg prop(erad), let (X, d_X) an arbitrary dg $\mathcal{P}$-gebra and let $(\mathcal{P}_\infty := \Omega(\mathcal{C}), \partial)$ be a cofibrant quasi-free resolution of $\mathcal{P}$ and

$$\begin{tikzcd}
\Omega(\mathcal{C}) \arrow[r, "\varepsilon"] \arrow[dr, "\gamma"'] & \mathcal{P} \arrow[d, "\varphi"] \\
& \mathrm{End}_X .
\end{tikzcd}$$

Definition (Deformation complex)**.** We define the *deformation complex of the $\mathcal{P}$-gebra structure of X* by $C_\bullet(\mathcal{P}, X) := \big(\mathrm{Der}_\bullet(\Omega(\mathcal{C}), \mathrm{End}_X), D\big)$.

Theorem 69. *The deformation complex $\big(\mathrm{Der}_\bullet(\Omega(\mathcal{C}), \mathcal{Q}), D\big)$ is isomorphic to $\mathrm{Hom}^{\mathbb{S}}_\bullet(\bar{\mathcal{C}}, \mathcal{Q})$ with $D = Q^\gamma$ for $\gamma = \varphi \circ \varepsilon_{|\bar{\mathcal{C}}}$.*

Proof. Lemma 14 proves the identification between the two spaces. Since γ is a morphism of dg prop(erad)s from a quasi-free prop(erad), it is a solution of the Maurer-Cartan equation $Q(\gamma) = 0$

in the convolution L_∞-algebra $\mathrm{Hom}^{\mathbb{S}}_{\bullet}(\bar{\mathcal{C}}, \mathcal{Q})$ by Theorem 62. Let f be an element of $\mathrm{Hom}^{\mathbb{S}}_{n}(\bar{\mathcal{C}}, \mathcal{Q})$. Following Lemma 14, we denote by ∂_f the unique derivation of $\mathrm{Der}_n(\Omega(\mathcal{C}), \mathcal{Q})$ induced by f. We have to show that $D(\partial_f)_{s^{-1}\bar{\mathcal{C}}} = Q^\gamma(f)$. For an element $s^{-1}c \in s^{-1}\mathcal{C}$, we use the Sweedler type notation for $\partial(s^{-1}c) = \sum_{\mathcal{G}} \mathcal{G}(s^{-1}c_1, \ldots, s^{-1}c_n)$. By Lemma 14, we have

$$\partial_f\big(\mathcal{G}(s^{-1}c_1, \ldots, s^{-1}c_n)\big) =$$
$$\sum_{i=1}^{n} (-1)^{n(|c_1|+\cdots+|c_{i-1}|+i-1)} \mu_{\mathcal{Q}}\big(\mathcal{G}(\gamma(s^{-1}c_1), \ldots, \gamma(s^{-1}c_{i-1}), f(s^{-1}c_i), \gamma(s^{-1}c_{i+1}), \ldots, \gamma(s^{-1}c_n)\big).$$

Therefore, $D(\partial_f)_{s^{-1}\bar{\mathcal{C}}}$ is equal to

$$D(\partial_f)(s^{-1}c) = (d_{\mathcal{Q}} \circ \partial_f - (-1)^n \partial_f \circ \partial)(s^{-1}c) = d_{\mathcal{Q}}(f(s^{-1}c)) -$$
$$(-1)^n \sum_{\mathcal{G}} \sum_{i=1}^{n} (-1)^{n(|c_1|+\cdots+|c_{i-1}|+i-1)} \mu_{\mathcal{Q}}\big(\mathcal{G}(\gamma(s^{-1}c_1), \ldots, \gamma(s^{-1}c_{i-1}), f(s^{-1}c_i), \gamma(s^{-1}c_{i+1}), \ldots, \gamma(s^{-1}c_n)\big)$$
$$= Q^\gamma(f).$$

□

In order words, the deformation complex is equal to the convolution L_∞-algebra $\mathrm{Hom}^{\mathbb{S}}_{\bullet}(\bar{\mathcal{C}}, \mathcal{Q})$ twisted by the Maurer-Cartan element γ.

Remark. It is natural to consider the augmentation of this chain complex by $\mathrm{Hom}^{\mathbb{S}}(I, \mathcal{Q})$, that is $\mathrm{Hom}^{\mathbb{S}}_{\bullet}(\mathcal{C}, \mathcal{Q})$.

In summary, by Theorem 62 the vector space $\mathrm{Hom}^{\mathbb{S}}_{\bullet}(\mathcal{C}, \mathcal{Q})$ has a canonical filtered L_∞-structure, Q whose Maurer-Cartan elements are morphisms of dg prop(erad)s, $\mathcal{P}_\infty \to \mathcal{Q}$, that is representations of $\mathcal{P}_\infty$ in $\mathcal{Q}$. Then let γ be one of these morphisms, and let Q^γ be the associated twisting of the canonical L_∞-algebra by γ (see §7.3). This defines the deformation complex of γ.

Definition (Deformation Complex). The deformation complex of a morphism of prop(erad)s $\gamma : \mathcal{P}_\infty \to \mathcal{Q}$ is the twisted L_∞-algebra $(\mathrm{Hom}^{\mathbb{S}}_{\bullet}(\mathcal{C}, \mathcal{Q}), Q^\gamma)$.

This definition extends to the case of prop(erad)s the deformation complex of algebras over operads introduced in [KS00, VdL02].

With the results on the model category structure on prop(erad)s (see Appendix), we can now prove the independence of this construction in the homotopy category of homotopy prop(erad)s and homotopy Lie algebras.

Theorem 70. *Let $\Omega(\mathcal{C}_1)$ and $\Omega(\mathcal{C}_2)$ be two quasi-free cofibrant resolutions of a dg prop(erad) $\mathcal{P}$. For any dg prop(erad) $\mathcal{Q}$, the homotopy convolution prop(erad)s $\mathrm{Hom}(\mathcal{C}_1, \mathcal{Q})$ and $\mathrm{Hom}(\mathcal{C}_2, \mathcal{Q})$ are linked by two quasi-isomorphisms of homotopy prop(erad)s*

$$\mathrm{Hom}(\mathcal{C}_1, \mathcal{Q}) \leftrightarrows \mathrm{Hom}(\mathcal{C}_2, \mathcal{Q}),$$

and the natural maps

$$(\mathrm{Hom}^{\mathbb{S}}_{\bullet}(\mathcal{C}_1, \mathcal{Q}), Q^{\gamma_1}) \leftrightarrows (\mathrm{Hom}^{\mathbb{S}}_{\bullet}(\mathcal{C}_2, \mathcal{Q}), Q^{\gamma_2})$$

is a quasi-isomorphism of homotopy Lie algebras.

Proof. We apply the left lifting property in the model category of dg prop(erad)s to the following diagram :

$$\begin{tikzcd}
0 \arrow[r] \arrow[d, tail] & \Omega(\mathcal{C}_1) \arrow[d, two heads, "\sim"] \arrow[dl, dotted, shift left] \\
\Omega(\mathcal{C}_2) \arrow[r, two heads, "\sim"] \arrow[ur, dotted, shift left] & \mathcal{P},
\end{tikzcd}$$

to get the two dotted quasi-isomorphisms of prop(erad)s. By Proposition 100, they induce quasi-isomorphisms of the level of the homotopy coprop(erad)s $\mathcal{C}_1 \leftrightarrows \mathcal{C}_2$. We conclude by Theorem 33 and by Corollary 35. □

The homology groups,

$$H^{\gamma}_{\bullet}(\mathcal{Q}) := H_{\bullet}(\mathrm{Hom}^{\mathbb{S}}(\mathcal{C}, \mathcal{Q}), Q^{\gamma}_{1}),$$

are independent of the choice of a cofibrant quasi-free resolution of $\mathcal{P}$ and are called *homology groups of the $\mathcal{P}_\infty$-representation $\mathcal{Q}$.* In the case where $\mathcal{Q} = \mathrm{End}_X$, they are called the homology groups of the $\mathcal{P}_\infty$-gebra (X, γ).

Proposition 71. *The Maurer-Cartan elements, Γ, of Q^γ are in one-to-one correspondence with those $\mathcal{P}_\infty$-structures, on X,*

$$\rho : (\mathcal{P}_\infty, \partial) \longrightarrow (\mathrm{End}_X, d),$$

whose restrictions to the generating space, $s^{-1}\mathcal{C}$, of $\mathcal{P}_\infty$ are equal precisely to the sum $\gamma + \Gamma$.

This proposition justifies the name 'deformation complex' because the L_∞-algebra Q^γ *controls the deformations of γ in the class of homotopy $\mathcal{P}$-structures.* When applied to $\mathcal{Q} = \mathrm{End}_X$ and $\gamma : \mathcal{P}_\infty \to \mathrm{End}_X$, this defines the deformation complex of the $\mathcal{P}_\infty$-gebra structure γ on X. (Some author call this the "cohomology of X with coefficients into itself" but we are reluctant to make this choice and prefer to view it as a deformation complex). This definition applies to any homotopy algebra over an operad (associative algebras, Lie algebras, commutative algebras, PreLie algebras, Poisson or Gerstenhaber algebras, etc ...) as well as to any homotopy (bial)gebra over a properad (Lie bialgebras, associative bialgebras, etc ...) in order to give, for the first time, a cohomology theory for homotopy $\mathcal{P}$-(bial)gebras.

8.3. **Koszul case and cohomology operations.** In Theorem 39, we have seen that a properad $\mathcal{P}$ is Koszul if and only if it admits a quadratic model $\Omega(\mathcal{P}^{\text{¡}}) \xrightarrow{\sim} \mathcal{P}$, where $\mathcal{P}^{\text{¡}}$ the Koszul dual (strict) coproperad. In this case, by Theorem 69, the deformation complex of a $\mathcal{P}_\infty$-gebra $\mathrm{Hom}^{\mathbb{S}}(\mathcal{P}^{\text{¡}}, \mathrm{End}_X)$ is dg Lie algebra where the boundary map is equal to the twisted differential $D(f) = d(f) + [\gamma, f]$.

The first definition of this kind of preLie operation appeared in the seminal paper of M. Gerstenhaber [Ger63] in the case of the cohomology of associative algebras. In the case treated by M. Gerstenhaber, the cooperad $\mathcal{C}$ is the Koszul dual cooperad $\mathcal{A}s^{\text{¡}}$ of the operad $\mathcal{A}s$ coding associative algebras and the operad $\mathcal{P}$ is the endomorphism operad End_A. The induced Lie bracket is the *intrinsic Lie bracket* of Stasheff [Sta93]. It is equal to the Lie bracket of Gerstenhaber [Ger63] on Hochschild cochain complex of associative algebras, the Lie bracket of Nijenhuis-Richardson [NR67] on Chevalley-Eilenberg cochain complex of Lie algebras and the Lie bracket of Stasheff on Harrison cochain complex of commutative algebras. It is proven by Balavoine in [Bal97] that the deformation complex of algebras over any Koszul operad admits a Lie structure. This statement was made more precise by Markl, Shnider and Stasheff in Section 3.9 Part II of [MSS02] where they proves that this Lie bracket comes from a Prelie product. This result on the level of operads was proved using the space of coderivations of the cofree $\mathcal{P}^{\text{¡}}$-coalgebra, which is shown to be a PreLie algebra. Such a method is impossible to generalize to prop(erad)s simply because there exists no notion of (co)free gebra.

As explained here, one has to work with convolution prop(erad) to prove a similar result. Actually, this method gives a stronger statement.

Theorem 72. *Let $\mathcal{P}$ be a Koszul properad and let $\varphi : \mathcal{P} \to \mathcal{Q}$ be a morphism of properads, the deformation complex of φ is a LR-algebra.*
In the non-symmetric case, when $\mathcal{P}$ and $\mathcal{Q}$ are non-symmetric properads, the deformation complex is a non-symmetric properad.

Proof. It is direct consequence of the definition of the deformation complex and Theorem 13. In the non-symmetric case, the deformation complex is directly a non-symmetric convolution properad, since it is not restricted to invariant elements. □

This result provides higher braces or LR-operations (see Section 2.4). Recall that non-symmetric braces play a fundamental role in the proof of Deligne's conjecture for associative algebras (see [Tam98, Vor00, KS00, MS02, BF04]) and in the extension of it to other kind of algebras (see

[Val06b] Section 5.5). From this rich structure, we derive a Lie-admissible bracket and then a Lie bracket which can be used to study the deformations of X. We expect the LR-operations to be used in the future for a better understanding of deformation theory (in the context of a Deligne conjecture for associative bialgebras and Gerstenhaber-Schack bicomplex, for instance).

Notice that this Lie bracket was found by hand in one example of gebras over a properad before this general theory. The properad of Lie bialgebras is Koszul. Therefore, on the deformation (bi)complex of Lie bialgebras, there is a Lie bracket. The construction of this Lie bracket was given by Kosmann-Schwarzbach in [KS91]. (See also Ciccoli-Guerra [CG03] for the interpretation of this bicomplex in terms of deformations.)

8.4. **Definition à la Quillen.** In the previous sections, we defined the deformation complex of representations of a prop(erad) $\mathcal{P}$ that admits a quasi-free model and proved the independence of this definition in the categories of homotopy prop(erad)s (and homotopy Lie algebras). In this section, we generalize the definition of the deformation theory of a morphism of commutative rings due to Quillen [Qui70] to the case of prop(erad)s. (See L. Illusie [Ill71] for a generalization in the context of topoi and schemes). Hence, it defines a (relative) deformation complex for representations of any prop(erad). It also gives rise to the *cotangent complex* associated to any morphism of prop(erad)s.
Since a commutative algebra is an associative algebra, an associative algebra an operad and an operad a prop(erad), this generalization of Quillen theory can be seen as a way to extend results of (commutative) algebraic geometry to non-commutative non-linear geometry. It is non-linear because the monoidal product $\boxtimes$ defining prop(erad)s is neither linear on the left nor on the right, contrary to the tensor product $\otimes$ of vector spaces.

Let $\mathcal{I}$ be a 'ground' prop(erad) (to recover the previous section, consider $\mathcal{I} = I$, the unit of the monoidal category of $\mathbb{S}$-bimodules). We look at prop(erad)s $\mathcal{P}$ under $\mathcal{I}$, $\mathcal{I} \to \mathcal{P}$. And for such a prop(erad) $\mathcal{P}$, we consider the category of prop(erad)s over $\mathcal{P}$, that is

$$\begin{tikzcd}
& \mathcal{X} \arrow[d, "f"] \\
\mathcal{I} \arrow[ru, "u"] \arrow[r] & \mathcal{P}.
\end{tikzcd}$$

We denote this category by Prop(erad)/$\mathcal{P}$. Let M be an infinitesimal bimodule over $\mathcal{P}$ (see Section 3.1). The infinitesimal $\mathcal{P}$-bimodule M is also an infinitesimal bimodule over any prop(erad) $\mathcal{X}$ over $\mathcal{P}$, by pulling back along $\mathcal{X} \to \mathcal{P}$. Hence, we can consider the space of $\mathcal{I}$-derivations from $\mathcal{X}$ to M, that is derivations from $\mathcal{X}$ to M which vanish on $\mathcal{I}$. We denoted this space by $\mathrm{Der}_{\mathcal{I}}(\mathcal{X}, M)$. We aim now to represent this bifunctor on the left and on the right. To represent it on the left, we introduce the *square-zero (or infinitesimal) extension of* $\mathcal{P}$ *by* M : $\mathcal{P} \ltimes M := \mathcal{P} \oplus M$ with the following structure of prop(erad) over $\mathcal{P}$. The monoidal product $(\mathcal{P} \oplus M) \boxtimes (\mathcal{P} \oplus M)$ is equal to

$$\mathcal{P} \boxtimes \mathcal{P} \bigoplus \mathcal{P} \boxtimes (\mathcal{P} \oplus \underbrace{M}_{1}) \bigoplus (\mathcal{P} \oplus \underbrace{M}_{1}) \boxtimes \mathcal{P} \bigoplus \mathcal{M},$$

where $\mathcal{M}$ is the sub-$\mathbb{S}$-bimodule of $(\mathcal{P} \oplus M) \boxtimes (\mathcal{P} \oplus M)$ composed by at least two elements from M. On the first component $\mathcal{P} \boxtimes \mathcal{P}$, the product of $\mathcal{P} \ltimes M$ is defined by the product of $\mathcal{P}$. On the second component, it is defined by the left action of $\mathcal{P}$ on M. On the third one, it is defined by the right action of $\mathcal{P}$ on M. Finally, the product on $\mathcal{M}$ is null.

Lemma 73. *For any prop(erad)* $\mathcal{P}$ *and any infinitesimal* $\mathcal{P}$*-bimodule* M*, the infinitesimal extension* $\mathcal{P} \ltimes M$ *is a prop(erad).*

PROOF. The definition of infinitesimal $\mathcal{P}$-bimodule directly implies the associativity of the prop(erad)ic composition of $\mathcal{P} \ltimes M$. □

The purpose of this definition is in the following result, which states that infinitesimal $\mathcal{P}$-bimodules are abelian group objects in the category of prop(erad)s over $\mathcal{P}$.

Proposition 74. *There is a natural bijection*

$$\mathrm{Hom}_{Prop(erad)/\mathcal{P}}(\mathcal{X}, \mathcal{P} \ltimes M) \cong \mathrm{Der}_{\mathcal{I}}(\mathcal{X}, M),$$

where $\mathrm{Der}_{\mathcal{I}}(\mathcal{X}, M)$ *is the space of* $\mathcal{I}$*- derivations from* $\mathcal{X}$ *to* M*.*

Proof. Let us denote by $f : \mathcal{X} \to \mathcal{P}$. Any morphism $\mathcal{X} \to \mathcal{P} \ltimes M = \mathcal{P} \oplus M$ the category of prop(erad)s over $\mathcal{P}$ is the sum of f with its component on M, which we denote by D. Finally, $f \oplus D : \mathcal{X} \to \mathcal{P} \ltimes M$ is a morphism of prop(erad)s if and only if D is a derivation $\mathcal{X} \to M$. ☐

To represent the space of derivations on the right, we introduce the *module of Kähler differentials of a prop(erad).* It is a quotient of the free infinitesimal $\mathcal{X}$-bimodule over $\mathcal{I}$ on $\mathcal{X}$ by suitable relations. We recall from Section 2.5 of [Val07a] that the relative composition product is defined by the following coequalizer

$$\begin{tikzcd} M \boxtimes \mathcal{P} \boxtimes N \arrow[r, shift left, "M\boxtimes\lambda"] \arrow[r, shift right, "\rho\boxtimes N"'] & M \boxtimes N \arrow[r, two heads] & M \boxtimes_{\mathcal{P}} N \end{tikzcd} ,$$

where λ is the left action of $\mathcal{P}$ on N, $\mathcal{P} \boxtimes N \to N$ and ρ the right action of $\mathcal{P}$ on M, $M \boxtimes \mathcal{P} \to M$.

Let $f : \mathcal{P} \to \mathcal{Q}$ be a morphism of prop(erad)s. There is a natural functor from the category of infinitesimal $\mathcal{Q}$-bimodules to the category of infinitesimal $\mathcal{P}$-bimodules by pulling back along f. We denote it by $f^* : \mathrm{Inf.}\, \mathcal{Q}\text{-biMod} \to \mathrm{Inf.}\, \mathcal{P}\text{-biMod}$.

Proposition 75. *The functor* $f^* : \mathit{Inf.}\, \mathcal{Q}\text{-}\mathit{biMod} \to \mathit{Inf.}\, \mathcal{P}\text{-}\mathit{biMod}$ *admits a left adjoint*

$$f_! \ : \ \mathit{Inf.}\, \mathcal{P}\text{-}\mathit{biMod} \leftrightharpoons \mathit{Inf.}\, \mathcal{Q}\text{-}\mathit{biMod} \ : \ f^*,$$

which is explicitly given by $f_!(M) = \mathcal{Q} \boxtimes_{\mathcal{P}} \underbrace{M}_{1} \boxtimes_{\mathcal{P}} \mathcal{Q}$*, for any infinitesimal* $\mathcal{P}$*-bimodule* M*. The* $\mathbb{S}$*-bimodule* $\mathcal{Q} \boxtimes_{\mathcal{P}} \underbrace{M}_{1} \boxtimes_{\mathcal{P}} \mathcal{Q}$ *is the coequalizer*

$$\begin{tikzcd} (\mathcal{Q} \boxtimes \mathcal{P}) \boxtimes \underbrace{M}_{1} \boxtimes (\mathcal{P} \boxtimes \mathcal{Q}) \arrow[r, shift left, "\rho_{\mathcal{Q}}\boxtimes M\boxtimes\lambda_{\mathcal{Q}}"] \arrow[r, shift right, "\mathcal{Q}\boxtimes(\lambda\circ(\mathcal{P}\boxtimes\rho))\boxtimes\mathcal{Q}"'] & \mathcal{Q} \boxtimes \underbrace{M}_{1} \boxtimes \mathcal{Q} \arrow[r, two heads] & \mathcal{Q} \boxtimes_{\mathcal{P}} \underbrace{M}_{1} \boxtimes_{\mathcal{P}} \mathcal{Q}, \end{tikzcd}$$

where the notation $\mathcal{Q} \boxtimes \underbrace{M}_{1} \boxtimes \mathcal{Q}$ *stands for 3-levels graphs with only one element of* M *labelling a vertex on the second level and such that every element of* Q *on the first and third level have a common internal edge with this element of* M*. (The action of* $\mathcal{P}$ *on* $\mathcal{Q}$ *is given by the morphism* f*.)*

Proof. We have the natural bijection

$$\mathrm{Hom}_{\mathrm{Inf.}\, \mathcal{Q}\text{-biMod}}(\mathcal{Q} \boxtimes_{\mathcal{P}} \underbrace{M}_{1} \boxtimes_{\mathcal{P}} \mathcal{Q}, N) \cong \mathrm{Hom}_{\mathrm{Inf.}\, \mathcal{P}\text{-biMod}}(M, f^*(N)).$$

Let $\Phi : \mathcal{Q} \boxtimes_{\mathcal{P}} \underbrace{M}_{1} \boxtimes_{\mathcal{P}} \mathcal{Q} \to N$ be a morphism a infinitesimal $\mathcal{Q}$-bimodules. It is characterized by the image of the projection of the element of $I \boxtimes \underbrace{M}_{1} \boxtimes I$ in $\mathcal{Q} \boxtimes_{\mathcal{P}} \underbrace{M}_{1} \boxtimes_{\mathcal{P}} \mathcal{Q}$. Let us call $\varphi : M \to N$ this map. It is then easy to see that φ is a morphism of infinitesimal $\mathcal{P}$-bimodules. ☐

Example. If we apply the preceding proposition to the unit $u : I \to \mathcal{P}$ of a prop(erad) $\mathcal{P}$, the functor $u^* : \mathrm{Inf.}\, \mathcal{P}\text{-biMod} \to \mathbb{S}\text{-biMod}$ is the classical forgetful functor. Hence $u_!(M) = \mathcal{P} \boxtimes \underbrace{M}_{1} \boxtimes \mathcal{P}$ is the free infinitesimal $\mathcal{P}$-bimodule associated to any $\mathbb{S}$-bimodule M.

Definition (Module of Kähler differentials)**.** Let us denote by $u : \mathcal{I} \to \mathcal{X}$. The module of Kähler differentials of a prop(erad) $\mathcal{X}$ over $\mathcal{I}$ is the quotient of the free infinitesimal $\mathcal{X}$-bimodule over the infinitesimal $\mathcal{I}$-bimodule $u^*(\mathcal{X})$, that is

$$u_!(u^*(\mathcal{X})) = \mathcal{X} \boxtimes_{\mathcal{I}} \underbrace{u^*(\mathcal{X})}_{1} \boxtimes_{\mathcal{I}} \mathcal{X},$$

by the relations

$$\pi\big(I \boxtimes \mu_{\mathcal{X}}(x_1 \boxtimes_{(1,1)} x_2) \boxtimes I - I \boxtimes x_1 \boxtimes_{(1,1)} x_2 - (-1)^{|x_1|} x_1 \boxtimes_{(1,1)} x_2 \boxtimes I\big),$$

where π is the canonical projection of $\mathcal{X} \boxtimes \underbrace{\mathcal{X}}_{1} \boxtimes \mathcal{X}$ on the coequalizer $\mathcal{X} \boxtimes_{\mathcal{I}} \underbrace{u^*(\mathcal{X})}_{1} \boxtimes_{\mathcal{I}} \mathcal{X}$. We denote it by $\Omega_{\mathcal{X}/\mathcal{I}}$.

We define the universal derivation $D : \mathcal{X} \to \Omega_{\mathcal{X}/\mathcal{I}}$ by $D(x)$ equal to the class of $I \boxtimes x \boxtimes I$ in $\Omega_{\mathcal{X}/\mathcal{I}}$. Like in the case of commutative algebras (see J.-L. Loday [Lod98] Section 1.3) or associative algebras (see [Con85, Kar87, Lod98], the module of Kähler differentials represents the derivations.

Proposition 76. *There is a natural bijection*

$$\mathrm{Der}_{\mathcal{I}}(\mathcal{X}, M) \cong \mathrm{Hom}_{\textit{Inf. } \mathcal{X}\textit{-biMod}}(\Omega_{\mathcal{X}/\mathcal{I}}, M).$$

Proof. Let d be a derivation in $\mathrm{Der}_{\mathcal{I}}(\mathcal{X}, M)$. There is a unique morphism of infinitesimal $\mathcal{X}$-bimodules $\theta : \Omega_{\mathcal{X}/\mathcal{I}} \to M$ such that the following diagram commutes

$$\begin{tikzcd}
\mathcal{X} \arrow[r, "D"] \arrow[dr, "d"'] & \Omega_{\mathcal{X}/\mathcal{I}} \arrow[d, dashed, "\theta"] \\
 & M.
\end{tikzcd}$$

The image of the class of $I \boxtimes x \boxtimes I$ in $\Omega_{\mathcal{X}/\mathcal{I}}$ under θ is defined by $d(x)$. It extends freely to the infinitesimal $\mathcal{X}$-bimodule $\mathcal{X} \boxtimes_{\mathcal{I}} \underbrace{u^*(\mathcal{X})}_{1} \boxtimes_{\mathcal{I}} \mathcal{X}$ and then passes to the quotient thanks to the Leibniz relation verified by d. ☐

The module of Kähler differentials of an associative algebra is the non-commutative analog of classical differential forms (see A. Connes [Con85]). Since operads and prop(erad)s can also be used to encode geometry (see [Mer05, Mer06]), the module of Kähler differentials for prop(erad)s seems a promising tool to study non-linear properties in non-commutative geometry.

Theorem 77. *For any infinitesimal $\mathcal{P}$-bimodule M, the following adjunction holds*

$$\mathrm{Hom}_{\textit{Prop(erad)}/\mathcal{P}}(\mathcal{X}, \mathcal{P} \ltimes M) \cong \mathrm{Der}_{\mathcal{I}}(\mathcal{X}, M) \cong \mathrm{Hom}_{\textit{Inf. } \mathcal{P}\textit{-biMod}}(\mathcal{P} \boxtimes_{\mathcal{X}} \underbrace{\Omega_{\mathcal{X}/\mathcal{I}}}_{1} \boxtimes_{\mathcal{X}} \mathcal{P}, M).$$

Proof. It is a direct corollary of Proposition 74 and Proposition 76. The last natural bijection

$$\mathrm{Hom}_{\mathrm{Inf.}\ \mathcal{X}\text{-biMod}}(\Omega_{\mathcal{X}/\mathcal{I}}, f^*(M)) \cong \mathrm{Hom}_{\mathrm{Inf.}\ \mathcal{P}\text{-biMod}}(f_!(\Omega_{\mathcal{X}/\mathcal{I}}), N)$$

is provided by Proposition 75 applied to the morphism $f : \mathcal{X} \to \mathcal{P}$. ☐

In other words, the following functors form a pair of adjoint functors

$$\mathcal{P} \boxtimes_{-} \underbrace{\Omega_{-/\mathcal{I}}}_{1} \boxtimes_{-} \mathcal{P} \ : \ \mathrm{Prop(erad)}/\mathcal{P} \ \rightleftharpoons \ \mathrm{Inf.}\, \mathcal{P}\text{-biMod} \ : \mathcal{P} \ltimes -.$$

The model category structure on prop(erad)s induces a model category structure on Prop(erad)/$\mathcal{P}$.

Lemma 78. *The category of infinitesimal $\mathcal{P}$-bimodules is endowed with a cofibrantly generated model category structure.*

Proof. We use the same arguments as in Appendix A, that is the Transfer Theorem 89 along the free infinitesimal $\mathcal{P}$-bimodule functor $\eta_!$: $\mathbb{S}$-biMod $\to$ Inf.$\mathcal{P}$-biMod. The forgetful functor η^* creates limits and colimits which proves (1) and (2). A relative $\eta_!(J)$-cell complex has the form $A_0 \to A_0 \oplus \mathcal{P} \boxtimes \underbrace{(\oplus_{i\geq 0} D^{k_i}_{m_i,n_i})}_{1} \boxtimes \mathcal{P}$, which is a quasi-isomorphism of dg $\mathbb{S}$-bimodules since the right hand term $\mathcal{P} \boxtimes \underbrace{(\oplus_{i\geq 0} D^{k_i}_{m_i,n_i})}_{1} \boxtimes \mathcal{P}$ is acyclic. $\square$

Proposition 79. *The pair of adjoint functors*

$$\mathcal{P} \boxtimes_- \underbrace{\Omega_{-/\mathcal{I}}}_{1} \boxtimes_- \mathcal{P} \ : \ \mathit{Prop(erad)}/\mathcal{P} \ \rightleftharpoons \ \mathit{Inf.}\,\mathcal{P}\text{-}\mathit{biMod} \ : \mathcal{P} \ltimes -$$

form a Quillen adjunction.

Proof. By Lemma 1.3.4 of [Hov99], it is enough to prove that the right adjoint $\mathcal{P} \ltimes -$ preserves fibrations and acyclic fibrations. Let $f : M \twoheadrightarrow M'$ be a fibration (resp. acyclic fibration) between two infinitesimal $\mathcal{P}$-bimodules, that is f is degreewise surjective (resp. and a quasi-isomorphism). Since $\mathcal{P} \ltimes (f)$ is the morphism of properads on $Id_{\mathcal{P}} \oplus f : \mathcal{P} \oplus M \to \mathcal{P} \oplus M'$, it is degreewise surjective (resp. and a quasi-isomorphism), which concludes the proof. $\square$

Thereofre, we can derive them in the associated homotopy categories.

This proves that the homology of $\mathrm{Der}_{\mathcal{I}}(\mathcal{R}, M)$ is independent of the choice of the cofibrant resolution of $\mathcal{P}$ because it is well defined in the homotopy category of prop(erad)s over $\mathcal{P}$ and in the homotopy category infinitesimal $\mathcal{P}$-bimodules.

$$\mathrm{Hom}_{\mathsf{Ho}(\mathrm{Prop(erad)}/\mathcal{P})}(\mathcal{X}, \mathcal{P} \ltimes M) \cong \mathrm{Der}_{\mathcal{I}}(\mathcal{X}, M) \cong \mathrm{Hom}_{\mathsf{Ho}(\mathrm{Inf.}\,\mathcal{P}\text{-biMod})}(\mathcal{P} \boxtimes_{\mathcal{X}} \underbrace{\Omega_{\mathcal{X}/\mathcal{I}}}_{1} \boxtimes_{\mathcal{X}} \mathcal{P}, M).$$

Definition (Cotangent complex)**.** The *cotangent complex* of $\mathcal{P}$ is the total left derived functor of the right adjoint, that is

$$\mathbb{L}_{\mathcal{P}/\mathcal{I}} := \mathcal{P} \boxtimes_{\mathcal{R}} \underbrace{\Omega_{\mathcal{R}/\mathcal{I}}}_{1} \boxtimes_{\mathcal{R}} \mathcal{P},$$

for $\mathcal{R}$ a cofibrant resolution of $\mathcal{P}$.

Since on the homology of the cotangent complex, in the classical case of commutative rings, one can read the properties of the morphism $\mathcal{I} \to \mathcal{P}$ (smooth, locally complete intersection,etc ...), we expect to be able to read such properties on the generalized version defined here. In the same way, transitivity and flat base change theorems should be proved for this cotangent complex but it is not our aim here and will be studied in a future work.

Remark. This section is written in the category of dg-prop(erad)s since we work in this paper over a field of characteristic 0. Therefore, to explicit the cotangent complex and the (co)homology of prop(erad)s, we have to use cofibrant resolutions in the category of dg prop(erad)s, for instance quasi-free resolutions (Koszul or homotopy Koszul). One can extend this section and the Appendix when the characteristic of the ground ring is not 0. In this case, one has to use simplicial resolutions like in M. André [And74] and D. Quillen [Qui70].

9. Examples of deformation theories.

In this section, we show that the conceptual deformation theory defined here coincide to well known theories in the case of associative algebras, Lie algebras, commutative algebras, Poisson algebras. As a corollary, we get classical Lie brackets on these cohomology theory as well as classical Lie brackets in differential geometry. More surprisingly, we make deformation theory explicit in the case of associative bialgebras and show that it corresponds to Gerstenhaber-Schack type bicomplex.

9.1. **Associative algebras.** If P is the properad, Ass, of associative algebras, it is generated by a non-symmetric operad still denoted by Ass. This operad is Koszul that is, its minimal resolution exists and is generated by the (strict) cooperad $\mathsf{Ass}^{\mathsf{i}}$ with

$$\mathsf{Ass}^{\mathsf{i}}(m,n) = \begin{cases} s^{n-2}\mathbb{K}[\mathbb{S}_1]\otimes\mathbb{K}[\mathbb{S}_n] & \text{for } m=1, n\geq 2 \\ 0 & \text{otherwise.}\end{cases}$$

We represent the generating element of $\mathsf{Ass}(1,n)$ by a corolla . The partial coproduct of this cooperad is given by the formula



$$\Delta_{(1,1)}\left(\ \right) = \sum_{k=0}^{n-1}\sum_{l=1}^{n-k}(-1)^{(l-1)(n-k-l)} \quad .$$

Proposition 80. *Let* $\mathsf{Ass}\xrightarrow{\varphi}\mathcal{Q}$ *be a map of (non-symmetric) operads. The deformation complex of this map is isomorphism to* $\mathcal{Q}$ *up to the following shift of degree*

$$C^{\varphi}_{\bullet}(\mathsf{Ass},\mathcal{Q})(n) = s^{-1}\mathrm{Hom}^{\mathbb{S}}_{\bullet}(\mathsf{Ass}^{i},\mathcal{Q})(n) = s^{-n}\mathcal{Q}_{\bullet}(n).$$

The boundary map is given by

$$D(q) = d(q) + \mu_{\mathcal{Q}}(\varphi(\nu); I, q) + \sum_{i=1}^{n}(-1)^i\mu_{\mathcal{Q}}(q; I,\ldots,I,\underbrace{\varphi(\nu)}_{i},I,\ldots,I) + (-1)^{n+1}\mu_{\mathcal{Q}}(\varphi(\nu); q, I),$$

for $q\in\mathcal{Q}(n)$ *if we denote by* ν *the generating binary operation of* $\mathsf{Ass}(2)$.

Proof. There is a one-to-one correspondence between $\mathbb{S}_n$-equivariant maps from $\mathsf{Ass}^{\mathsf{i}}(n)$ to $\mathcal{Q}(n)$ and elements of $\mathcal{Q}(n)$. Let us denote by f_q the unique map determined by $q\in\mathcal{Q}(n)$. Since $\mathsf{Ass}^{\mathsf{i}}$ is a coooperad and $\mathcal{Q}$ is an operad, the convolution operad $\mathrm{Hom}(\mathsf{Ass}^{\mathsf{i}},\mathcal{Q})$ is preLie algebra with product denote $\star$ (see Section 2). By Theorem 69 and Section 8.3, we have $D(f_q) = d(f_q) + \gamma\star f_q - (-1)^{|f_q|}f_q\star\gamma$. Since f_q vanishes on $\mathsf{Ass}^{\mathsf{i}}(m)$ for $m\neq n$ and since γ vanishes on $\mathsf{Ass}^{\mathsf{i}}(m)$ for $m\neq 2$, the only non-vanishing component of $\gamma\star f_q = \mu_{\mathcal{Q}}\circ(\gamma\boxtimes_{(1,1)}f_q)\circ\Delta_{(1,1)}$ is $-\mu_{\mathcal{Q}}(\varphi(\nu); I, q) + (-1)^n\mu_{\mathcal{Q}}(\varphi(\nu); q, I)$ on $\mathrm{Hom}^{\mathbb{S}}_{\bullet}(\mathsf{Ass}^{\mathsf{i}},\mathcal{Q})$. And the only non-vanishing component of $f_q\star\gamma = \mu_{\mathcal{Q}}\circ(f_q\boxtimes_{(1,1)}\gamma)\circ\Delta_{(1,1)}$ is $\sum_{i=1}^{n}(-1)^{i+1}\mu_{\mathcal{Q}}(q; I,\ldots,I,\underbrace{\varphi(\nu)}_{i},I,\ldots,I)$ on $\mathrm{Hom}^{\mathbb{S}}_{\bullet}(\mathsf{Ass}^{\mathsf{i}},\mathcal{Q})$, which concludes the proof. □

This deformation complex appears in many places in the literature under different names. When $\mathcal{Q} = \mathrm{End}_X$ with X an associative algebra, it is the Hochschild (co)chain complex of X (*with coefficient in* X): $s^{-1}\mathrm{Hom}(V,\mathrm{End}_X) = \oplus_{n\geq 2}s^{1-n}\mathrm{Hom}(X^{\otimes n}, X)$. The induced L_∞-algebra, Q on it is strict since the operad Ass is Koszul. It is precisely the Gerstenhaber Lie algebra [Ger63] and Q^γ is the Hochschild dg Lie algebra controlling deformations of a particular associative algebra structure, $\gamma : \mathsf{Ass}\to\mathrm{End}_X$, on a vector space X.

In the work of McClure-Smith on Deligne's conjecture [MS02], an operad $\mathcal{Q}$ with a morphism of operads $\mathsf{Ass}\to\mathcal{Q}$ is called a *multiplicative operad.* The simplicial complex that they define on such an operad is exactly the deformation complex of this map. For the operad $\mathcal{Q} = \mathit{Poisson}$, this complex is related to the homology of long knots (see [Tou04]). More generally, Maxim Kontsevich proposed the conjecture that the deformation complex of $\mathsf{Ass}\to\mathrm{End}_X$ is a $d+1$-algebra when X is a d-algebra in [Kon99]. This conjecture was proved by Tamarkin in [Tam00], see also Hu, Kriz and Voronov [HKV06]. In this context, this chain complex is often called the *Hochschild complex of* $\mathcal{Q}$.

Since this (co)chain complex comes from the general theory of (co)homology of Quillen, it would be better to call its (co)homology the *cohomology of* Ass *with coefficients in* $\mathcal{Q}$ or the chain complex, the *deformation complex of the map* φ.

Analogously one recovers other classical examples — Harrison complex/cohomology and Chevalley-Eilenberg complex/cohomology — from the operads of commutative algebras and, respectively, Lie algebras.

9.2. **Poisson structures.** A *Lie 1-bialgebra* is, by definition, a graded vector space V together with two linear maps,

$$\begin{array}{rccc} \delta: & V & \longrightarrow & \wedge^2 V \\ & a & \longrightarrow & \sum a_1\wedge a_2 \end{array} \quad , \quad \begin{array}{rccc} [\bullet]: & \odot^2 V & \longrightarrow & V \\ & a\otimes b & \longrightarrow & (-1)^{|a|}[a\bullet b] \end{array}$$

of degrees 0 and -1 respectively which satisfy the identities,

(i) $(\delta\otimes \mathrm{Id})\delta a + \tau(\delta\otimes\mathrm{Id})\delta a + \tau^2(\delta\otimes\mathrm{Id})\delta a = 0$, where τ is the cyclic permutation (123) represented naturally on $V\otimes V\otimes V$ (co-Jacobi identity);
(ii) $[[a\bullet b]\bullet c] = [a\bullet[b\bullet c]] + (-1)^{|b||a|+|b|+|a|}[b\bullet[a\bullet c]]$ (Jacobi identity);
(iii) $\delta[a\bullet b] = \sum a_1\wedge[a_2\bullet b] - (-1)^{|a_1||a_2|}a_2\wedge[a_1\bullet b] + [a\bullet b_1]\wedge b_2 - (-1)^{|b_1||b_2|}[a\bullet b_2]\wedge b_1$ (Leibniz type identity).

This notion of Lie 1-bialgebras is similar to the well-known notion of Lie bialgebras except that in the latter case both operations, Lie and co-Lie brackets, have degree 0.

Let $\mathcal{L}ieBi$ be the properad whose representations are Lie 1-bialgebras. It is Koszul contractible, that is its minimal resolution, $(\mathcal{L}ieBi_\infty, \delta)$, exists and is generated by the $\mathbb{S}$-bimodule $V = \{V(m,n)\}_{m,n\geq 1, m+n\geq 3}$ with

$$V(m,n) := s^{m-2} sgn_m \otimes \mathbf{1}_n = \mathrm{span}$$



,

where $sign_m$ stands for the sign representation of $\mathbb{S}_m$ and $\mathbf{1}_n$ for the trivial representation of $\mathbb{S}_n$. The differential is given on generators by [Mer06]

$$\delta$$



$$= \sum_{\substack{I_1\sqcup I_2 = (1,\dots,m)\\ J_1\sqcup J_2=(1,\dots,n)\\ |I_1|\geq 0, |I_2|\geq 1\\ |J_1|\geq 1, |J_2|\geq 0}} (-1)^{\sigma(I_1\sqcup I_2)+|I_1||I_2|}$$



where $\sigma(I_1\sqcup I_2)$ is the sign of the shuffle $I_1\sqcup I_2 = (1,\dots,m)$.

Hence, for an arbitrary dg vector space X,

$$s^{-1}\mathrm{Hom}(V,\mathrm{End}_X) = \bigoplus_{m,n\geq 1} s^{1-m}\wedge^m X\otimes \odot^n X \simeq \wedge^\bullet T_X,$$

where $\wedge^\bullet T_X$ is the vector space of formal germs of polyvector fields at $0\in X$ when we view X as a formal graded manifold. It is not hard to show using the above explicit formula for the differential δ that the canonically induced, in accordance with Theorem 62(i), L_∞-structure on $s^{-1}\mathrm{Hom}(V,\mathrm{End}_X)$ is precisely the classical Schouten Lie algebra structure on polyvector fields. Thus our theory applied to Lie 1-bialgebras reproduces deformation theory of Poisson structures, and $\mathcal{L}ieBi$-homology is precisely Poisson homology.

In a similar way one can check that our construction of L_∞-algebras applied to the minimal resolution of so called pre-Lie^2-algebras [Mer05] gives rise to another classical geometric object — the Frölicher-Nijenhuis Lie brackets on the sheaf, $T_X\otimes\Omega_X^\bullet$, of tangent vector bundle valued differential forms. Thus the associated deformation theory describes deformations of integrable Nijenhuis structures.

9.3. **Associative bialgebras.** In this section, we make explicit the deformation theory of representation of the properad $\mathcal{A}ss\mathcal{B}i$ of associative bialgebras. As this example has never been rigorously treated in the literature before, we show full details here.

As the properad $\mathcal{A}ss\mathcal{B}i$ is homotopy Koszul (see Section 5.4) it admits a minimal resolution, $\mathcal{A}ss\mathcal{B}i_\infty = (\mathcal{F}(\mathcal{C}), \partial)$, which is generated by a relatively small $\mathbb{S}$-bimodule $\mathcal{C} = \{\mathcal{C}(m,n)\}_{m,n\geq 1, m+n\geq 3}$,

$$\mathcal{C}(m,n) := s^{m+n-3}\mathbb{K}[\mathbb{S}_m]\otimes\mathbb{K}[\mathbb{S}_n] = \mathrm{span}$$



The differential ∂ in $\mathcal{A}ss\mathcal{B}i_\infty$ is neither quadratic nor of genus 0. The derivation ∂ on $\mathcal{F}(\mathcal{C})$ is equivalent to a structure of homotopy coproperad on $s^{-1}\mathcal{C}$. The values of ∂ on $(1,n)$- and $(m,1)$-corollas are given, of course, by the well-known A_∞-formulae, while


∂
1
2
1
2
...
n–1
n
=
$\sum_{k=0}^{n-2}\sum_{l=2}^{n-k}(-1)^{k+l(n-k-l)+1}$
1
2
1 ... k
k+l+1 ... n
k+1...k+l
$-\sum_{k=2}^{\infty}\sum_{r_1+\cdots+r_k=n}(-1)^s$
Δ^{SU}
...
k
...
...
...
...
r_1
r_2
r_k

where

$$s = (k-1)(r_1-1) + (k-2)(r_2-1) + \ldots + 1(r_k-1),$$

Δ^{SU} is the Saneblidze-Umble diagonal, and the horizontal line means *fraction* composition from [Mar06]. The meaning of this part of the differential is clear: it describes A_∞-morphisms between an A_∞-structure on X and the associated Saneblidze-Umble diagonal A_∞-structure on $X\otimes X$. Explicitly, this formula is obtained by first considering the quasi-free resolution of the 2-colored operad coding two associative algebras and a morphism between them. While the resolution of the associative operad is given by the associahedra, this resolution is given by the multiplihedra. This resolution gives the relaxed notion of A_∞-algebra and morphism of A_∞-algebras at the same time. Then, to get the formula above, we applied this resolution to the A_∞-algebra X and to $X\otimes X$ with the A_∞-algebra structure induced by the Saneblidze-Umble diagonal.

The values of ∂ on corollas of the form



describe a *homotopy* between two natural A_∞-morphisms from X to $X\otimes X\otimes X$, values on corollas with 4 output legs — homotopies between homotopies etc. We conjecture that $(\mathcal{A}ss\mathcal{B}i_\infty, \partial)$ is a one coloured version of a certain $\mathbb{N}$-coloured properad describing A_∞-algebras, morphisms of A_∞-algebras, homotopies between morphisms of A_∞-algebras, homotopies of homotopies etc., and we hope to describe it in a future publication.

It was proven in [Mar06, MV03] that there exists a minimal model $(\mathcal{A}ss\mathcal{B}i_\infty, \partial)$ such that the differential preserves Kontsevich's path grading of $\mathcal{A}ss\mathcal{B}i_\infty$ and has the form $\partial = \partial_0 + \partial_{pert}$, where ∂_0 describes the minimal resolution, $\frac{1}{2}\mathcal{A}ss\mathcal{B}_\infty$ of the prop of $\frac{1}{2}$-bialgebras (these facts follow also immediately from our Corollary 42). The perturbation part, ∂_{pert}, is a linear combination of so called *fractions* and their compositions. We shall assume from now on that ∂ has all these properties. By checking genus of these fractions (or by referring to our proof of homotopy Koszulness of $\mathcal{A}ss\mathcal{B}i$ in Section 5.4) one can easily obtain the following useful (for our purposes)

Fact 81. *The differential ∂_0 is precisely the quadratic part of ∂, i.e. it is equal to the composition,*

$$\partial_0 : \mathcal{C} \xrightarrow{\partial} \mathcal{F}(\mathcal{C}) \xrightarrow{proj} \mathcal{F}(\mathcal{C})^{(2)}.$$

Let $\mathcal{Q}$ be a dg properad. By Theorem 62, the vector space,

$$s^{-1}\mathrm{Hom}^{\mathbb{S}}(\mathcal{C}, \mathcal{Q}) = \bigoplus_{\substack{n,m\geq 1\\ m+n\geq 3}} s^{2-m-n}\mathcal{Q}(m,n) =: \mathfrak{g}_{GS}(\mathcal{Q}),$$

has a canonical homotopy non-symmetric properad and L_∞-structure Q, whose Maurer-Cartan elements are morphisms of properads $\mathcal{F}(\mathcal{C}) \to \mathcal{Q}$.

If $\gamma : \mathcal{A}ss\mathcal{B}i \to \mathcal{Q}$ is a representation of $\mathcal{A}ss\mathcal{B}i$, or more generally of $\mathcal{A}ss\mathcal{B}i_\infty : \mathcal{A}ss\mathcal{B}i_\infty \to \mathcal{Q}$, then, by Definition 8.2, there exists an associated twisted L_∞-structures, $Q^\gamma = \{Q^\gamma_n\}_{n\geq 1}$, on $\mathfrak{g}_{GS}(\mathcal{Q})$ which controls deformations of γ in the class of representations of $\mathcal{A}ss\mathcal{B}i_\infty$. An explicit formula for the differential ∂ would induce an explicit L_∞-structure. Once again, our main example of this deformation theory is given by $\mathcal{Q} = \mathrm{End}_X$. In this case, the complex above is the deformation complex of associative bialgebra, or more generally of $\mathcal{A}ss\mathcal{B}i_\infty$-gebra, structure on X.

When X is an associative bialgebra, Gerstenhaber and Schack defined in [GS90] a bicomplex whose homology has nice properties with respect to the deformations of the associative bialgebra structure (see also [LM91]). Let us first extend this definition to any properad $\mathcal{Q}$ and not only End_X.

Definition. Let $\gamma : \mathcal{A}ss\mathcal{B}i \to \mathcal{Q}$ be a representation of $\mathcal{A}ss\mathcal{B}i$. We define the *Gerstenhaber-Schack bicomplex of* γ by $C^{m,n} := \mathcal{Q}(m,n)$ and the differentials by





where the general (m,n)-corollas have to be understand as elements of $\mathcal{Q}(m,n)$. The binary corollas are the image under γ of the generating product and coproduct of $\mathcal{A}ss\mathcal{B}i$. Finally, this pictures represent the composition of all these elements in $\mathcal{Q}$.

Let us compare these with the Gerstenhaber-Schack differential, d_{GS}, in the bicomplex $C^{m,n}_{GS} := \mathrm{Hom}(X^{\otimes n}, X^{\otimes m})$ which is defined by [GS90]

$$d_{GS} = d_h + d_v,$$

with $d_h : \mathrm{Hom}(X^{\otimes n}, X^{\otimes m}) \to \mathrm{Hom}(X^{\otimes n+1}, X^{\otimes m})$ given on an arbitrary $f \in \mathrm{Hom}(X^{\otimes n}, X^{\otimes m})$ by

$$\begin{aligned}(d_h f)(a_0, a_1, \ldots, a_n) \quad &:= \quad \Delta^m(a_0)\Box f(a_1, a_2, \ldots, a_n) - \sum_{i=0}^{n-1}(-1)^i f(a_1, \ldots, a_i a_{i+1}, \ldots, a_n) \\ &\qquad +(-1)^{n+1} f(a_1, a_2, \ldots, a_n)\Box\Delta^m(a_n) \quad \forall \;\; a_i \in X.\end{aligned}$$

Here the multiplication in X is denoted by juxtaposition, the induced multiplication in the algebra $X^{\otimes m}$ by $\Box$, the comultiplication in X by Δ, and

$$\Delta^n : (\Delta \otimes \mathrm{Id}^{\otimes m-2}) \circ (\Delta \otimes \mathrm{Id}^{\otimes m-3}) \circ \ldots \circ \Delta : X \to X^{\otimes m}.$$

The expression for d_v is an obvious "dual" analogue of d_h. Now let us represent d_h in graphical terms by associating the graphs



and


to comultiplication and, respectively, multiplication while the corolla



to f. Then the r.h.s of the formula for d_h reads,



which is precisely the first three summands in the previous definition. The other three terms correspond to d_v. Therefore, when $\mathcal{Q} = \mathrm{End}_X$ and $\gamma : \mathcal{A}ss\mathcal{B}i \to \mathrm{End}_X$ is an associative bialgebra structure on X, the preceding bicomplex is exactly Gerstenhaber-Schack bicomplex [GS90].

However, even without an explicit minimal model of $\mathcal{A}ss\mathcal{B}i$, we can show the following general result.

Theorem 82. *Let* $(\mathcal{A}ss\mathcal{B}i_\infty, \partial) \xrightarrow{\pi} \mathcal{A}ss\mathcal{B}i$*, be a minimal model of the properad of bialgebras and* $\gamma : \mathcal{A}ss\mathcal{B}i \to \mathcal{Q}$ *an arbitrary representation of* $\mathcal{A}ss\mathcal{B}i$*. Then the differential,*

$$Q_1^\gamma = Q \circ e^{\gamma\odot}$$

associated to this minimal model in the twisted L_∞*-structure,* Q^γ*, on* $\mathfrak{g}_{GS}$*, is isomorphic to the Gerstenhaber-Schack differential. Hence the deformation complex of representation of* $\mathcal{A}ss\mathcal{B}i$ *is isomorphic to the Gerstenhaber-Schack bicomplex.*

Proof. Let $(\mathcal{A}ss\mathcal{B}i_\infty = \mathcal{F}(\mathcal{C}), \partial)$ be a minimal model of the properad of bialgebras, and let I be the ideal in $\mathcal{F}(\mathcal{C})$ generated by graphs in $\mathcal{F}(\mathcal{C})^{(\geq 2)}$ with at least two non-binary (i.e. neither  nor ) vertices, and let

$$\mathsf{B} := \frac{\mathcal{A}ss\mathcal{B}i_\infty}{(I, \partial I)}$$

be the associated quotient dg properad. The induced differential in B we denote by ∂_{ind}. It is precisely this quotient part, ∂_{ind}, of the total differential ∂ which completely determines the L_∞-differential differential Q_1^γ. Thus our plan is the following: in the next Lemma we present an explicit, up to an automorphism, form of the differential ∂_{ind} (despite the fact that ∂ is not explicit !) and thereafter compare the resulting Q_1^γ with the Gerstenhaber-Schack definition.

The major step in the proof is the following Lemma (in its formulation we use fraction notations again).

Lemma 83. *(i) The derivation, d, of* B *given on generators by,*

(9)


(10)


and, for all other generators with $m + n \geq 4$, by

(11)


is a differential.

(ii) The dg properads $(\mathsf{B}, \partial_{ind})$ *and* (B, d) *are isomorphic.*

Proof. (i) It is easy to see that among for 2-vertex connected binary graphs[6] attached to any other graph in B the bialgebra relations,



hold. Using this fact it is an easy and straightforward calculation to check that $d^2 = 0$. We omit the details. (In fact we shall show below that d is essentially a graph encoding of the Gerstenhaber-Schack differential d_{GS} so this calculation is essentially identical to the one which establishes $d^2_{GS} = 0$.)

(ii) We begin our proof of Lemma 83(ii) with the following

Claim 1. *The natural projection $p : (\mathsf{B}, d) \to \mathcal{A}ss\mathcal{B}i$ is a quasi-isomorphism.*

Indeed, the dg properad (B, d) has a natural increasing and bounded above filtration[7], $\{\mathsf{F}_{-p}\mathsf{B}\}_{p\geq 0}$, with $\mathsf{F}_{-p}\mathsf{B}$ being the span of equivalence classes of graphs which admit a representative in $\mathcal{A}ss\mathcal{B}i_\infty^{(\geq p)}$. As the differential d is connected and preserves the induced path gradation, the associated spectral sequence $(\mathsf{E}_r, \mathsf{d}_r)$ converges to $H^\bullet(\mathsf{B}, d)$. The 0th term $(\mathsf{E}_0, \mathsf{d}_0)$ has the differential given on generators by

(12)


and, for all other generators with $m + n \geq 4$,

(13)


We want to compute homology, $\mathsf{E}_1 = H^\bullet(\mathsf{E}_0, \mathsf{d}_0)$, of this complex and show that $\mathsf{E}_1 \simeq \mathcal{A}ss\mathcal{B}i$. For this purpose consider a 3-step filtration, $0 \subset F_{-2} \subset F_{-1} \subset F_0 = \mathsf{E}_0$, of the complex $(\mathsf{E}_0, \mathsf{d}_0)$ with



and let $(\mathcal{E}_r, \partial_r)$ be the associated spectral sequence. The differential ∂_0 is zero on the generators of F_{-1} and is equal to d_0 on all the other generators. Thus, modulo shifts of gradings, actions of finite groups and tensor products by trivial (i.e. with zero differential) complexes, the complex $(\mathcal{E}_0, \partial_0)$ is isomorphic to the tensor product of two isomorphic operadic complexes (one with "time" flow reversed upside down relative to another) which were studied on page 40 of [MMS06] and which have the differential (in notations of that paper) given by



It is shown in [MMS06] that the cohomology of this complex is concentrated in degree 0 and is isomorphic to the operad of associative algebras. In our context this result immediately implies that $(\mathcal{E}_1, \partial_1)$ is isomorphic to F_{-1} with the differential ∂_1 given on generators by (12). Its cohomology is obviously concentrated in degree 0 (and is equal, in fact, to the properad, $\frac{1}{2}\mathsf{B}$, of infinitesimal bialgebras). Hence $H^\bullet(\mathsf{B}, d)$ is concentrated in degree 0 proving Claim 1.

---

[6] Equivalence classes of graphs in B we call simply graphs for shortness.

[7] One might prove Claim 1 using another filtration of B by the number of vertices and Fact 81 provided one assumes (without any losses) that B is completed with respect to this filtration.

Claim 2. *The natural projection $\pi : (\mathsf{B}, \partial_{ind}) \to \mathcal{A}ss\mathcal{B}i$ is a quasi-isomorphism.*

Indeed, the defined above filtration, $\{\mathsf{F}_{-p}\mathsf{B}\}_{p\geq 0}$, by the number of vertices is also compatible with the differential ∂_{ind}. Let (E_r, d_r) be the associated spectral sequence. Its first nontrivial term, (E_1, d_1) is, by Fact 81, isomorphic to the complex $(\mathsf{E}_1, \mathsf{d}_1)$ above. Hence we can apply the same reasoning as in the proof of Claim 1.

Claim 3. *The exits a morphism of dg properads Φ making the diagram*

$$\begin{tikzcd} & (\mathsf{B}, d) \arrow[d, "p"] \\ (\mathcal{A}ss\mathcal{B}i_\infty, \partial) \arrow[ur, "\Phi"] \arrow[r, "\pi"'] & (\mathcal{A}ss\mathcal{B}i, 0) \end{tikzcd}$$

commutative.

Since $\mathcal{A}ss\mathcal{B}i$ is a properad concentrated in degree 0, the map p is surjective. Since p is a quasi-isomorphism by Claim 2, it is an acyclic fibration in the model category of properads (see Appendix A). By Corollary 97, $\mathcal{A}ss\mathcal{B}i_\infty$ is a cofibrant properad. Finally, the morphism Φ is given by the left lifting property in the model category of properads. Hence, the existence of Φ is clear but we need to make it more precise. We construct it as follows and refine it in Claim 4.

As $\mathcal{A}ss\mathcal{B}i_\infty = \mathcal{F}(\mathcal{C})$ is a free properad, a morphism Φ is completely determined by its values on the generating (m,n)-corollas which span the vector space $\mathcal{C}$, and one can construct Φ by a simple induction [8] on the degree, $r := m + n - 3 \geq 0$, of such corollas. For $r = 0$ we set Φ to be identity, i.e.



Assume we constructed values of Φ on all corollas of degree $r \leq N$. Let e be a generating corolla of non-zero weight $r = N + 1$. Note that δe is a linear combination of graphs whose vertices are decorated by corollas of weight $\leq N$ (as differential δ has degee -1). Then, by induction, $\Phi(\delta e)$ is a well-defined element in B. As $\pi(e) = 0$, the element,

$$\Phi(\delta e)$$

is a closed element in B which projects under p to zero. By Claim 1, the surjection p is a quasi-isomorphism. Hence this element is exact and there exists $\mathfrak{e} \in \mathsf{B}$ such that

$$d\mathfrak{e} = \Phi(\delta e).$$

We set $\Phi(e) := \mathfrak{e}$ completing thereby inductive construction of Φ.

Claim 4. *A morphism Φ can be chosen so that*



$+$ terms with ≥ 2 number of vertices.

Indeed, the differential d in B has the form,

$$d = \mathsf{d}_1 + d_{rest},$$

where d_1 is the quadratic *differential* in B defined by (13) and the part d_{part} corresponds to graphs lying in $F_{-3}\mathsf{B}$. We shall prove Claim 4 by induction on the degree $r = m+n-3$ of the generating (m,n)-corollas in $\mathcal{A}ss\mathcal{B}i_\infty$ (cf. proof of Theorem 43 in [Mar06]). For $r = 0$ the Claim is true. Assume we have already constructed Φ such that the claim is true for values of Φ on corollas with non-zero degree $\leq N$ and consider a generating corolla, e, of degree $N+1$. The value, $\mathfrak{e} := \Phi(e)$, is a solution of the equation,

---

[8] this induction is a straightforward analogue of the Whitehead lifting trick in the theory of CW-complexes in algebraic topology.

$$\mathsf{d}_1\mathfrak{e} + d_{rest}\mathfrak{e} = \Phi(\partial_0 e) + \Phi(\partial_{pert} e). \tag{14}$$

Let π_1 and π_2 denote projections in B to the subspaces spanned by equivalence classes of graphs with 1 and, respectively, 2 vertices. Then equation (14) implies,

$$\pi_2 \circ \mathsf{d}_1(\mathfrak{e}) = \mathsf{d}_1 \circ \pi_1(\mathfrak{e}) = \pi_2 \circ \Phi(\partial_0 e),$$

as both $d_{rest}\mathfrak{e}$ and $\Phi(\partial_{pert}e)$ are spanned by graphs lying in $F_{-3}\mathsf{B}$. Using now the explicit form for the differential ∂_0 (given, e.g., by formula (14) in [Mar06]) and the induction assumption we immediately conclude that

$\pi_1(\mathfrak{e}) =$ 

completing the proof of Claim 4.

Claim 5. *The morphism Φ induces a dg isomorphism $(\mathsf{B}, \partial_{ind}) \to (\mathsf{B}, d)$.*

Indeed, Φ sends the ideal I to zero. Since Φ respects differentials, it sends the ideal $(I, \partial I)$ to zero as well and hence induces, by Claims 3 and 4, a required isomorphism. This completes proof of Lemma 83. ☐

Now we continue with the proof of Theorem 82. The differential Q_1^γ in the graded vector space $\mathfrak{g}_{GS}(\mathcal{Q}) = \oplus_{m,n} s^{2-m-n}\mathcal{Q}$ is completely determined by the quotient differential, ∂_{ind}, of the full differential ∂ in $\mathcal{A}ss\mathcal{B}i_\infty$. By Lemma 83, this quotient differential is given, up to automorphims, by formulae (9)-(11). The proof of the Theorem 82 is completed. ☐

As a direct corollary, we have

Corollary 84. *The Gerstenhaber-Schack bicomplex of an associative bialgebra X is a homotopy non-symmetric properad and a twisted L_∞-algebra whose Maurer-Cartan elements are deformations of the first structure.*

The homotopy non-symmetric properad structure induces, on this chain complex, (homotopy) LR-operations which play the same role than the non-symmetric braces for Hochschild cochain complex. They are are expected to be used in the proof of a Deligne conjecture for associative bialgebras.

9.4. **Twisted L_∞-algebras and dg prop(erad)s.** For any quasi-free prop(erad) $(\mathcal{P}_\infty = \mathcal{F}(s^{-1}\mathcal{C}), \partial_\mathcal{P})$ and any prop(erad) $\mathcal{Q}$ there exists, in accordance with Theorem 62, a canonical L_∞ structure, Q, on the graded vector space $s^{-1}\mathrm{Hom}_\bullet^{\mathbb{S}}(\mathcal{C}, \mathcal{Q})$ whose Maurer-Cartan elements are in one-to-one correspondence with representation of P_∞ in $\mathcal{Q}$. If γ is any particular representation of P_∞, then the associated twisted L_∞-algebra, Q^γ, describes deformation theory of γ within the class of representation of $\mathcal{P}_\infty$ (see §7.2). Remarkably, *there always exists a quasi-free prop(erad) $(\mathcal{P}_\infty^{(2)}, \partial)$ whose representations in $\mathcal{Q}$ are in one-to-one correspondence with pairs, (γ, Γ), where γ is a representation of $\mathcal{P}_\infty$ on $\mathcal{Q}$ and Γ is a MC element in Q^γ.* Thus the dg prop(erad) $\mathcal{P}_\infty^{(2)}$ gives a complete description of the deformation theory of a generic representation of $\mathcal{P}_\infty$. In fact, this constructions can be obviously iterated giving rise to quasi-free prop(erad)s $\mathcal{P}_\infty^{(3)}$, $\mathcal{P}_\infty^{(4)}$ etc.

By definition, $\mathcal{P}_\infty^{(2)}$ is a free prop(erad) on the $\mathbb{S}$-bimodule $s^{-1}\mathcal{C} \oplus s^{-1}\mathcal{C}$ but the differential, ∂, in $\mathcal{P}_\infty^{(2)}$ is *not* a direct sum, $\partial_\mathcal{P} \oplus \partial_\mathcal{P}$, of differentials in $\mathcal{P}_\infty$. We illustrate the above claim in the case of $\mathcal{P} = \mathcal{A}ss$, the operad of associative algebras, before giving the general definition.

Let $\mathcal{A}ss_\infty^{(2)}$ be a quasi-free operad generated by an $\mathbb{S}$-module,


$s^{n-2}\mathbb{K}[\Sigma_n] \oplus s^{n-2}\mathbb{K}[\Sigma_n] \simeq$ span
$\sigma(1)$ $\sigma(2)$ ... $\sigma(n)$
,
$\sigma(1)$ $\sigma(2)$ ... $\sigma(n)$
$\sigma\in\mathbb{S}_n$

and equipped with a differential given on generators by,


∂
1 2 ... n–1 n
$= \sum_{k=0}^{n-1}\sum_{l=1}^{n-k}(-1)^{(l-1)(n-k-l)}$
1 ... k k+l+1 ... n
k+1...k+l
,


∂
1 2 ... n–1 n
$= \sum_{k=0}^{n-1}\sum_{l=1}^{n-k}(-1)^{(l-1)(n-k-l)}$
1 ... k k+l+1 ... n
k+1...k+l
+
+
1 ... k k+l+1 ... n
k+1...k+l
+
1 ... k k+l+1 ... n
k+1...k+l

Proposition 85. *There is a one-to-one correspondence between representations of the dg operad $\mathcal{A}ss_\infty^{(2)}$ in an operad $\mathcal{Q}$ and degree -1 elements, γ and Γ, in the deformation complex (Hochschild complex) $\oplus_{n\geq 2}s^{1-n}\mathcal{Q}(n)$ such that,*

$$\begin{aligned} [d+\gamma, d+\gamma]_G &= 0 \\ [d+\gamma,\Gamma]_G + \frac{1}{2}\,[\Gamma,\Gamma]_G &= 0, \end{aligned}$$

where $[\ ,\ ]_G$ stands for the Gerstenhaber brackets.

Proof is obvious and hence is omitted. The data (d,γ) describes a representation of $\mathcal{A}ss_\infty$ on $\mathcal{Q}$, and the data Γ describes a deformation of this representation.

In general, $\mathcal{P}_\infty^{(2)}$ is the prop(erad) given by $\mathcal{F}(s^{-1}\mathcal{C}^\bullet \oplus s^{-1}\mathcal{C}^\blacktriangledown)$. We denote by $\partial_\mathcal{P}(c) = \sum \mathcal{G}(c_1,\ldots,c_n)$ the image under the differential $\partial_\mathcal{P}$ of an element c of $s^{-1}\mathcal{C}$, with $c_1,\ldots,c_n \in \mathcal{C}$. The differential ∂ of $\mathcal{P}_\infty^{(2)}$ is defined by

$$\begin{aligned} \partial(c^\bullet) &:= \sum \mathcal{G}(c_1^\bullet,\ldots,c_n^\bullet) \quad \text{for} \quad c^\bullet \in s^{-1}\mathcal{C}^\bullet, \\ \partial(c^\blacktriangledown) &:= \sum \mathcal{G}(c^{i_1},\ldots,c_n^{i_n}) \quad \text{for} \quad c^\blacktriangledown \in s^{-1}\mathcal{C}^\blacktriangledown, \end{aligned}$$

where the $i_1,\ldots,i_n$ are in $\{\bullet, \blacktriangledown\}$ with at least one equal to $\blacktriangledown$. It is easy to see that $\partial^2 = 0$.

Proof. The formula for $\partial(c^\bullet)$ gives the first relation. With the formula for $\partial(c^\blacktriangledown)$, it gives the second one. ☐

Proposition 86. *There is a one-to-one correspondence between representations of the dg prop(erad) $\mathcal{P}_\infty^{(2)}$ in an prop(erad) $\mathcal{Q}$ and degree -1 elements, γ and Γ, in the deformation complex $s^{-1}\mathrm{Hom}_\bullet^\mathbb{S}(\mathcal{C},\mathcal{Q})$ such that*

$$\begin{aligned} Q(\gamma) &= 0 \\ Q^\gamma(\Gamma) = Q(\gamma+\Gamma) &= 0, \end{aligned}$$

where Q stands for the L_∞-algebra structure.

In the following proposition, we interpret $\mathcal{A}ss_\infty^{(2)}$ as the Koszul resolution of a new operad, denoted by $\mathcal{A}ss^{(2)}$.

Proposition 87. *The dg operad $(\mathcal{A}ss_\infty^{(2)}, \partial)$ is a quadratic resolution of a quadratic operad $\mathcal{A}ss^{(2)}$ defined as the quotient of the free operad on the $\mathbb{S}$-module*

$A(n) := \mathbb{K}[\mathbb{S}_2] \oplus \mathbb{K}[\mathbb{S}_2] = \textit{span}$



$_{\sigma\in\Sigma_2}$ *for $n = 2$*

0 *otherwise,*

modulo the ideal generated by relations,



$$= 0, \qquad \forall \sigma \in \mathbb{S}_3.$$

and



$$= 0, \qquad \forall \sigma \in \mathbb{S}_3.$$

Proof. Let $F_{-p}(\mathcal{A}ss_\infty^{(2)})$ be the subspace of $\mathcal{A}ss_\infty^{(2)}$ spanned by trees with a least p internal edges between one vertex labelled by $\bullet$ and the other one labelled by $\blacktriangledown$. This defines an increasing filtration with is bounded on $\mathcal{A}ss_\infty^{(2)}(n)$ for each n. Therefore it converges to the homology of $\mathcal{A}ss_\infty^{(2)}(n)$ by the Classical Convergence Theorem 5.5.1 of [Wei94]. The first term E^0_{pq} is equal to the subspace of $\mathcal{A}ss_\infty^{(2)}$ spanned by trees with exactly p internal edges between one vertex labelled by $\bullet$ and the other one labelled by $\blacktriangledown$. And the differential d^0 is equal to the sum of the differentials $\partial_{\mathcal{A}ss_\infty^\bullet} + \partial_{\mathcal{A}ss_\infty^\blacktriangledown}$, that it is splits $\bullet$ and $\blacktriangledown$ vertices into pure $\bullet$ and $\blacktriangledown$ trees. Hence (E^0, d^0) is the coproduct $\mathcal{A}ss_\infty^\bullet \vee \mathcal{A}ss_\infty^\blacktriangledown$ (see Section A.3) of two resolutions of $\mathcal{A}ss$, which is acyclic. Finally, we have $E^1_{pq} = 0$ for $p + q \neq 0$ and $\bigoplus_{p\geq 0} E^1_{-p\,p} = \mathcal{A}ss^\bullet \vee \mathcal{A}ss^\blacktriangledown$. The spectral sequence collapses and the homology of $\mathcal{A}ss_\infty^{(2)}$ is concentrated in degree 0. Another presentation of this homology group is given by the quotient of the free operad on degree 0 elements, namely the two binary products $\bullet$ and $\blacktriangledown$, by ideal generated by the image under ∂ of the degree 1 elements of $\mathcal{A}ss_\infty^{(2)}$. ☐

In other words, the operad $\mathcal{A}ss^{(2)}$ is Koszul. A representation of $\mathcal{A}ss^{(2)}$ in a vector space X is equivalent to a pair of linear maps $\mu : X \otimes X \to X$ and $\nu : X \otimes X \to X$ such that both (X, μ) and $(X, \mu + \nu)$ are associative algebras. As a corollary, we get the following isomorphism of $\mathbb{S}$-modules $\mathcal{A}ss^{(2)} \cong \mathcal{A}ss \vee \mathcal{A}ss$.

Remark. The example of $\mathcal{A}ss^{(2)}$ is also interesting from the viewpoint of Koszul operad. It comes from a set theoretic operad. It is Koszul whereas the method of [Val06a] cannot be applied because $\mathcal{A}ss^{(2)}$ is not basic set, that is the composition of operations is not injective. The product $\blacktriangledown$ has an "absorbing" effect.

In the same way, we define the operad $\mathcal{L}ie^{(2)}$ by $\mathcal{F}([\,,\,]_\bullet \oplus [\,,\,]_\blacktriangledown)/(\mathrm{Jac}^\bullet_\bullet \oplus (\mathrm{Jac}^\bullet_\blacktriangledown + \mathrm{Jac}^\blacktriangledown_\bullet + \mathrm{Jac}^\blacktriangledown_\blacktriangledown))$, where $[\,,\,]_\bullet$ and $[\,,\,]_\blacktriangledown$ stand for two skew-symmetric brackets and where Jac^b_a stands for the "Jacobi" relation $[[X,Y]_a, Z]_b + [[Y,Z]_a, X]_b + [[Z,X]_a, Y]_b = 0$. This operad enjoys the same properties with $\mathcal{L}ie$ than the non-symmetric operad $\mathcal{A}ss^{(2)}$ with $\mathcal{A}ss$ explained above. More generally, to any binary quadratic operad $\mathcal{P}$ (eventually non-symmetric) with its minimal model $\mathcal{P}_\infty$, we can associate an operad $\mathcal{P}^{(2)}$ such that $\mathcal{P}_\infty^{(2)}$ is its minimal model.

We summary this result with the explicit form of $\mathcal{P}^{(2)}$ in terms of Manin products in the following theorem. Let $\mathcal{P} = \mathcal{F}(V)/(R)$ and $\mathcal{Q} = \mathcal{F}(W)/(S)$ be two binary quadratic non-symmetric operads. There exists a morphism of $\psi : \mathcal{F}(V)(3) \otimes \mathcal{F}(W)(3) \to \mathcal{F}(V \otimes W)(3)$. *Manin's black square product of $\mathcal{P}$ and $\mathcal{Q}$* is equal to $\mathcal{P} \blacksquare \mathcal{Q} := \mathcal{F}(V \otimes W)/(\psi(R \otimes S))$. In the symmetric case, the definition is similar but the morphism ψ is more involved and requires the use of signature representations (see [Val06b] for more details).

Theorem 88. *For any binary quadratic non-symmetric operad $\mathcal{P}$ which admits a minimal model $\mathcal{P}_\infty$, the non-symmetric operad $\mathcal{P}_\infty^{(2)}$ is a minimal model (resolution) of $\mathcal{P} \blacksquare \mathcal{A}ss^{(2)}$, which is isomorphic, as a graded module, to $\mathcal{P} \vee \mathcal{P}$, where the coproduct has to be taken in the category of non-symmetric operads .*
For any binary quadratic operad $\mathcal{P}$ which admits a minimal model $\mathcal{P}_\infty$, the operad $\mathcal{P}_\infty^{(2)}$ is a minimal model (resolution) of $\mathcal{P} \bullet \mathcal{L}ie^{(2)}$ which is isomorphic, as an $\mathbb{S}$-module to $\mathcal{P} \vee \mathcal{P}$.

Proof. By the same argument as in Proposition 87 above, the homology of $\mathcal{P}_\infty^{(2)}$ is concentrated in degree 0. If we denote the quadratic operad $\mathcal{P}$ by $F(V)/(\mathcal{R})$, this non-trivial homology group is equal to $F(V_\bullet \oplus V_\blacktriangledown)/(\mathcal{R}_\bullet^\bullet \oplus \mathcal{R}_\blacktriangledown^\bullet + \mathcal{R}_\bullet^\blacktriangledown + \mathcal{R}_\blacktriangledown^\blacktriangledown)$, which is equal to the black product of $\mathcal{P}$ with $\mathcal{L}ie^{(2)}$. □

Appendix A. Model category structure for prop(erad)s

In this appendix, we prove that the categories of props and properads have a cofibrantly generated model category structure. We make precise coproducts, pushouts, cofibrations and cofibrant objects. We refer the reader to the book [Hov99] of M. Hovey for a comprehensive treatment of model categories. (In order to be self-contained in this appendix, with we will not avoid the prefix dg here.)

Let us denote by *dg Properads* the category of dg properads and by *dg Props* the category of dg props. It means either the category of reduced dg prop(erad)s ($\mathcal{P}(m, n) = 0$ for $n = 0$ or $m = 0$) or the category of dg prop(erad)s over a field of characteristic 0. By default, we work over unbounded chain complexes but the following proofs hold over bounded chain complexes as well. We transfer the cofibrantly generated model category structure of $\mathbb{S}$-bimodules to the category of prop(erad)s via the free prop(erad) functor.

A.1. **Model category structure on $\mathbb{S}$-bimodules.**

The category of dg $\mathbb{S}$-bimodules is endowed with a cofibrantly generated model category structure coming from the cofibrantly generated model category structure on dg $\mathbb{K}$-modules.

Recall from [Hov99] Theorem 2.3.11 that the category of dg R-modules has a cofibrantly generated model category structure for any ring R. Quasi-isomorphisms form the class of weak equivalences and degreewise surjective maps form the class of fibrations. Let us make explicit the (generating) acyclic cofibrations. The model category of dg $\mathbb{K}$-modules is cofibrantly generated by the acyclic cofibrations $J^k : 0 \to D^k$, where D^k is the chain complex

$$\cdots \to 0 \to \underbrace{\mathbb{K}}_{k} \xrightarrow{\mathrm{Id}} \underbrace{\mathbb{K}}_{k-1} \to 0 \to \cdots$$

and by the cofibrations $I^k : S^{k-1} \to D^k$, where S^{k-1} is the following chain complex

$$\cdots \to 0 \to \underbrace{\mathbb{K}}_{k-1} \to 0 \to \cdots .$$

For any $m, n \in \mathbb{N}$, the category of left $\mathbb{S}_m$ and right $\mathbb{S}_n$-bimodules is the category of dg modules over the group ring $\mathbb{K}[\mathbb{S}_m^{\mathrm{op}} \times \mathbb{S}_n]$. By Theorem 2.3.11 of [Hov99] the preceding theorem, it has a cofibrantly generated model category structure, where the generating acyclic cofibrations are the maps $J^k_{m,n}\ :\ 0 \to D^k_{m,n}$, where $D^k_{m,n}$ is the acyclic dg $\mathbb{K}[\mathbb{S}^{\mathrm{op}}_m \times \mathbb{S}_n]$-module

$$\cdots \to 0 \to \underbrace{\mathbb{K}[\mathbb{S}^{\mathrm{op}}_m \times \mathbb{S}_n]}_{k} \xrightarrow{\mathrm{Id}} \underbrace{\mathbb{K}[\mathbb{S}^{\mathrm{op}}_m \times \mathbb{S}_n]}_{k-1} \to 0 \to \cdots$$

and where the generating cofibrations are the maps $I^k_{m,n}\ :\ S^{k-1}_{m,n} \to D^k_{m,n}$, with $S^{k-1}_{m,n}$ the following dg $\mathbb{K}[\mathbb{S}^{\mathrm{op}}_m \times \mathbb{S}_n]$-module

$$\cdots \to 0 \to \underbrace{\mathbb{K}[\mathbb{S}^{\mathrm{op}}_m \times \mathbb{S}_n]}_{k-1} \to 0 \to \cdots .$$

Since the category of dg $\mathbb{S}$-bimodules is the product over $(m,n) \in \mathbb{N}^2$ of the model categories of left $\mathbb{S}_m$ and right dg $\mathbb{S}_n$-bimodules, it is naturally endowed with a term-by-term cofibrantly generated model category structure. The set of generating acyclic cofibrations can be chosen to be $J = \{\widetilde{J}^k_{m,n};\ k \in \mathbb{Z}, m,n \in \mathbb{N}\}$, where $\widetilde{J}^k_{m,n}$ is equal to $J^k_{m,n}\ :\ 0 \to D^k_{m,n}$ in arity (m,n) and 0 elsewhere. Similarly, the set of generating cofibrations can be chosen to be $I = \{\widetilde{I}^k_{m,n};\ k \in \mathbb{Z}, m,n \in \mathbb{N}\}$, where $\widetilde{I}^k_{m,n}$ is equal to $I^k_{m,n}\ :\ S^{k-1}_{m,n} \to D^k_{m,n}$ in arity (m,n) and 0 elsewhere. Notice that the domains of elements of I or J are sequentially small with respect to any map in the category of dg $\mathbb{S}$-bimodules.

A.2. **Transfer Theorem.** In the section, we recall the theorem of transfer, mainly due to Quillen [Qui67] Section II.4 (see also S. E. Crans [Cra95] Theorem 3.3 and M. Hovey [Hov99] Proposition 2.1.19). We will use it to endow the category of dg prop(erad)s with a model category structure.

Definition (Relative I-cell complexes)**.** For every class I of maps of a category, a *relative I-cell complexes* is a sequential colimit of pushouts of maps of I.

Let us make explicit this type of morphisms. A relative I-cell complex is a map $A_0 \xrightarrow{\varphi} A_\infty$ which comes from a sequential colimit

$$\begin{tikzcd}
A_0 \arrow[r, "\imath_0"] \arrow[d, "\varphi"] & A_1 \arrow[r, "\imath_1"] \arrow[dl] & \cdots \arrow[r] & A_n \arrow[r, "\imath_n"] \arrow[dlll] & A_{n+1} \arrow[r] \arrow[dllll] & \cdots \\
A_\infty := Colim_{\mathbb{N}}\ A_n, & & & & &
\end{tikzcd}$$

where each map $A_n \xrightarrow{i_n} A_{n+1}$ is defined by a pushout

$$\begin{tikzcd}
\vee_\alpha S_\alpha \arrow[r] \arrow[d, "\vee_\alpha j_\alpha"] & A_n \arrow[d, "i_n"] \\
\vee_\alpha T_\alpha \arrow[r] & A_{n+1},
\end{tikzcd}$$

with $j_\alpha \in I$. As usual, we denote the collection of relative I-cell complexes by I-cell.

Theorem 89 ([Qui67] Section II.4, [Cra95] Theorem 3.3, [Hov99] Proposition 2.1.19)**.** *Let $\mathcal{C}$ be a cofibrantly generated model category with I as the set of generating cofibrations and J as the set of generating acyclic cofibrations. Let $F\ :\mathcal{C} \rightleftharpoons \mathcal{D}\ :\ U$ be an adjunction, where F is the left adjoint and U the right adjoint. Suppose that*

1. *$\mathcal{D}$ has finite limits and colimits,*
2. *the functor U preserves filtered colimits,*
3. *the image under U of any relative $F(J)$-cell complex is a weak equivalence in $\mathcal{C}$.*

A map f in $\mathcal{D}$ is defined to be a weak equivalence (resp. fibration) if the associated map $U(f)$ is a weak equivalence (resp. fibration) in $\mathcal{C}$. The class of cofibrations in $\mathcal{D}$ is the class of map that verify the left lifting property (LLP) with respect to acyclic fibrations.

These three classes of maps provide the category $\mathcal{D}$ with a model category structure cofibrantly generated by $F(I)$ as the the set of generating cofibrations and $F(J)$ as the set of generating acyclic cofibrations.

We also refer the reader to Section 2.5 of [BM03] for the application of this Theorem with stronger and sometimes more convenient hypotheses. Remark that Transfer Theorem 89 was used (and rephrased) by V. Hinich in [Hin97] to provide a model category structure to the category of operads over unbounded chain complexes (see Theorem 2.2.1 of [Hin97] and the corrected version of Theorem 6.6.1 in [Hin03]). M. Spitzweck also applied this theorem to prove a general result about model category structures on categories of algebras over a triple (Theorem 1 of [Spi01]).

A.3. **Limits and Colimits of prop(erad)s.** In this section, we prove that the category of prop(erad)s has all limits and finite colimits. We also make explicit the coproducts and pushouts of prop(erad)s.

Proposition 90. *The category of prop(erad)s has all limits.*

Proof. We recall from D. Borisov and Y.I. Manin [BM06] that the free prop(erad) functor induces a triple $\mathcal{F} : \mathbb{S}\text{-biMod} \to \mathbb{S}\text{-biMod}$ such that an algebra over it is a prop(erad). Since the underlying category of $\mathbb{S}$-bimodules has limits, the category of prop(erad)s has all limits (Section 1.5 of [GJ94]). □

To prove that the category of prop(erad)s has finite colimits, we first make explicit coproducts and pushouts. This section is the generalization of Section 1.5 of [GJ94] from operads to prop(erad)s. Once again, the situation is more subtle for prop(erad)s than for operads since it requires the notion of adjacent vertices of a graph (see Section 4.2).

Let $\mathcal{P}$ and $\mathcal{Q}$ be two prop(erad)s. The coproduct of $\mathcal{P}$ and $\mathcal{Q}$ is given by a quotient of the free prop(erad) on their sum $\mathcal{F}(\mathcal{P} \oplus \mathcal{Q})$. On this space, we define an equivalence relation by the following generating relation : if a graph g, with vertices indexed by elements of $\mathcal{P}$ and $\mathcal{Q}$ has two adjacent vertices indexed elements of $\mathcal{P}$ (or $\mathcal{Q}$), it is equivalent to the same graph, where the two adjacent vertices are contracted and the new vertex is labelled by the composition in $\mathcal{P}$ (or $\mathcal{Q}$) of the two associated elements of $\mathcal{P}$ (or $\mathcal{Q}$). The quotient of $\mathcal{F}(\mathcal{P} \oplus \mathcal{Q})$ by this relation is the coproduct of $\mathcal{P}$ and $\mathcal{Q}$. We denote it by $\mathcal{P} \vee \mathcal{Q}$. This $\mathbb{S}$-bimodule has the following basis. It can be represented by the sum over (connected) graphs with vertices indexed by elements of $\mathcal{P}$ and $\mathcal{Q}$ such that no adjacent pair of vertices are labelled by the same kind of elements (see Figure 5).



Figure 5. Element of the coproduct $\mathcal{P} \vee \mathcal{Q}$

Let $\mathcal{P}$, $\mathcal{Q}$ and $\mathcal{R}$ be three prop(erad)s. Let $f : \mathcal{P} \to \mathcal{Q}$ and $g : \mathcal{P} \to \mathcal{R}$ be two morphisms of prop(erad)s. Their pushout is isomorphic to the quotient of $\mathcal{Q} \vee \mathcal{R}$ by the ideal generated by $\{f(p) - g(p),\, p \in \mathcal{P}\}$. (We refer to Appendix B of [Val06b] for the notion of ideal of a prop(erad). The notion of ideal generated by a sub-$\mathbb{S}$-bimodule is also made explicit there.) The pushout $\mathcal{Q} \vee_{\mathcal{P}} \mathcal{R}$ is represented by labelled (connected) graphs as above but further quotient by the following relation : if a vertex is labelled by an element of the form $f(p)$ for $p \in \mathcal{P}$, it can be replaced by the same vertex labelled by the corresponding element $g(p)$ and vice-versa. When this operation generates two adjacent vertices indexed by elements of the same prop(erad), there are to be composed.

Proposition 91. *The category of prop(erad)s has finite colimits.*

Proof. This result can be proved with two methods.
First, recall that the free properad on an $\mathbb{S}$-bimodule V is given by the sum on (connected) graphs without level whose vertices are coherently labelled by elements of V (see Section 2.7 of [Val07a]). We denote it by

$$\mathcal{F}(V) = \left(\bigoplus_{g \in \mathcal{G}_c} \bigotimes_{\nu \in \mathcal{N}(g)} V(|Out(\nu)|,\, |In(\nu)|) \right) \Big/ \approx ,$$

on Theorem 2.3 of [Val07a], where $\mathcal{N}(g)$ is the set of vertices of a graph g. Since the tensor product of dg $\mathbb{S}$-bimodules preserves colimits, the functor

$$\begin{array}{ccc} \mathbb{S}\text{-biMod} & \to & \mathbb{S}\text{-biMod} \\ V & \mapsto & \displaystyle\bigotimes_{\nu \in \mathcal{N}(g)} V(|Out(\nu)|,\, |In(\nu)|), \end{array}$$

associated to any graph g, preserves filtered colimits (see Lemma 1.14 of [GJ94]). Then the triple $\mathcal{F} : \mathbb{S}\text{-biMod} \to \mathbb{S}\text{-biMod}$ associated to the free prop(erad) functor preserves filtered colimits. The argument of Page 16 of [GJ94] proves that the category of prop(erad)s has filtered colimits. Since it has pushouts and filtered colimits, it has finite colimits by Chapter IX of [ML98].
We can also construct coequalizers in this category. Since it is an additive category, it is enough to construct cokernels. Let $f : \mathcal{P} \to \mathcal{Q}$ be a morphism of prop(erad)s. Its cokernel is given by the quotient of $\mathcal{Q}$ with the ideal generated by the image of f. Since it has coproducts and coequalizers, this category has finite colimits by Theorem 2.1 Chapter V of [ML98]. ☐

A.4. **Model category structure.** In this section, we apply the Transfer Theorem 89 to provide a cofibrantly generated model category structure on the category of prop(erad)s.

We consider the free prop(erad) adjunction $\mathcal{F} : \text{dg } \mathbb{S}\text{-biMod} \rightleftharpoons \text{dg Prop(erad)s} : U$. We proved in A.1 that the category on the left hand side is a cofibrantly generated model category. We apply the Transfer Theorem 89 to this adjunction as follows. The generating acyclic cofibrations are $\mathcal{F}(J) = \{I \to \mathcal{F}(D^k_{m,n})\}$ and the generating cofibrations are $\mathcal{F}(I) = \{\mathcal{F}(S^{k-1}_{m,n}) \to \mathcal{F}(D^k_{m,n})\}$.

Lemma 92. *A morphism of dg properads is a relative $\mathcal{F}(J)$-cell complex if and only is it is a map $\mathcal{P} \to \mathcal{P} \vee \mathcal{F}(D)$, where $D = \bigoplus_{d \geq 1} D_i$ is an acyclic dg $\mathbb{S}$-bimodule whose components are free $\mathbb{S}$-bimodules with each D_i equal to a direct sum of dg $\mathbb{S}$-bimodules $D^k_{m,n}$.*
A morphism of dg properads is a relative $\mathcal{F}(I)$-cell complex if and only if it is a map $\mathcal{P} \to \mathcal{P} \vee \mathcal{F}(S)$, where S is a dg $\mathbb{S}$-bimodule, whose components are free $\mathbb{S}$-bimodules, endowed with an exhaustive filtration

$$S_0 = \{0\} \subset S_1 \subset S_2 \subset \cdots \subset Colim_i S_i = S$$

such that $d : S_i \to \mathcal{F}(S_{i-1})$ and such that $S_{i-1} \rightarrowtail S_i$ are split monomorphisms of dg $\mathbb{S}$-bimodules with cokernels isomorphic to a free $\mathbb{S}$-bimodule.

Proof. Pushouts of elements of $\mathcal{F}(J)$ are as follows:

$$\begin{tikzcd} I \arrow[r] \arrow[d, tail, "\vee_\alpha \mathcal{F}(J^\alpha)"] & \mathcal{P} \arrow[d] \\ \bigvee_\alpha \mathcal{F}(D^\alpha) \arrow[r] & \mathcal{P} \vee \left(\bigvee_\alpha \mathcal{F}(D^\alpha)\right), \end{tikzcd}$$

with each D^α equal to a $D^k_{m,n}$. Since the coproduct of free prop(erad)s is the free prop(erad) on the sum of their generating spaces, $\mathcal{F}(V) \vee \mathcal{F}(V') \cong \mathcal{F}(V \oplus V')$, the composite of two such maps

is equal to $\mathcal{P} \to \mathcal{P} \vee \mathcal{F}(\bigoplus_\alpha D^\alpha \oplus \bigoplus_\beta D^\beta)$. Hence a sequential colimit of such pushouts has the form $\mathcal{P} \to \mathcal{P} \vee \mathcal{F}(D)$, with $D = \bigoplus_{d\geq 1} D_i$ an acyclic dg $\mathbb{S}$-bimodule whose components are free $\mathbb{S}$-bimodules.
A pushout of an element of $\mathcal{F}(I)$ is

$$\begin{tikzcd}
\vee_\alpha \mathcal{F}(S^\alpha) \arrow[r, "f"] \arrow[d, tail, "\vee_\alpha \mathcal{F}(I^\alpha)"] & \mathcal{P} \arrow[d] \\
\vee_\alpha \mathcal{F}(D^\alpha) \arrow[r] & \mathcal{Q},
\end{tikzcd}$$

with each S^α equal to an $S^k_{m,n}$ and D^α equal to a $D^k_{m,n}$. We denote by z the image under f of the generating element of $S^{k-1}_{m,n}$. Notice that z is a cycle in $\mathcal{P}$. If we denote by ξ and $d\xi$ the generating elements of $D^k_{m,n}$, the pushout $\mathcal{Q}$ is equal to $\mathcal{P} \vee \mathcal{F}(\bigoplus_\alpha \xi^\alpha.\mathbb{K}[\mathbb{S}^{\mathrm{op}}_m \times \mathbb{S}_n])$ with $d\xi = z$. Therefore a relative $\mathcal{F}(I)$-cell complex is a map $\mathcal{P} \to \mathcal{P} \vee \mathcal{F}(S)$, with S a dg $\mathbb{S}$-bimodule whose components are free $\mathbb{S}$-bimodules. Since a relative $\mathcal{F}(I)$-cell complex is a sequential colimit of such pushouts, the filtration of S is given by this sequential guing of cells. ☐

Theorem 93. *The category of prop(erad)s has a cofibrantly model category structure provided by the following three classes of morphisms. A map* $\mathcal{P} \xrightarrow{f} \mathcal{Q}$ *is a*

- *weak equivalence if and only if it is a quasi-isomorphism of dg $\mathbb{S}$-bimodules, that is a quasi-isomorphism in any arity,*
- *fibration if and only if it is a degreewise surjection in any arity,*
- *cofibration if and only if it has the left lifting property with respect to acyclic fibrations.*

The generating cofibrations are the maps $\mathcal{F}(I) = \{\mathcal{F}(S^{k-1}_{m,n}) \to \mathcal{F}(D^k_{m,n})\}$ *and the generating acyclic cofibrations are the maps* $\mathcal{F}(J) = \{I \to \mathcal{F}(D^k_{m,n})\}$.

Proof. The category of prop(erad)s has finite limits and colimits (1) by the preceding section. To any dg $\mathbb{S}$-bimodule M, we can consider the trivial (abelian) prop(erad) structure on $I \oplus M$, that is the composite product is zero on M. So, it is easy to check that the forgetful functor preserves filtered colimits (2). Recall from A.3 that the coproduct $\mathcal{P} \vee \mathcal{F}(D)$ admits a basis composed by (connected) graphs with vertices indexed elements of $\mathcal{P}$ and D such that there is no pair of adjacent vertices indexed by two elements of $\mathcal{P}$. Therefore, $\mathcal{P} \vee \mathcal{F}(D)$ is equal to the directed sum $\mathcal{P} \oplus X$, where X has a basis given by graphs indexed by elements coming from $\mathcal{P}$ and at least one element from D. The map $\mathcal{P} \to \mathcal{P} \vee \mathcal{F}(D)$ is the inclusion of $\mathcal{P}$ into the first summand so that it is enough to prove that X is an acyclic chain complex. For every graph g indexed by elements of $\mathcal{P}$ and at least one element of D, the resulting chain complex is isomorphic to a quotient by the action of some symmetric groups of tensor products of $\mathcal{P}$ and at least one D. Since D is an acyclic chain complex made of free $\mathbb{K}[\mathbb{S}^{\mathrm{op}}_m \times \mathbb{S}_n]$-modules, it is an acyclic projective chain complex over any ring of symmetric subgroup . Hence the chain complex associated to any graph g indexed by elements of $\mathcal{P}$ and at least one element of D is acyclic, which proves hypothesis (3) of Transfer Theorem 89. ☐

A.5. **Cofibrations and Cofibrant objects.** In this section, we make explicit the cofibrations and the cofibrant objects in the model category of dg prop(erad)s. We refer to the Appendix of [Fre04] for the case of operads.

Proposition 94. *A map* $f : \mathcal{P} \rightarrowtail \mathcal{Q}$ *is a cofibration in the model category of dg prop(erad)s if and only if it is a retract of a map* $\mathcal{P} \to \mathcal{P} \vee \mathcal{F}(S)$, *with isomorphisms on domains, where S is a dg $\mathbb{S}$-bimodule whose components are free $\mathbb{S}$-bimodules, endowed with an exhaustive filtration*

$$S_0 = \{0\} \subset S_1 \subset S_2 \subset \cdots \subset Colim_i S_i = S$$

such that $d : S_i \to \mathcal{F}(S_{i-1})$ *and such that* $S_{i-1} \rightarrowtail S_i$ *are split monomorphisms of dg $\mathbb{S}$-bimodules with cokernels isomorphic to a free $\mathbb{S}$-bimodule.*

A map $f : \mathcal{P} \overset{\sim}{\rightarrowtail} \mathcal{Q}$ *is an acyclic cofibration in the model category of dg prop(erad)s if and only if it is a retract of a map* $\mathcal{P} \to \mathcal{P} \vee \mathcal{F}(D)$*, with isomorphisms on domains, where* $D = \bigoplus_{d \geq 1} D_i$ *is an acyclic dg* $\mathbb{S}$*-bimodule whose components are free* $\mathbb{S}$*-bimodules with each* D_i *equal to a direct sum of dg* $\mathbb{S}$*-bimodules* $D^k_{m,n}$*.*

Proof. The proposition follows from general results on the (acyclic) cofibrations of cofibrantly generated model categories. Explicitely, we apply Proposition 2.1.18 of [Hov99] to the cofibrantly generated model category of prop(erad)s. This proposition gives explicitly that (acyclic) cofibrations of prop(erad)s are retracts of relative $\mathcal{F}(I)$-cell complexes (relative $\mathcal{F}(J)$-cell complexes). We conclude by Lemma 92. ☐

Applied to $\mathcal{P} = I$, this proposition gives to following corollary.

Proposition 95. *A dg prop(erad) is cofibrant for this model category structure if and only if it is a retract of a quasi-free prop(erad)* $\mathcal{F}(S)$*, where the components of* S *are free* $\mathbb{S}$*-bimodules, endowed with an exhaustive filtration*

$$S_0 = \{0\} \subset S_1 \subset S_2 \subset \cdots \subset Colim_i S_i = S$$

such that $d : S_i \to \mathcal{F}(S_{i-1})$ *and such that* $S_{i-1} \rightarrowtail S_i$ *are split monomorphisms of dg* $\mathbb{S}$*-bimodules with cokernels isomorphic to a free* $\mathbb{S}$*-bimodule.*

Remark. In the model category of dg prop(erads) on non-negatively graded dg $\mathbb{S}$-bimodules, a dg prop(erad) is cofibrant if and only if it is retract of a quasi-free prop(erad) $\mathcal{F}(S)$ whose components are free $\mathbb{S}$-bimodules. The extra assumption on the filtration is automatically given by the homological degree.

Recall that we are working over a field of characteristic 0.

Lemma 96. *Any quasi-free prop(erad)* $\mathcal{F}(X)$ *is a retract of a quasi-free prop(erad)* $\mathcal{F}(S)$*, where the components of* S *are free* $\mathbb{S}$*-bimodules. Moreover, if* X *is endowed with an exhaustive filtration*

$$X_0 = \{0\} \subset X_1 \subset X_2 \subset \cdots \subset Colim_i X_i = X$$

such that $d : X_i \to \mathcal{F}(X_{i-1})$ *and such that* $X_{i-1} \rightarrowtail X_i$ *are split monomorphisms of dg* $\mathbb{S}$*-bimodules, then* S *can be chosen with the same property and such that the cokernels of the* $S_{i-1} \rightarrowtail S_i$ *are free* $\mathbb{S}$*-bimodules.*

Proof. Let $\bar{X}(m,n)$ denote the set of equivalence classes under the action of $\mathbb{S}^{\mathrm{op}}_m \times \mathbb{S}_n$. For simplicity, we use the generic notation $\bar{X}$. We choose a set of representatives $\{\bar{x}_i\}_{i \in \mathcal{I}}$ of $\bar{X}$. Let S be the free $\mathbb{S}$-bimodule generated by the $\{\bar{x}_i\}_{i \in \mathcal{I}}$. The generator associated to $\bar{x}_i$ will be denoted by s_i. For any x in X, we consider the sub-group $\mathbb{S}_x := \{\sigma \in \mathbb{S}^{\mathrm{op}}_m \times \mathbb{S}_n \,|\, x.\sigma = \chi(\sigma)x,\ \chi(\sigma) \in \mathbb{K}\}$. In this case, χ is a character of $\mathbb{S}_x$. We define the following element of S :

$$N(\bar{x}_i) := \frac{1}{|\mathbb{S}_{\bar{x}_i}|} \sum \chi(\sigma^{-1}).s_i\sigma,$$

where the sum runs over $\sigma \in \mathbb{S}_{\bar{x}_i}$. The image under the boundary map ∂ of an $\bar{x}_i$ is a sum of graphs of the form $\sum \mathcal{G}(\bar{x}_{i_1}, \ldots, \bar{x}_{i_k})$. We define the boundary map ∂' on $\mathcal{F}(S)$ by

$$\partial'(s_i) := \sum \frac{1}{|\mathbb{S}_{\bar{x}_i}|} \sum \chi(\sigma^{-1}).\mathcal{G}(N(\bar{x}_{i_1}), \ldots, N(\bar{x}_{i_k}))\sigma,$$

where the second sum runs over $\sigma \in \mathbb{S}_{\bar{x}_i}$. Finally, we define the maps of dg prop(erad)s $\mathcal{F}(S) \to \mathcal{F}(X)$ by $s_i \mapsto \bar{x}_i$ and $\mathcal{F}(X) \to \mathcal{F}(S)$ by $\bar{x}_i \mapsto N(\bar{x}_i)$. They form a deformation retract, which preserves the filtration of X when it exists. ☐

Corollary 97. *In the model category of dg prop(erads), any quasi-free properad* $\mathcal{F}(X)$*, where* X *is endowed with an exhaustive filtration*

$$X_0 = \{0\} \subset X_1 \subset X_2 \subset \cdots \subset Colim_i X_i = X$$

such that $d : X_i \to \mathcal{F}(X_{i-1})$ *and such that the* $X_{i-1} \rightarrowtail X_i$ *are split monomorphisms of dg* $\mathbb{S}$*-bimodules is cofibrant.*

Remark. In the non-negatively graded case, any quasi-free prop(erad) is cofibrant.

Proof. It is a direct corollary of Proposition 95 and Lemma 96. □

Theorem 98. *Any dg properad* $\mathcal{Q}$ *admits a cofibrant replacement of the form* $\mathcal{F}(S) \overset{\sim}{\twoheadrightarrow} \mathcal{Q}$ *, where the components of* S *are free* $\mathbb{S}$*-bimodules, endowed with an exhaustive filtration*

$$S_0 = \{0\} \subset S_1 \subset S_2 \subset \cdots \subset Colim_i S_i = S$$

such that $d : S_i \to \mathcal{F}(S_{i-1})$ *and such that* $S_{i-1} \rightarrowtail S_i$ *are split monomorphisms of dg* $\mathbb{S}$*-bimodules with cokernels isomorphic to a free* $\mathbb{S}$*-bimodule.*

Proof. Any dg properad $\mathcal{Q}$ admits a cofibrant replacement $I \rightarrowtail \mathcal{P} \overset{\sim}{\twoheadrightarrow} \mathcal{Q}$. Since $\mathcal{P}$ is cofibrant, it is retract $\mathcal{P} \xrightarrow{\sim} \mathcal{F}(S) \overset{\sim}{\twoheadrightarrow} \mathcal{P}$ of such an $\mathcal{F}(S)$ by Proposition 95. □

We can simply such a cofibrant replacement as follows.

Theorem 99. *A quasi-free cofibrant replacement* $\mathcal{F}(S) \overset{\sim}{\twoheadrightarrow} \mathcal{Q}$ *induces a quasi-free cofibrant replacement* $\mathcal{F}(X) \overset{\sim}{\twoheadrightarrow} \mathcal{Q}$ *, where the action of the symmetric groups on the components of* X *is the same then the action on their image in* $\mathcal{Q}$*. Moreover,* X *is endowed with an exhaustive filtration*

$$X_0 = \{0\} \subset X_1 \subset X_2 \subset \cdots \subset Colim_i X_i = X$$

such that $d : X_i \to \mathcal{F}(X_{i-1})$ *and such that* $X_{i-1} \rightarrowtail X_i$ *are split monomorphisms of dg* $\mathbb{S}$*-bimodules.*

Proof. The dg $\mathbb{S}$-bimodule which generates the quasi-free cofibrant replacement $\mathcal{F}(S) \overset{\sim}{\twoheadrightarrow} \mathcal{Q}$ is a free $\mathbb{S}$-bimodule. Let us denote by s_α the generators and q_α their image in $\mathcal{Q}$. We define X to be the $\mathbb{S}$-bimodule generated by the q_α and we consider the free properad $\mathcal{F}(X)$ on X. In $\mathcal{F}(S)$, the image of s_α under the differential map d is equal to $d(s_\alpha) = \sum \mathcal{G}(s_{\alpha_1}, \ldots, s_{\alpha_k})$. We define the differential map of $\mathcal{F}(X)$ by $d(q_\alpha) = \sum \mathcal{G}(q_{\alpha_1}, \ldots, q_{\alpha_k})$. The map $\mathcal{F}(S) \overset{\sim}{\twoheadrightarrow} \mathcal{Q}$ factors through $\mathcal{F}(S) \overset{\sim}{\twoheadrightarrow} \mathcal{F}(X) \overset{\sim}{\twoheadrightarrow} \mathcal{Q}$. Finally, $\mathcal{F}(X)$ is cofibrant by Corollary 97. □

The difference between resolution $\mathcal{F}(S)$ and $\mathcal{F}(X)$ is that in $\mathcal{F}(S)$, the symmetry of the operations of $\mathcal{Q}$ is deformed up to homotopy whereas in $\mathcal{F}(X)$ only the relations are deformed up to homotopy. (The same phenomenon appears for resolutions of the operad Com of commutative algebras where the former corresponds to E_∞ operads and the later to C_∞).
We can now choose to work with such cofibrant models. The extra filtration on the space of generators, which appears conceptually here, is similar to the one used by Sullivan [Sul77] in rational homotopy theory and by Markl in [Mar96b] for operads.

Let $\mathcal{P}$ be a dg properad. Its space of *indecomposable elements* is the cokernel of the composite map with non-trivial elements, $\mu : \bar{\mathcal{P}} \boxtimes \bar{\mathcal{P}} \to \bar{\mathcal{P}}$. The space of indecomposable elements inherits a differential map from the one of $\mathcal{P}$ which makes it into a dg $\mathbb{S}$-bimodule. The associated functor Indec : dg properads $\to$ dg$\mathbb{S}$-bimodules is left adjoint to the augmentation functor $M \mapsto M \oplus I$, where the properad structure on $M \oplus I$ is the trivial one.
The following last result will allow us to proof that the deformation complex defined in ... does not depend on the quasi-free model chosen to make it explicit.

Proposition 100. *Any weak equivalence (quasi-isomorphism) between two quasi-free cofibrant dg properads* $\mathcal{F}(X) \xrightarrow{\sim} \mathcal{F}(Y)$ *induces a weak equivalence (quasi-isomorphism) between the spaces of indecomposable elements* $X \xrightarrow{\sim} Y$*.*

Proof. The two categories of dg properads and dg $\mathbb{S}$-bimodules have model categories structures. Since the augmentation functor preserves fibrations and acyclic fibrations, by Lemma 1.3.4 of [Hov99] the indecomposable functors, being its left adjoint, preserves cofibrations and acyclic cofibrations. And by Brown's Lemma (Lemma 1.1.12 of [Hov99]), it preserves weak equivalences between cofibrant objects. □

Acknowledgements

This work was partially supported by the Göran Gustafsson foundation. The first author expresses his thanks to the Max Planck Institute in Mathematics in Bonn and École Normale Supérieure in Paris for stimulating conditions during the work on this project. The second author would like to thank Jean-Louis Loday, Dennis Sullivan, Benoit Fresse, Clemens Berger, Bernhard Keller and Domenico Fiorenza for useful discussions. The second author expresses his thanks to Northwestern University for the invitation in Spring quarter 2007. B.V. is supported by the ANR OBTH.

References

[Ada78] John Frank Adams, *Infinite loop spaces*, Annals of Mathematics Studies, vol. 90, Princeton University Press, Princeton, N.J., 1978.

[Agu00] M. Aguiar, *Infinitesimal Hopf algebras*, New trends in Hopf algebra theory (La Falda, 1999), Contemp. Math., vol. 267, Amer. Math. Soc., Providence, RI, 2000, pp. 1–29.

[AL07] Marcelo Aguiar and Muriel Livernet, *The associative operad and the weak order on the symmetric groups*, J. Homotopy Relat. Struct. **2** (2007), no. 1, 57–84 (electronic).

[And74] M. André, *Homologie des algèbres commutatives*, Springer-Verlag, Berlin, 1974, Die Grundlehren des mathematischen Wissenschaften, Band 206.

[Bal97] D. Balavoine, *Deformations of algebras over a quadratic operad*, Operads: Proceedings of Renaissance Conferences (Hartford, CT/Luminy, 1995) (Providence, RI), Contemp. Math., vol. 202, Amer. Math. Soc., 1997, pp. 207–234.

[Bar06] S. Barannikov, *Modular operads and BV geometry*, MPIM preprint (2006).

[Bar07] V. Baranovsky, *A universal enveloping for L_∞-algebras*, `arXiv:0706.1396` (2007).

[BF04] Clemens Berger and Benoit Fresse, *Combinatorial operad actions on cochains*, Math. Proc. Cambridge Philos. Soc. **137** (2004), no. 1, 135–174.

[BM03] C. Berger and I. Moerdijk, *Axiomatic homotopy theory for operads*, Comment. Math. Helv. **78** (2003), no. 4, 805–831.

[BM06] D. Borisov and Yu I. Manin, *Generalized operads and their inner cohomomorphisms*, `arXiv:math.CT/0609748` (2006).

[Bro59] E. H. Brown, Jr., *Twisted tensor products. I*, Ann. of Math. (2) **69** (1959), 223–246.

[CG03] N. Ciccoli and L Guerra, *The variety of Lie bialgebras*, J. Lie Theory **13** (2003), no. 2, 579–590.

[Con85] A. Connes, *Noncommutative differential geometry*, Inst. Hautes Études Sci. Publ. Math. (1985), no. 62, 257–360.

[Cra95] S. E. Crans, *Quillen closed model structures for sheaves*, J. Pure Appl. Algebra **101** (1995).

[Dol07] V. A. Dolgushev, *Erratum to: "A Proof of Tsygan's Formality Conjecture for an Arbitrary Smooth Manifold"*, `arXiv:math.QA/0703113` (2007).

[EE05] Benjamin Enriquez and Pavel Etingof, *On the invertibility of quantization functors*, J. Algebra **289** (2005), no. 2, 321–345.

[Fre04] B. Fresse, *Koszul duality of operads and homology of partition posets*, Homotopy theory: relations with algebraic geometry, group cohomology, and algebraic K-theory, Contemp. Math., vol. 346, Amer. Math. Soc., Providence, RI, 2004, pp. 115–215.

[Gan03] W. L. Gan, *Koszul duality for dioperads*, Math. Res. Lett. **10** (2003), no. 1, 109–124.

[Gan05] ———, *On a theorem of Conant-Vogtmann*, Math. Res. Lett. **12** (2005), no. 5-6, 749–757.

[Ger63] M. Gerstenhaber, *The cohomology structure of an associative ring*, Ann. of Math. (2) **78** (1963), 267–288.

[GJ94] E. Getzler and J. D. S. Jones, *Operads, homotopy algebra and iterated integrals for double loop spaces*, `hep-th/9403055` (1994).

[GK98] E. Getzler and M. M. Kapranov, *Modular operads*, Compositio Math. **110** (1998), no. 1, 65–126.

[GO04] D. Guin and J.-M. Oudom, *On the lie envelopping algebra of a pre-lie algebra*, `arXiv.org:math/0404457` (2004).

[Gra07] Johan Granåker, *Strong homotopy properads*, Int. Math. Res. Not. IMRN (2007), no. 14.

[GS90] M. Gerstenhaber and S.D. Schack, *Bialgebra cohomology, deformations, and quantum groups*, Proc. Nat. Acad. Sci. USA **87** (1990), 478–481.

[GV95] M. Gerstenhaber and A. Voronov, *Homotopy G-algebras and moduli space operad*, Internat. Math. Res. Notices (1995), no. 3, 141–153 (electronic).

[Hin97] V. Hinich, *Homological algebra of homotopy algebras*, Comm. Algebra **25** (1997), no. 10.

[Hin03] V. Hinich, *Erratum to "homological algebra of homotopy algebras"*, `arXiv:math.QA/0309453` (2003).

[HKV06] Po Hu, Igor Kriz, and Alexander A. Voronov, *On Kontsevich's Hochschild cohomology conjecture*, Compos. Math. **142** (2006), no. 1, 143–168.

[Hov99] M. Hovey, *Model categories*, Mathematical Surveys and Monographs, vol. 63, American Mathematical Society, Providence, RI, 1999.

[Ill71] L. Illusie, *Complexe cotangent et déformations. I II*, Springer-Verlag, Berlin, 1971, Lecture Notes in Mathematics, Vol. 239.

[Kar87] M. Karoubi, *Homologie cyclique et K-théorie*, Astérisque (1987), no. 149, 147.

[Kau07] R.M. Kaufmann, *On spineless cacti, deligne's conjecture and connes-kreimer's hopf algebra*, Topology **46** (2007), no. 1, 39.

[KM01] M. Kapranov and Yu. Manin, *Modules and Morita theorem for operads*, Amer. J. Math. **123** (2001), no. 5, 811–838.

[Kon99] M. Kontsevich, *Operads and motives in deformation quantization*, Lett. Math. Phys. **48** (1999), no. 1, 35–72, Moshé Flato (1937–1998).

[Kon03] ———, *Deformation quantization of Poisson manifolds*, Lett. Math. Phys. **66** (2003), no. 3, 157–216.

[KS91] Y. Kosmann-Schwarzbach, *Grand crochet, crochets de Schouten et cohomologies d'algèbres de Lie*, C. R. Acad. Sci. Paris Sér. I Math. **312** (1991), no. 1, 123–126.

[KS00] M. Kontsevich and Y. Soibelman, *Deformations of algebras over operads and the Deligne conjecture*, Conférence Moshé Flato 1999, Vol. I (Dijon), Math. Phys. Stud., vol. 21, Kluwer Acad. Publ., Dordrecht, 2000, pp. 255–307.

[LFCG01] J.-L. Loday, A. Frabetti, F. Chapoton, and F. Goichot, *Dialgebras and related operads*, Lecture Notes in Mathematics, vol. 1763, Springer-Verlag, Berlin, 2001.

[LM91] A. Yu. Lazarev and M. V. Movshev, *Deformations of Hopf algebras*, Uspekhi Mat. Nauk **46** (1991), no. 1(277), 211–212.

[LM05] T. Lada and M. Markl, *Symmetric brace algebras*, Appl. Categ. Structures **13** (2005), no. 4, 351–370.

[Lod98] J.-L. Loday, *Cyclic homology*, second ed., Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 301, Springer-Verlag, Berlin, 1998, Appendix E by María O. Ronco, Chapter 13 by the author in collaboration with Teimuraz Priashvili.

[Lod06] J.-L. Loday, *Generalized bialgebras and triples of operads*, `arXiv:math.QA/0611885` (2006).

[LR06] J.-L. Loday and M. Ronco, *On the structure of cofree Hopf algebras*, J. Reine Angew. Math. **592** (2006), 123.

[Mar96a] M. Markl, *Cotangent cohomology of a category and deformations*, J. Pure Appl. Algebra **113** (1996), no. 2, 195–218.

[Mar96b] Martin Markl, *Models for operads*, Comm. Algebra **24** (1996), no. 4, 1471–1500.

[Mar04a] M. Markl, *Homotopy algebras are homotopy algebras*, Forum Math. **16** (2004), no. 1, 129–160.

[Mar04b] ———, *Intrinsic brackets and the l_∞-deformation theory of bialgebras*, `arXiv.org:math/0411456` (2004).

[Mar06] ———, *A resolution (minimal model) of the PROP for bialgebras*, J. Pure Appl. Algebra **205** (2006), no. 2, 341–374.

[Mer00] S. A. Merkulov, *An L_∞-algebra of an unobstructed deformation functor*, Internat. Math. Res. Notices (2000), no. 3, 147–164.

[Mer04] ———, *Prop profile of deformation quantization and graph complexes with loops and wheels*, 2004.

[Mer05] ———, *Nijenhuis infinity and contractible dg manifolds, math.ag/0403244*, Compositio Mathematica (2005), no. 141, 1238–1254.

[Mer06] ———, *Prop profile of poisson geometry, math.dg/0401034*, Commun. Math. Phys. (2006), no. 262, 117–135.

[ML65] Saunders Mac Lane, *Categorical algebra*, Bull. Amer. Math. Soc. **71** (1965), 40–106.

[ML98] S. Mac Lane, *Categories for the working mathematician*, second ed., Graduate Texts in Mathematics, vol. 5, Springer-Verlag, New York, 1998.

[MMS06] M. Markl, S. Merkulov, and S. Shadrin, *Wheeled PROPs, graph complexes and the master equation*, `arXiv:math.AG/0610683` (2006).

[Moo71] John C. Moore, *Differential homological algebra*, Actes du Congrès International des Mathématiciens (Nice, 1970), Tome 1, Gauthier-Villars, Paris, 1971, pp. 335–339.

[MS02] J. E. McClure and J. H. Smith, *A solution of Deligne's Hochschild cohomology conjecture*, Recent progress in homotopy theory (Baltimore, MD, 2000), Contemp. Math., vol. 293, Amer. Math. Soc., Providence, RI, 2002, pp. 153–193.

[MSS02] M. Markl, S. Shnider, and J. Stasheff, *Operads in algebra, topology and physics*, Mathematical Surveys and Monographs, vol. 96, American Mathematical Society, Providence, RI, 2002.

[MV03] M. Markl and A. Voronov, *PROPped up graph cohomology*, `arXiv:math.QA/0307081` (2003).

[NR67] A. Nijenhuis and R. W. Richardson, Jr., *Deformations of Lie algebra structures*, J. Math. Mech. **17** (1967), 89–105.

[Qui67] D. G. Quillen, *Homotopical algebra*, Lecture Notes in Mathematics, No. 43, Springer-Verlag, Berlin, 1967.

[Qui69] D. Quillen, *Rational homotopy theory*, Ann. of Math. (2) **90** (1969), 205–295.

[Qui70] ———, *On the (co-) homology of commutative rings*, Applications of Categorical Algebra (Proc. Sympos. Pure Math., Vol. XVII, New York, 1968), Amer. Math. Soc., Providence, R.I., 1970, pp. 65–87.

[Sho03] B. Shoikhet, *On the Cambell-Baker-Hausdorf deformation quatization of a linear Poisson structure*, `arXiv:math/9903036` (2003).

[Spi01] M. Spitzweck, *Operads, algebras and modules in general model categories*, `arXiv:math.AT/0101102` (2001).

- [Sta63] J. D. Stasheff, *Homotopy associativity of H-spaces. I, II*, Trans. Amer. Math. Soc. 108 (1963), 275-292; ibid. **108** (1963), 293–312.
- [Sta93] J. Stasheff, *The intrinsic bracket on the deformation complex of an associative algebra*, J. Pure Appl. Algebra **89** (1993), no. 1-2, 231–235.
- [Sul77] Dennis Sullivan, *Infinitesimal computations in topology*, Inst. Hautes Études Sci. Publ. Math. (1977), no. 47, 269–331 (1978).
- [Tam98] D. E. Tamarkin, *Another proof of M. Kontsevich formality theorem*, `arXiv:math/9803025` (1998).
- [Tam00] ———, *Deformation complex of a d-algebra is a (d+1)-algebra*, `arXiv:math/0010072` (2000).
- [Tou04] V. Tourtchine, *On the other side of the bialgebra of chord diagrams*, `math.QA/0411436` (2004).
- [Val04] B. Vallette, *Free monoid in monoidal abelian categories*, `arXiv:math/0411543`, to appear in Applied Categorical Structures (2004).
- [Val06a] ———, *Homology of generalized partition posets*, J. Pure Appl. Algebra **208** (2006), no. 2, 699–725.
- [Val06b] ———, *Manin products, Koszul duality, Loday algebras and Deligne conjecture*, `arXiv:math/0609002`, to appear in J. Reine Angew. Math. (2006).
- [Val07a] ———, *A Koszul duality for props*, Trans. of Amer. Math. Soc. **359** (2007), 4865–4993.
- [Val07b] ———, *Transfer of algebraic homotopy structures*, in preparation (2007).
- [VdL02] P. Van der Laan, *Operads up to Homotopy and Deformations of Operad Maps*, `arXiv:math.QA/0208041` (2002).
- [VdL03] ———, *Coloured Koszul duality and strongly homotopy operads*, `arXiv:math.QA/0312147` (2003).
- [Vor00] Alexander A. Voronov, *Homotopy Gerstenhaber algebras*, Conférence Moshé Flato 1999, Vol. II (Dijon), Math. Phys. Stud., vol. 22, Kluwer Acad. Publ., Dordrecht, 2000, pp. 307–331.
- [Wei94] C. A. Weibel, *An introduction to homological algebra*, Cambridge Studies in Advanced Mathematics, vol. 38, Cambridge University Press, Cambridge, 1994.

SERGEI MERKULOV, DEPARTMENT OF MATHEMATICS, STOCKHOLM UNIVERSITY,
10691 STOCKHOLM, SWEDEN
E-mail address : `sm@math.su.se`
URL : `http://www.math.su.se/∼sm/`

BRUNO VALLETTE, LABORATOIRE J.A. DIEUDONNÉ, UNIVERSITÉ DE NICE SOPHIA-ANTIPOLIS,
PARC VALROSE, 06108 NICE CEDEX 02, FRANCE
E-mail address : `brunov@math.unice.fr`
URL : `http://math.unice.fr/∼brunov`